

Brightside Specialty Pharmacy

431 N. Tustin Ave., STE C, Santa Ana, CA 92705

Phone 714-707-5115

Date :8/10/2022

Patient Education for

PREGABALIN 75 MG CAPSULE

Check with doctor before discontinuing Avoid drinking alcohol Risk of angioedema, anaphylaxis, or allergy. Tell Pt to report STAT Avoid taking with other CNS depressant drugs or alcohol Dispense Medication Guide and explain to patient before use May cause drowsiness or dizziness Use caution driving. To maintain constant blood levels, space doses evenly throughout day. Depression or thoughts of suicide should be reported to MD ASAP.

Do not stop medicine without calling doctor Avoid drinking alcohol Call Dr now if swelling, rash or breathing problem Avoid alcohol/other drugs that make you sleepy Before use, important to read the Medication Guide May make you drowsy or dizzy. Drive with caution Space doses evenly throughout the day. Immediately report to MD any thoughts of suicide.

.....Education Section.....

IMPORTANT:

HOW TO USE THIS INFORMATION: This is a summary and

does NOT have all possible information about this product. This information does not assure that this product is safe, effective, or appropriate for you. This information is not individual medical advice and does not substitute for the advice of your health care professional. Always ask your health care professional for complete information about this product and your specific health needs.

PREGABALIN - ORAL

(pree-GAH-ba-lin)

COMMON BRAND NAME(S):

Lyrica

USES:

This medication is used to treat pain caused by nerve

damage due to diabetes, shingles (herpes zoster) infection, or spinal cord injury. This medication is also used to treat pain in people with fibromyalgia.

It is also used with other medications to treat certain types of seizures (focal seizures).

OTHER USES:

This section contains uses of this drug that are not

listed in the approved professional labeling for the drug but that may be prescribed by your health care professional. Use this drug for a condition that is listed in this section only if it has been so prescribed by your health care professional.

This drug may also be used for hot flashes that occur with menopause, restless legs syndrome (RLS), or anxiety.

HOW TO USE:

Read the Medication Guide and, if available, the

Patient Information Leaflet provided by your pharmacist before you start using pregabalin and each time you get a refill. If you have any questions regarding the information, consult your doctor or pharmacist.

Take this medication by mouth as directed by your doctor,

usually 2 to 3 times a day with or without food. The dosage is based on your medical condition, kidney function, and response to treatment. Children's dosage is also based on weight.

If you are using the liquid form of this medication, carefully measure the dose using a special measuring device/spoon. Do not use a household spoon because you may not get the correct dose.

To reduce your risk of side effects (such as dizziness and drowsiness), your doctor may direct you to start this medication at a low dose and gradually increase your dose. Follow your doctor's instructions carefully.

Use this medication regularly to get the most benefit from it. To help you remember, take it at the same times each day. This drug works best when the amount of medicine in your body is kept at a constant level. It is best to take pregabalin at evenly spaced intervals throughout the day and night.

Do not stop taking this medication without consulting your doctor. Some conditions (such as seizures) may become worse when this drug is suddenly stopped. Also, you may experience symptoms such as difficulty sleeping, nausea, headache and diarrhea. To prevent these symptoms while you are stopping treatment with this drug, your doctor may reduce your dose gradually. Report any new or worsening symptoms right away.

Though it helps many people, this medication may sometimes cause addiction. This risk may be higher if you have a substance use disorder (such as overuse of or addiction to drugs/alcohol). Take this medication exactly as prescribed to lower the risk of addiction. Ask your doctor or pharmacist for more details.

Tell your doctor if your condition lasts or gets worse.

SIDE EFFECTS:

Drowsiness, dizziness, dry mouth, constipation,

difficulty concentrating, or weight gain may occur. If any of these effects last or get worse, tell your doctor or pharmacist promptly.

Remember that this medication has been prescribed because your doctor has judged that the benefit to you is greater than the risk of side effects. Many people using this medication do not have serious side effects.

Tell your doctor right away if you have any unlikely but serious side effects, including: blurred vision, unusual

bleeding/bruising, unsteadiness, confusion, muscle pain/tenderness/weakness (especially if you are tired or have a fever), swelling of hands/legs/feet, signs of kidney problems (such as change in the amount of urine).

A small number of people who take anticonvulsants for any condition (such as seizure, bipolar disorder, pain) may experience depression, suicidal thoughts/attempts, or other mental/mood problems. Tell your doctor right away if you or your family/caregiver notice any unusual/sudden changes in your mood, thoughts, or behavior including signs of depression, suicidal thoughts/attempts, thoughts about harming yourself.

Get medical help right away if you have any very serious side effects, including: slow/shallow breathing.

A very serious allergic reaction to this drug is rare. However, get medical help right away if you notice any symptoms of a serious allergic reaction, including: rash, itching/swelling (especially of the face/tongue/throat), severe dizziness, trouble breathing.

This is not a complete list of possible side effects. If you notice other effects not listed above, contact your doctor or pharmacist.

In the US -

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or at www.fda.gov/medwatch.

In Canada - Call your doctor for medical advice about side effects. You may report side effects to Health Canada at 1-866-234-2345.

PRECAUTIONS:

Before taking pregabalin, tell your doctor or

pharmacist if you are allergic to it; or if you have any other allergies. This product may contain inactive ingredients, which can cause allergic reactions or other problems. Talk to your pharmacist for more details.

Before using this medication, tell your doctor or pharmacist your medical history, especially of: heart problems (such as heart failure), kidney disease, history of an allergic reaction which included itching/swelling of the face/lips/tongue/throat (angioedema), personal or family history of a substance use disorder (such as overuse of or addiction to drugs/alcohol), breathing problems (such as COPD).

This drug may make you dizzy or drowsy or blur your vision. Alcohol or marijuana (cannabis) can make you more dizzy or drowsy. Do not drive, use machinery, or do anything that needs alertness or clear vision until you can do it safely. Avoid alcoholic beverages. Talk to your doctor if you are using marijuana (cannabis).

Before having surgery, tell your doctor or dentist about all the products you use (including prescription drugs, nonprescription drugs, and herbal products).

Older adults may be more sensitive to the side effects of this drug, especially drowsiness, dizziness, unsteadiness, confusion, and slow/shallow breathing.

During pregnancy, this medication should be used only when clearly needed. Discuss the risks and benefits with your doctor.

This drug passes into breast milk. Consult your doctor before

breast-feeding.

DRUG INTERACTIONS:

Drug interactions may change how your

medications work or increase your risk for serious side effects. This document does not contain all possible drug interactions. Keep a list of all the products you use (including prescription/nonprescription drugs and herbal products) and share it with your doctor and pharmacist. Do not start, stop, or change the dosage of any medicines without your doctor's approval.

A product that may interact with this drug is: orlistat.

The risk of serious side effects (such as slow/shallow breathing, severe drowsiness/dizziness) may be increased if this medication is taken with other products that may also cause drowsiness or breathing problems. Tell your doctor or pharmacist if you are using other products such as opioid pain or cough relievers (such as codeine, hydrocodone), alcohol, marijuana (cannabis), drugs for sleep or anxiety (such as alprazolam, lorazepam, zolpidem), muscle relaxants (such as carisoprodol, cyclobenzaprine), or antihistamines (such as cetirizine, diphenhydramine).

Check the labels on all your medicines (such as allergy or cough-and-cold products) because they may contain ingredients that cause drowsiness. Ask your pharmacist about using those products safely.

OVERDOSE:

If someone has overdosed and has serious symptoms such

as passing out or trouble breathing, call 911. Otherwise, call a poison control center right away. US residents can call their local poison control center at 1-800-222-1222. Canada residents can call a provincial poison control center.

NOTES:

Do not share this medication with others. Sharing it is

against the law.

Laboratory tests may be done to measure kidney function.

MISSED DOSE:

If you miss a dose, use it as soon as you remember.

If it is near the time of the next dose, skip the missed dose.

Use your next dose at the regular time. Do not double the dose to catch up.

STORAGE:

Store at room temperature away from light and moisture.

Do not store in the bathroom. Keep all medications away from children and pets.

Do not flush medications down the toilet or pour them into a drain unless instructed to do so. Properly discard this product when it is expired or no longer needed. Consult your pharmacist or local waste disposal company.

MEDICAL ALERT:

Your condition can cause complications in a

medical emergency. For information about enrolling in MedicAlert, call 1-888-633-4298 (US) or 1-800-668-1507 (Canada).

