

Brightside Specialty Pharmacy

431 N. Tustin Ave., STE C, Santa Ana, CA 92705

Phone 714-707-5115

Date :8/10/2022

Patient Education for TRIAMCINOLONE 0.1% CREAM



Use for specified length of therapy only. Do not apply occlusive bandage unless directed. Do not use for other skin problems unless directed by doctor. Do not use on face, genitals or axilla unless directed by doctor. Following prolonged use, adrenal suppression is possible. Infections can become worse due to immune system suppression. Clean and dry skin, then apply a thin layer to affected area.

Use for specified length of therapy only. Do not bandage unless told by doctor. Use for other problems only if directed by MD. Use on face/under arms/or groin only if directed. After prolonged use, call doctor before stopping. Call doctor if condition does not improve. Clean and dry skin, then use a thin layer on area.

.....Education Section.....

IMPORTANT:

HOW TO USE THIS INFORMATION: This is a summary and

does NOT have all possible information about this product. This information does not assure that this product is safe, effective, or appropriate for you. This information is not individual medical advice and does not substitute for the advice of your health care professional. Always ask your health care professional for complete information about this product and your specific health needs.

TRIAMCINOLONE - TOPICAL

(trye-am-SIN-oh-lone)

COMMON BRAND NAME(S):

Aristocort, Aristocort HP, Kenalog, Triderm

USES:

This medication is used to treat a variety of skin

conditions (such as eczema, dermatitis, allergies, rash).

Triamcinolone reduces the swelling, itching, and redness that can occur in these types of conditions. This medication is a medium-to strong-potency corticosteroid. The potency depends on the strength and the form that you use. For details on the potency of your product, ask your pharmacist.

HOW TO USE:

Use this medication only on the skin. Do not use it

on the face, groin, or underarms unless directed to do so by your doctor.

Wash and dry your hands. Before applying the medication, clean and dry the affected area. Apply a thin film of the medication to the affected area and gently rub in, usually 2 to 4 times daily or as directed by your doctor. Do not cover, bandage or wrap the area unless directed to do so by your doctor.

After applying the medication, wash your hands unless you are using this medication to treat the hands. When applying this medication near the eyes, avoid getting it in the eyes as this may worsen or cause glaucoma. Also, avoid getting this medication in the nose or mouth. If you get the medication in your eyes, nose, or mouth, rinse with plenty of water.

Use this medication only for the condition prescribed. Do not use it for longer than prescribed.

Inform your doctor if your condition lasts or gets worse.

SIDE EFFECTS:

Burning, itching, irritation, or dryness may occur

when this medication is first applied to the skin. This should disappear in a few days as your body adjusts to the medication. If any of these effects last or get worse, notify your doctor or pharmacist promptly.

Remember that this medication has been prescribed because your doctor has judged that the benefit to you is greater than the risk of side effects. Many people using this medication do not have serious side effects.

Tell your doctor right away if you have any serious side effects, including: stretch marks, skin thinning/discoloration, acne, greatly increased hair growth, "hair bumps" (folliculitis).

Skin infections can become worse when this medication is used. Notify your doctor if redness, swelling, or irritation do not improve.

Rarely, it is possible this medication will be absorbed from the skin into the bloodstream. This can lead to side effects of too much corticosteroid. These side effects are more likely in children, and in people who use this medication for a long time or over large areas of the skin. Tell your doctor right away if any of the following side effects occur: unusual/extreme tiredness, weight loss, headache, swelling ankles/feet, increased thirst/urination, vision problems.

A very serious allergic reaction to this drug is rare. However, get medical help right away if you notice any symptoms of a serious allergic reaction, including: rash, itching/swelling (especially of the face/tongue/throat), severe dizziness, trouble breathing.

This is not a complete list of possible side effects. If you notice other effects not listed above, contact your doctor or pharmacist.

In the US -

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or at www.fda.gov/medwatch.

In Canada - Call your doctor for medical advice about side effects. You may report side effects to Health Canada at 1-866-234-2345.

PRECAUTIONS:

Before using triamcinolone, tell your doctor or

pharmacist if you are allergic to it; or to other corticosteroids (such as hydrocortisone, prednisone); or if you have any other allergies. This product may contain inactive ingredients, which can cause allergic reactions or other problems. Talk to your pharmacist for more details.

Before using this medication, tell your doctor or pharmacist your medical history, especially of: poor blood circulation, immune system problems.

Do not use if there is an infection or sore present in the area to be treated.

Rarely, using corticosteroid medications for a long time or over large areas of skin can make it more difficult for your body to respond to physical stress. Before having surgery or emergency treatment, or if you get a serious illness/injury, tell your doctor or dentist that you are using this medication or have used this medication within the past few months.

Though it is unlikely, this medication may temporarily slow down a child's growth if used for a long time. See the doctor regularly so your child's height can be checked.

During pregnancy, this medication should be used only when clearly needed. Discuss the risks and benefits with your doctor.

It is not known whether this drug passes into breast milk when applied to the skin. Other medications in this class pass into breast milk when taken by mouth. Consult your doctor before breast-feeding.

DRUG INTERACTIONS:

Drug interactions may change how your

medications work or increase your risk for serious side effects. This document does not contain all possible drug interactions. Keep a list of all the products you use (including prescription/nonprescription drugs and herbal products) and share it with your doctor and pharmacist. Do not start, stop, or change the dosage of any medicines without your doctor's approval.

Some products that may interact with this drug are: corticosteroids taken by mouth (such as prednisone), drugs that suppress the immune system (such as cyclosporine).

OVERDOSE:

This medicine may be harmful if swallowed. If someone

has overdosed and has serious symptoms such as passing out or trouble breathing, call 911. Otherwise, call a poison control center right away. US residents can call their local poison control center at 1-800-222-1222. Canada residents can call a provincial poison control center.

NOTES:

Do not share this medication with others.

This medication has been prescribed for your current condition only. Do not use it later for other skin problems unless told to do so by your doctor. A different medication may be necessary in

those cases.

Laboratory and/or medical tests (such as adrenal gland function tests) may be performed periodically to monitor your progress or check for side effects, especially if you use this drug for an extended period of time or apply it over large areas of the body. Consult your doctor for more details.

Inform all your doctors that you use (or have used) this medication.

MISSED DOSE:

If you miss a dose, use it as soon as you remember.

If it is near the time of the next dose, skip the missed dose.

Use your next dose at the regular time. Do not double the dose to catch up.

STORAGE:

Store at room temperature away from light and moisture.

Do not freeze. Do not store in the bathroom. Keep all medications away from children and pets.

Do not flush medications down the toilet or pour them into a drain unless instructed to do so. Properly discard this product when it is expired or no longer needed. Consult your pharmacist or local waste disposal company.

Information last revised January 2022.

Copyright(c) 2022 First Databank, Inc.