

Brightside Specialty Pharmacy

431 N. Tustin Ave., STE C, Santa Ana, CA 92705

Phone 714-707-5115

Date :8/10/2022

Patient Education for VITAMIN D3 125 MCG (5000 UNIT)

Read directions carefully before start of therapy Take as directed. Some formulations best with food, improves absorption. Bile acid sequestrant(s) interfere w/absorption (e.g. cholestyramine) Check labels on OTC medications for contraindications. Discuss with doctor your needs for calcium, vitamin D, and diet. Signs of excess vitamin D: emesis, pain, drowsiness, and constipation

Read directions carefully before start of therapy Take as directed. Ask RPh if your brand is best taken with food. If on bile resins, take drug 2 hours before/after. Check warnings on OTC medications. Discuss with MD your need for calcium and vitamin D. Call MD for constipation, vomiting/pain/drowsiness.

.....Education Section.....

IMPORTANT:

HOW TO USE THIS INFORMATION: This is a summary and

does NOT have all possible information about this product. This information does not assure that this product is safe, effective, or appropriate for you. This information is not individual medical advice and does not substitute for the advice of your health care professional. Always ask your health care professional for complete information about this product and your specific health needs.

VITAMIN D - ORAL

COMMON BRAND NAME(S):

Calciferol, Drisdol

USES:

Vitamin D (ergocalciferol-D2, cholecalciferol-D3,

alfacalcidol) is a fat-soluble vitamin that helps your body absorb calcium and phosphorus. Having the right amount of vitamin D, calcium, and phosphorus is important for building and keeping strong bones. Vitamin D is used to treat and prevent bone disorders (such as rickets, osteomalacia). Vitamin D is made by the body when skin is exposed to sunlight. Sunscreen, protective clothing, limited exposure to sunlight, dark skin, and age may prevent getting enough vitamin D from the sun.

Vitamin D with calcium is used to treat or prevent bone loss (osteoporosis). Vitamin D is also used with other medications to treat low levels of calcium or phosphate caused by certain disorders (such as hypoparathyroidism, pseudohypoparathyroidism, familial hypophosphatemia). It may be used in kidney disease to keep calcium levels normal and allow normal bone growth. Vitamin D drops (or other supplements) are given to breast-fed infants because breast milk usually has low levels of vitamin D.

HOW TO USE:

Take vitamin D by mouth as directed. Vitamin D is

best absorbed when taken after a meal but may be taken with or without food. Alfacalcidol is usually taken with food. Follow all directions on the product package. If you have any questions, ask your doctor or pharmacist.

If your doctor has prescribed this medication, take as directed by your doctor. Your dosage is based on your medical condition, amount of sun exposure, diet, age, and response to treatment.

If you are using the liquid form of this medication, carefully measure the dose using a special measuring device/spoon. Do not use a household spoon because you may not get the correct dose.

If you are taking the chewable tablet or wafers, chew the medication thoroughly before swallowing. Do not swallow whole wafers.

If you are taking the rapidly-dissolving tablets, dry your hands before handling the medication. Place each dose on the tongue, allow it to dissolve completely, and then swallow it with saliva or water. You do not need to take this medication with water.

Certain medications (bile acid sequestrants such as cholestyramine/colestipol, mineral oil, orlistat) can decrease the absorption of vitamin D. Take your doses of these medications as far as possible from your doses of vitamin D (at least 2 hours apart, longer if possible). It may be easiest to take vitamin D at bedtime if you are also taking these other medications. Ask your doctor or pharmacist how long you should wait between doses and for help finding a dosing schedule that will work with all your medications.

Take this medication regularly to get the most benefit from it. To help you remember, take it at the same time each day if you are taking it once a day. If you are taking this medication only once a week, remember to take it on the same day each week. It may help to mark your calendar with a reminder.

If your doctor has recommended that you follow a special diet (such as a diet high in calcium), it is very important to follow the diet to get the most benefit from this medication and to prevent serious side effects. Do not take other supplements/vitamins unless ordered by your doctor.

If you think you may have a serious medical problem, get medical help right away.

SIDE EFFECTS:

Vitamin D at normal doses usually has no side

effects. If you have any unusual effects, contact your doctor or

pharmacist promptly.

If your doctor has directed you to use this medication, remember that your doctor has judged that the benefit to you is greater than the risk of side effects. Many people using this medication do not have serious side effects.

Too much vitamin D can cause harmful high calcium levels. Tell your doctor right away if any of these signs of high vitamin

D/calcium levels occur:

nausea/vomiting, constipation, loss of

appetite, increased thirst, increased urination, mental/mood changes, unusual tiredness.

A very serious allergic reaction to this drug is rare.

However, get medical help right away if you notice any symptoms of a serious allergic reaction, including: rash, itching/swelling (especially of the face/tongue/throat), severe dizziness, trouble breathing.

This is not a complete list of possible side effects. If you notice other effects not listed above, contact your doctor or pharmacist.

In the US -

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or at www.fda.gov/medwatch.

In Canada - Call your doctor for medical advice about side effects. You may report side effects to Health Canada at 1-866-234-2345.

PRECAUTIONS:

Before taking vitamin D, tell your doctor or

pharmacist if you are allergic to it; or to other vitamin D products (such as calcitriol); or if you have any other allergies. This product may contain inactive ingredients (such as peanut/soy), which can cause allergic reactions or other problems. Talk to your pharmacist for more details.

Before using this medication, tell your doctor or pharmacist your medical history, especially of: high calcium/vitamin D levels (hypercalcemia/hypervitaminosis D), difficulty absorbing nutrition from food (malabsorption syndrome), kidney disease, liver disease.

Liquid products, chewable tablets, or dissolving tablets may contain sugar and/or aspartame. Liquid products may also contain alcohol. Caution is advised if you have diabetes, liver disease, phenylketonuria (PKU), or any other condition that requires you to limit/avoid these substances in your diet. Ask your doctor or pharmacist about using this product safely.

Before having surgery, tell your doctor or dentist about all the products you use (including prescription drugs, nonprescription drugs, and herbal products).

During pregnancy, doses of vitamin D greater than the recommended dietary allowance should be used only when clearly needed. Discuss the risks and benefits with your doctor.

This medication passes into breast milk. Consult your doctor before breast-feeding.

DRUG INTERACTIONS:

Drug interactions may change how your

medications work or increase your risk for serious side effects.

This document does not contain all possible drug interactions.

Keep a list of all the products you use (including prescription/nonprescription drugs and herbal products) and share it with your doctor and pharmacist. Do not start, stop, or change the dosage of any medicines without your doctor's approval.

Some products that may interact with this drug include:

burosumab, phosphate binders.

Check the labels on all your prescription and nonprescription/herbal products (such as antacids, laxatives, vitamins) because they may contain calcium, magnesium, phosphate, or vitamin D. Ask your pharmacist about using those products safely.

Vitamin D is very similar to calcitriol. Do not use medications containing calcitriol while using vitamin D.

This medication may interfere with certain laboratory tests (including cholesterol tests), possibly causing false test results. Make sure laboratory personnel and all your doctors know you use this drug.

OVERDOSE:

If someone has overdosed and has serious symptoms such

as passing out or trouble breathing, call 911. Otherwise, call a poison control center right away. US residents can call their local poison control center at 1-800-222-1222. Canada residents can call a provincial poison control center. Symptoms of overdose may include: seizures, confusion, irregular heartbeat.

NOTES:

Keep all regular medical and laboratory appointments. If

your doctor has prescribed this medication, laboratory and/or medical tests (such as calcium/magnesium/phosphorus levels) should be performed periodically to monitor your progress or check for side effects. Consult your doctor for more details.

Foods rich in vitamin D include: fortified dairy products, eggs, sardines, cod liver oil, chicken livers, and fatty fish.

MISSED DOSE:

If you miss a dose, take it as soon as you

remember. If it is near the time of the next dose, skip the missed dose. Take your next dose at the regular time. Do not double the dose to catch up.

STORAGE:

Store vitamin D products (except alfacalcidol) at room

temperature away from light and moisture. Do not store in the bathroom. Store alfacalcidol drops in the refrigerator between 36-46 degrees F (2-8 degrees C). Do not freeze. Keep all medications away from children and pets.

Do not flush medications down the toilet or pour them into a drain unless instructed to do so. Properly discard this product when it is expired or no longer needed. Consult your pharmacist or local waste disposal company for more details about how to safely discard your product.

Information last revised July 2021.

Copyright(c) 2022 First Databank, Inc.