

Brightside Specialty Pharmacy

431 N. Tustin Ave., STE C, Santa Ana, CA 92705

Phone 714-707-5115

Date :8/10/2022

Patient Education for

VYVANSE 30 MG CAPSULE

May be habit forming. Not recommended when breast feeding. Physician must know patient's complete medical history. Dispense Medication Guide and explain to patient before use. May cause dizziness. Use caution operating machinery or driving. Can induce very serious cardiovascular problems. Discuss gradual dose reduction with MD before stopping medication. Depression or thoughts of suicide should be reported to MD ASAP.

May be habit forming. Not recommended when breast feeding. Tell doctor your complete medical history. Before use, important to read the Medication Guide. May make you dizzy. Use caution when driving. Can increase heart rate and blood pressure. MD may need to reduce the dose before you stop it. Immediately report to MD any thoughts of suicide.

.....Education Section.....

IMPORTANT:

HOW TO USE THIS INFORMATION: This is a summary and

does NOT have all possible information about this product. This information does not assure that this product is safe, effective, or appropriate for you. This information is not individual medical advice and does not substitute for the advice of your health care professional. Always ask your health care professional for complete information about this product and your specific health needs.

LISDEXAMFETAMINE - ORAL

(lis-DEX-am-FET-a-meen)

COMMON BRAND NAME(S):

Vyvanse

WARNING:

Misuse or abuse of amphetamines may cause serious

(possibly fatal) heart and blood pressure problems.

Amphetamine-type medications can be habit-forming. Use only as directed. If you use this drug for a long time, you may become dependent on it and may have withdrawal symptoms after stopping the drug. Consult your doctor or pharmacist for more details.

(See also How to Use section.)

USES:

Lisdexamfetamine is used to treat attention deficit

hyperactivity disorder (ADHD) as part of a total treatment plan, including psychological, social, and other treatments. It may help to increase the ability to pay attention, stay focused, and stop fidgeting. Lisdexamfetamine may also be used to treat binge eating disorder (BED). It may help to reduce the number of binge eating days.

This medication is a stimulant. It is thought to work by restoring the balance of certain natural chemicals (neurotransmitters) in the brain.

This medication is not recommended for use for weight loss due to the risk of serious side effects.

HOW TO USE:

Read the Medication Guide provided by your

pharmacist before you start taking lisdexamfetamine and each time you get a refill. If you have any questions, ask your doctor or pharmacist.

Take this medication with or without food as directed by your doctor, usually once daily in the morning. Do not take this medication in the afternoon or evening because it may cause you to have trouble sleeping. The dosage is based on your medical condition and response to treatment. Your doctor may adjust your dose to find the dose that is best for you. Follow your doctor's instructions carefully.

If you are taking the chewable tablet, chew the tablet thoroughly and then swallow.

If you are taking the capsule form of this medication, swallow the capsule whole. However, if you have trouble swallowing the capsule, you may open the capsule and pour all of its contents (powder) in a glass of water or orange juice or mix it in yogurt. Use a spoon to break apart any powder that is stuck together. Stir well until the contents dissolve completely. Drink or eat the mixture right away. Do not prepare a supply in advance. It is normal to see a filmy coating on the inside of your glass or container after you drink or eat all of the medicine.

Use this medication regularly to get the most benefit from it. To help you remember, take it at the same time each day.

During treatment, your doctor may occasionally recommend stopping the medication for a short time to see whether there are any changes in your behavior and whether the medication is still needed.

If you suddenly stop using this medication, you may have withdrawal symptoms (such as severe tiredness, sleep problems, mental/mood changes such as depression). To help prevent withdrawal, your doctor may lower your dose slowly. Withdrawal is more likely if you have used lisdexamfetamine for a long time or in high doses. Tell your doctor or pharmacist right away if you have withdrawal.

Though it helps many people, this medication may sometimes cause addiction. This risk may be higher if you have a substance use disorder (such as overuse of or addiction to drugs/alcohol). Do not increase your dose, take it more often, or use it for a longer time than prescribed. Properly stop the medication when so directed.

When this medication is used for a long time, it may not work as well. Talk with your doctor if this medication stops working well.

Tell your doctor if your condition does not improve or if it worsens.

SIDE EFFECTS:

Nausea, vomiting, constipation, stomach/abdominal

pain, loss of appetite, dry mouth, headache, nervousness, dizziness, trouble sleeping, sweating, weight loss, irritability, and restlessness may occur. If any of these effects last or get worse, tell your doctor or pharmacist promptly.

Remember that this medication has been prescribed because your doctor has judged that the benefit to you is greater than the risk of side effects. Many people using this medication do not have serious side effects.

This medication may raise your blood pressure. Check your blood pressure regularly and tell your doctor if the results are high.

Tell your doctor right away if you have any serious side effects, including: blurred vision, fast/pounding/irregular heartbeat, mental/mood/behavior changes (such as agitation, aggression, mood swings, depression, hallucinations, abnormal thoughts/behavior, suicidal thoughts/attempts), uncontrolled movements, muscle twitching/shaking, signs of blood flow problems in the fingers or toes (such as coldness, numbness, pain, or skin color changes), unusual wounds on the fingers or toes, outbursts of words/sounds, change in sexual ability/interest, swelling ankles/feet, extreme tiredness, rapid/unexplained weight loss, frequent/prolonged erections (in males).

Get medical help right away if you have any very serious side effects, including: shortness of breath, fainting, chest/jaw/left arm pain, seizures, weakness on one side of the body, trouble speaking, confusion, sudden vision changes.

This medication may increase serotonin and rarely cause a very serious condition called serotonin syndrome/toxicity. The risk increases if you are also taking other drugs that increase serotonin, so tell your doctor or pharmacist of all the drugs you take (see Drug Interactions section). Get medical help right away if you develop some of the following symptoms: fast heartbeat, hallucinations, loss of coordination, severe dizziness, severe nausea/vomiting/diarrhea, twitching muscles, unexplained fever, unusual agitation/restlessness.

A very serious allergic reaction to this drug is rare. However, get medical help right away if you notice any symptoms of a serious allergic reaction, including: rash, itching/swelling (especially of the face/tongue/throat), severe dizziness, trouble breathing.

This is not a complete list of possible side effects. If you notice other effects not listed above, contact your doctor or pharmacist.

In the US -

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or at www.fda.gov/medwatch.

In Canada - Call your doctor for medical advice about side effects. You may report side effects to Health Canada at

1-866-234-2345.

PRECAUTIONS:

Before taking this medication, tell your doctor or

pharmacist if you are allergic to it; or to other sympathomimetic drugs (such as amphetamine or dextroamphetamine); or if you have any other allergies. This product may contain inactive ingredients, which can cause allergic reactions or other problems. Talk to your pharmacist for more details.

Before using this medication, tell your doctor or pharmacist your medical history, especially of: blood circulation problems (such as Raynaud's disease), certain mental/mood conditions (such as severe agitation, psychosis), personal/family history of mental/mood disorders (such as bipolar disorder, depression, psychotic disorder, suicidal thoughts), heart problems (including irregular heartbeat, coronary artery disease, previous heart attack, heart failure, cardiomyopathy, problems with heart structure such as valve problems), family history of heart problems (such as sudden death, irregular heartbeat), history of stroke, high blood pressure, overactive thyroid (hyperthyroidism), a certain eye problem (glaucoma), seizures, kidney disease, personal or family history of a substance use disorder (such as overuse of or addiction to drugs/alcohol), personal/family history of uncontrolled muscle movements (such as Tourette's syndrome).

This drug may make you dizzy. Alcohol or marijuana (cannabis) can make you more dizzy. Do not drive, use machinery, or do anything that needs alertness until you can do it safely. Limit alcoholic beverages. Talk to your doctor if you are using marijuana (cannabis).

Before having surgery, tell your doctor or dentist about all the products you use (including prescription drugs, nonprescription drugs, and herbal products).

Children may be more sensitive to the side effects of this drug, especially weight loss. This medication may slow down a child's growth. The doctor may recommend temporarily stopping the medication from time to time to reduce this risk. Monitor your child's weight and height. Consult your doctor or pharmacist for more details.

During pregnancy, this medication should be used only if clearly needed. Discuss the risks and benefits with your doctor. Infants born to mothers who are dependent on this medication may be born too soon (premature) and have low birth weight. They may also have withdrawal symptoms. Tell your doctor right away if you notice possible mood changes, agitation, or unusual tiredness in your newborn.

This medication passes into breast milk and may have undesirable effects on a nursing infant. Breast-feeding is not recommended while using this drug. Consult your doctor before breast-feeding.

DRUG INTERACTIONS:

Drug interactions may change how your

medications work or increase your risk for serious side effects. This document does not contain all possible drug interactions. Keep a list of all the products you use (including prescription/nonprescription drugs and herbal products) and share

it with your doctor and pharmacist. Do not start, stop, or change the dosage of any medicines without your doctor's approval.

Taking MAO inhibitors with this medication may cause a serious (possibly fatal) drug interaction. Avoid taking MAO inhibitors (isocarboxazid, linezolid, metaxalone, methylene blue, moclobemide, phenelzine, procarbazine, rasagiline, safinamide, selegiline, tranylcypromine) during treatment with this medication. Most MAO inhibitors should also not be taken for two weeks before treatment with this medication. Ask your doctor when to start or stop taking this medication.

Some products have ingredients that could raise your heart rate or blood pressure. Tell your pharmacist what products you are using, and ask how to use them safely (especially cough-and-cold products or diet aids).

The risk of serotonin syndrome/toxicity increases if you are also taking other drugs that increase serotonin. Examples include street drugs such as MDMA/ "ecstasy," St. John's wort, certain antidepressants (including SSRIs such as fluoxetine/paroxetine, SNRIs such as duloxetine/venlafaxine), among others. The risk of serotonin syndrome/toxicity may be more likely when you start or increase the dose of these drugs.

Lisdexamfetamine is very similar to amphetamine or dextroamphetamine. Do not use medications containing amphetamine or dextroamphetamine while using lisdexamfetamine.

This medication may interfere with certain laboratory tests (including blood and urine steroid levels), possibly causing false test results. Make sure laboratory personnel and all your doctors know you use this drug.

OVERDOSE:

If someone has overdosed and has serious symptoms such

as passing out or trouble breathing, call 911. Otherwise, call a poison control center right away. US residents can call their local poison control center at 1-800-222-1222. Canada residents can call a provincial poison control center. Symptoms of overdose may include: severe mental/mood changes (such as confusion, hallucinations), seizures, irregular heartbeat, severe restlessness, fast breathing.

NOTES:

Do not share this medication with others. Sharing it is

against the law.

Laboratory and/or medical tests (such as blood pressure, heart rate, growth monitoring in children) may be performed periodically to monitor your progress or check for side effects. Consult your doctor for more details.

MISSED DOSE:

If you miss a dose, take it as soon as you remember

in the morning hours. If it is afternoon or near the time of the next dose, skip the missed dose. Take your next dose at the regular time. Do not double the dose to catch up.

STORAGE:

Store at room temperature away from light and moisture.

Do not store in the bathroom. Keep all medications away from children and pets.

Do not flush medications down the toilet or pour them into a drain unless instructed to do so. Properly discard this product when it is expired or no longer needed. Consult your pharmacist or local waste disposal company.

Information last revised February 2022.

Copyright(c) 2022 First Databank, Inc.