



FOR ALL SEASONS II


16-0-8

Apply in early spring or fall 2 to 3 weeks prior to seed germination to obtain control of grassy and broadleaf weeds. For use on warm and cool season grasses. See back label for overseeding/reseeding restrictions.

Controls Henbit, Oxalis, Poa Annua, Chickweed, Crabgrass, Foxtail, Purslane, Shepherd's Purse, Prostrate Spurge, Knotweed and others listed on label.

Coverage: 5,000 sq. ft. For best results, application should be followed by 1/2 inch of rainfall or irrigation.

Always Read and Follow Label Directions



FOT All Scasons I Lawn Food Plus Grabgrass and Weed Preventer

For All Seasons II Lawn Food Plus Grabgrass and Weed Preventer

For All Seesons II Lawn Food Plus Grabstass and Weed Preventer

\sim	PRECAUTIONARY S	TATEMENTS		MAXIMUM ANNUAL RATES		
HAZARDS TO HUMANS AND DOMESTIC ANIMALS			This product is recommended for use on the kerf grass species listed in the following table. Do not exceed the maximum yearly rate as given in the following table.			
CAUTION: Causes moderate eye initiation. Avoid contact with eyes or clathing. Wear protective eyewear. Wash theroughly with scap and water after handling and before eating, drinking, chewing gum, using tobacco or using the beliet.			MAXIMUM APPLICATION RATE/CALENDAR VEAR OF			
FIRST AID			For All Seasons II Lawn Food Plus Crabgrass and Weed Preventer BY TUPE DRASS SPECIES ⁽⁷⁾ Turf Species: Ltbs. Products/1000 sq. ft. Ltbs. a.i/A			
If in Eyes:	Hold eye open and rinse slowly and gent Remove contact lenses, if present, after	ly with water for 15 to 20 minutes. 5 minutes, then continue rinsing.	Creeping Bentgrass	4.0 0.65		
If Swallowed:	Call a poison control center or doctor for Call a poison control center or doctor imm	treatment advice. nediately for treatment advice.	Creeping Red Fescue	4.7 0.75 6.2 1.0		
	Call a poisen control center or doctor imm Have person sip a glass of water if able t Do net induce vemiting unless told to do	o swallow. by a poison control center or doctor.	Buffalogitas Kentucky Bluegitas Perennial Ryegitasa	64 13		
H On Shin			Bermudaorass ²⁰	93 15		
If On Skin Or Cluthing:	 Rinse skin immediately with plenty of wa Call a poison control center or doctor for 	ter for 15 to 20 minutes. Institutes	Bahiagrass Centipedegrass Seashore Paspalum			
If Ishaled:	Move person to fresh air. If person is not breathing, call 911 or an preferably mouth to mouth if possible. Call a poison control center or doctor for	ambulance, then give artificial respiration,	St. Augustinegrass Tail Foscue (including turf-type) Zoysia			
Have this product In the event of a North America 1	Call a possion control center or doctor for container or label with you when calling a poison medical or chemical emergency contact -250-255-3924 or Werldwide Intl. + 01-813 ENVIRIONIMIENTAL	control center or doctor or when going for treatment. → 1/1 ≫. -248-0585	These are the memory notice per calculated year's typeople inhibitions. May to used on every compared or phanged memory and the second 45.5 h. a. Linene singular to 155 fbs/ A of this pended to respect to the second second second second second second second second second to any provide second			
This product has lo above the level of v organisms in adjace ditches, gutters or will help ensure the on a driveway, side or drainage system	w solubility in water. At the limits of solubility, th water solubility, it may be toxic to fish. Dirit are ent sites. To protect the environment, do not allo surface waters. Applying this product in cathr statiset or rain does not blow or wash pesificide water, or shreat, back onto the treated area of th 6.	is product is not taxic to fait. However, at encentrations renoff from twated ansais may be hazardous to equatic vegetidad to other or num off into atom draine, drainage wather when rain is not predicted for the next 24 hours of the brahamet rans. Sweeping my product that lands a lawn or garden will help prevent runoff to water bodies	Ibs./1000 sq.ft. of this product). • Use higher rates of this product to achie but do not exceed the maximum applica WHEN T Do not apply to overseeded tarf within 60	It cannot get you have been and a set of the large products by legal to do be higher levels of the get and begge products of set out hard type, after notice specified in the Maximum Arenael Rates Table. TO APPLY AFTER OVERSEEDUNG TUBF (dogs that seading secondary notes are in the second isch of set, net section applied before seeding secondary notes are in the second isch of set, net section applied before seeding secondary notes are in the second isch of set, net section applied before seeding secondary notes are in the second isch of set, net section applied before seeding secondary notes are in the second isch of set, net section applied before seeding secondary notes are in the second isch of set, net second second secon		
It is a violation of Fi	DIRECTIONS R idenal Law to use this product in a manner inco			WHEN TO OVERSEED AFTER APPLICATION		
 This product is a set Established to 	elective preemergence herbicide that provides r af grasses lexcluding golf course putting green	esidual control of many grass and broadleaf weeds in: and lawns	This product will inhibit the germination of turf species if overseeded too soon after application. Follow rates and intervals in the following table below for best overseeding/seeeding results.			
 Landscape or 	namentais		For All Seasons II Lawn Food Plas Cra and Weed Preventer (bs./1000 sq.			
This product control	ols susceptible weeds by inhibiting weed seeds	germination and root development. Most effective weed				
prior to weed seed	germination and within 14 days following appli-	germination and root development. Most effective weed awatal, impation, or shallow (1 to 2 inches) incorporation, ciai usa, (2) for commercial seed production, or (2) for halic purposes or climatic modification and being grown rounds. This product. Do not apply sensible to not apply to golf of infigation system.	3.1	0.50 4 4 4 4 0.65 5 4 4		
nex for use on pla research purposes.	ma owing grown for; (1) Sale or other comme For use on ornamental plants intended for aest	tow use, up of commercial seed production, or (3) for thetic purposes or climatic modification and being grown	47	0.75 6 5 5		
Do not graze or fee	ere or period, or on gon courses or silwits and g ed livestock forage cut from areas treated with set. Do not coole this another through	this product. Do not apply serially. Do not apply to golf	5.0	0.00 - 6 6		
FALURE TO FOLL	OW THE DIRECTIONS FOR USE AND PRECAU	of impation system. Itions on this label may result in poor weed	2.1	1.14 9		
CONTROL, GROP INJURY, OR ILLEGAL RESIDUES. WEEDS CONTROLLED			8.1	1.30 10		
When used in protestance with this label this numbert will ensuing control of the following wears-			2.5	1.50 12 SPREADER SETTINGS		
	Barnyardigeasa Bluegrass, Annual (Poa annua) Carpetweed Discloweed, common	Kochia Lambsquarter, common Lovegrass	Spreader settings vary by make and mode	el of spreader. It is recommended that individual spreaders are calibrated for the		
i i	Zhickweed, common Zhickweed, mouse ear (from seed)	Panicum (Texas, Fall, Browntop) Pinweart	specific product that is to be applied. A wa Spreader	iking speed of 3 miles per hour is recommended.		
6	Diabgrass (large, smooth) Diowfootgrass	Purslane, common Pusitey, Fiorida Boor semana ¹	spreaser	Spreader Settings Application Rates a.L.Ibs./ Acre (Product Ibs./1000 sq ft)		
Ē	Dickoved, common Dickoved, common eur (form seed) Dasbyrass (large, smooth) Drowfoodysias Dorograss, (foody instalis, Annual Doroograss' Senbit	Rescuegrass ¹ Shepherd's Purse ²	Scotts*/Republic Accupreen (Drop)	0.5 (3.1) 0.65 (4.0) 0.75 (4.7) 1.0 (6.2) 1.5 (8.3) 3-1/4 4 5 6-1/2 9-1/2		
	Soosegriks' tenbit	Septembrass, Broadleaf Speedwell, Persian	Scotts*/Republic Speedy Green (Broadca			
	tringsass Johnsongrass (Yom seed) Junglerice Knotweed	Rescuegrator Stephard's Pursa' Signalgrand, Riveran Spendened, Pursian Spangelan Spangelan Souray, Prostata Witchgraids Witchgraids Witchgraids	ferti+iome*/Earthway Ex-N-Spred (Broad			
Contained To Annuel			Book of the maximum annual application with of 13.9 Ju-1/2000 up, 15, 03.8 product/1000 up, 15, 03.0 product/1000 up, 15, 03.0 product/1000 up, 15, 03.0 product/1000 up, 15, 03.0 product/1000 up, 15, 05.0 product/1000			
	APPLICATION DI	RECTIONS	0.05% Water Soluble Copper (Ca) Inten (Fe)			
Apply uniformly with ESTABLISHED TUR	h suitable, calibrated application equipment. IF: This product is a selective preemerpence h	rhicide that, when properly applied, will control certain he maximum amount of this product that may be applied al Rates section of this label. Most effective weed control	0.10% Water Soluble Iron (Fe) Manganese (Me)			
grass and broadleat per year is given for	f weeds in established turf grasses and lawns. T r each turf grass species in the Maximum Annu	he maximum amount of this product that may be applied al Rates section of this label. Most effective weed control				
in turf grasses will I germination and will	be obtained when this product is activated by at thin 14 days following application. See the map I	least 0.5 inches of rainfall or irrigation prior to weed seed selow for approximate crabgrass seed germination dates.	2016 (201) 0.05% Water Soldele Zinc (Ze) Darived from: Ammonian Sulfak, Inrea, Mariala of Petaah, Sodiam Barate, Copper Sulfate, Iron Sulfak, Manganica Sulfak, and Zinc Sulfata. F370			
	CRABGRASS SEED GERM	MINATION DATES	Iron Sullate, Manganese Sullate, and Zinc Sullate. P370			
			STORAGE: Store this product in its origin feed, or seed by storage or disposal. PRODUCT DISPOSAL: As a responsible or of the bag be used, carefully libbowing lib CANTAINER DISPOSAL: Mourrelinitie	TORAGE AND DISPOSAL nal container in a dry, cool, secured area. Do not contaminate water, foodstuffe, nvironmental practico, where possible, it is recommended that all of the contents bol directions and processions.		
Presents tais Presents tais Presents tais Presents tais Presents tais Presents P						
Control and the second seco						
The product may be applied to a single application or in second applications to control service processing. For special service All applications are not applications of the service applications of						
Kenner Service Se						
11914 Viail la & www.fortione.com/vide/ Question? 355-278-47780fermation regaring the contents and leads of metals in the product is available on the internet at http://www.asplica.org/interlati.htm						