

LOC Tears Eye Drops

Instructions for use:

ERC/OLS/A/01



Unique Double-action Ocular Solution

LOC Tears is a Unique Double-action Lubricating and Osmoprotective Comfort Solution developed to fight the sensation of dryness, discomfort and irritation of eyes caused by external factors such as prolonged exposure to computer/LED screens, air-conditioning, dust, pollution, smoke, sunlight and other non-pathological factors.

CONTENTS:

LOC Tears contains Carmellose Sodium 0.5% preserved with the safer preservative Stabilised Oxychloro Complex, in a buffered isotonic solution with a neutral pH.

CONTRAINDICATIONS:

Hypersensitivity to carmellose sodium or to any of the ingredients listed below.

Other ingredients: Glycerin, Boric Acid, Calcium Chloride Dihydrate, Magnesium Chloride Hexahydrate, Erythritol, Levocarnitine, Potassium Chloride, Sodium Borate Decahydrate (Borax), Sodium Citrate Dihydrate.

INTENDED USE:

LOC Tears is uniquely formulated to have the following effects:

Lubrication and Washing- Supplements the aqueous layer of the natural tear film on the eye surface with additional water under conditions of significantly increased evaporation (due to external effects such as dry heat, air conditioning, etc.) and physically washes the eye surface (due to irritation caused by the external effects of dust, smoke, etc.).

Osmoprotection- Presence of compatible solutes in the solution helps restore the natural osmotic balance of the eye surface that is usually disrupted by external and non-pathological factors.

DIRECTIONS FOR USE:

Place 1 or 2 drops in the eye(s) as needed.



1



2

- Get the LOC Tears bottle and a mirror.
- Wash your hands.
- Take the bottle and twist off the cap.
- After cap is removed, if the tamper evident snap collar is loose, remove before using the product.
- Hold the bottle, pointing down, between your thumb and middle finger.
- Tilt your head back. Pull down your eyelid with a clean finger, until there is a 'pocket' between the eyelid and your eye. The drop will go in here (picture 1).
- Bring the bottle tip close to the eye. Use the mirror if it helps.
- Don't touch your eye or eyelid, surrounding areas or other surfaces with the dropper. It could infect the drops left in the bottle.
- Gently press on the base of the bottle to release one drop of LOC Tears at a time.
- Don't squeeze the bottle, it is designed so that just a gentle press on the bottom is needed (picture 2).
- If you use drops in both eyes, repeat the steps for your other eye.
- Put the bottle cap back on firmly immediately after use.

If a drop misses your eye, try again.

PRECAUTIONS:

- Do not use if allergic to any of the ingredients.
- Do not swallow solution.
- To avoid contamination or possible eye injury, do not touch bottle tip on any surface, and avoid direct contact with the eye.
- Close the bottle tightly immediately after use.
- Once opened, discard any unused

- solution after 1 month.
- Do not use after expiry date indicated on the carton & bottle.
- Do not use if package is damaged.
- Store in a dry, dark & well-ventilated place at a temperature not exceeding 25°C.
- Allow 5 minutes between the administration of other eye drops.
- Remove contact lenses before use.

KEEP OUT OF SIGHT AND REACH OF CHILDREN

SIDE EFFECTS:

Common: Eye irritation (including burning and discomfort), eye pain, eye pruritus, visual disturbance.

EXPECTED GROUP OF PATIENTS FOR OCULAR LUBRICATING & COMFORT SOLUTION :

Children above the age of 5 years, adults & geriatric population, who are affected by dry eyes (such as soreness, burning, irritation or dryness) or dry eye syndrome. Dry eye syndrome, also known as chronic dry eye, is characterised by a reduced ability to form tears or a reduction in the quality of those tears. Do not use if pregnant or breast-feeding.

WARNINGS: If you experience any problems with this product, please discuss with your pharmacist, GP or eye care specialist.

INSTRUCTION FOR DISPOSAL:

The device shall be disposed as per prevailing regulations where the devices are used.

HOW SUPPLIED:

LOC Tears is supplied in a 10 ml plastic bottle.

STERILE A

Sterilized by aseptic processing techniques



Keep away from sunlight



Do not sterilize



Do not use if package is damaged



Upper limit of temperature



See Package insert for more information.

10ml

EC REP

Kunze Indopharm BV,
Koningin Emmakade 171,
2518JN Den Haag,
Netherlands.



Entod Research Cell UK Ltd.
15 Tottenham Lane,
London N8 9DJ, UK.
Email: scientificell@entodpharma.com
Website: ophthalmicsuk.com

CE
0068