

INFECTION CONTROL 2025

1. Hands should be washed with soap and water after care of residents with infectious diarrhea like C-Diff. TRUE OR FALSE
2. Hand washing is less drying and damaging to skin than alcohol-based hand sanitizers. TRUE OR FALSE
3. Residents should be assisted with hand washing before meals, after toileting and intermittently throughout the day. TRUE OR FALSE
4. Gloves can replace hand washing in some circumstances. TRUE OR FALSE
5. Alcohol-based hand sanitizers are preferred over hand washing with soap and water when hands are not visible soiled. TRUE OR FALSE
6. Soap and water can be used to re-use gloves between residents as long as soap and water is used for 20 seconds and covers all surfaces. TRUE OR FALSE
7. The preferred maximum length for nail tips is ¼ inch. TRUE OR FALSE
8. One of the benefits of alcohol-based hand sanitizers is they are less dependent on technique and easier to coat all parts of your hands. TRUE OR FALSE
9. Skin becomes more fragile with age so resident hand washing should be limited to before meals and after toileting. TRUE OR FALSE
10. Name 3 ways to help prevent urinary tract infections (UTIs).

11. When wearing PPE, where do you take off your gloves and gown? _____
12. When wearing PPE, where do you take off your eye shield and mask? _____
13. In this video, what is one of the risks for taking antibiotics? _____
14. Name the 3 components that are how people get infection diseases.

15. What is the single most import defense against germs? _____

Name (print clearly): _____ Completed: AT HOME DURING WORK