



# ACTON Fe EDTA

## Guaranteed Analysis:

Iron (Fe) EDTA ..... 13.0%

Derived from: Iron-EDTA (ethylenediaminetetraacetic acid)

Information regarding the contents and levels of metals in this product is available on the internet at: <http://www.aapfco.org/metals.html>

ONLY A LICENSED AND EXPERIENCED APPLICATOR SHOULD APPLY THIS PRODUCT.

**Limited Warranty:** Manufacturer warrants that this product conforms to the guaranteed analysis for the growing season in which purchased. Manufacturer MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTIES. IN NO SUCH CASE SHALL MANUFACTURER OR THE SELLER BE LIABLE FOR CONSEQUENTIAL, INCIDENTAL, SPECIAL, EXEMPLARY OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. ANY AND ALL DAMAGES CLAIMED SHALL BE LIMITED TO THE AMOUNT PAID FOR THE PRODUCT.

Manufacturer offers this product and the Buyer/User accepts it subject to the forgoing Conditions of Sale and Limited Warranty.

### Precautionary Statement

Prevention: Observe good industrial hygiene practices.

Product may stain cloth and equipment. Response: Wash hands after handling.

Storage: Store away from incompatible materials.

Storage of bulk products is best suited for original manufacturing containers.

Harmful if swallowed. Avoid skin contact. Causes eye and skin irritations.

In case of contact, wash the affected areas with water immediately and continue washing for fifteen minutes. Seek medical attention if irritation persists.

Disposal: Dispose of contents in accordance with local regulations.

## EDTA CHELATED IRON FERTILIZER FOR SOIL AND FOLIAR APPLICATION

### Product information:

A water soluble EDTA chelated iron based product can be used for the prevention and correction of Fe deficiencies induced by lack of Fe or imbalances of Fe assimilation.

### Directions and General Recommendations

Can be applied via SOIL & FOLIAR APPLICATION

SOIL APPLICATION (MOST RECOMMENDED) \*Every 2-3 weeks\*:

- Vegetable 4 to 12 lbs/Acre.
- Field Crops 4 to 14 lbs/Acre.
- Tree crops, Fruits, Vines and Citrus: 4 to 14 lbs/Ac or 0.5 to 3 oz per tree depending on tree size
- Turf and Ornamentals: 0.5 lb/1000 sq ft or 20 lbs/Ac

FOLIAR APPLICATION \*Every 2-3 weeks\*:

- Vegetable: 0.25 - 1.5 lbs/Ac in 25 gal water
  - Field Crops: 0.25-1.2 lbs/Ac in 25 gal water
  - Tree crops, Fruits, Vines and Citrus: 0.25-2 lbs/Ac in 100 gal water. Do not apply during bloom.
  - Turf and Ornamentals: 1 - 4 oz/1000 sq ft in adequate spray water volume for full coverage
  - Optimum rate of application will vary depending on soil properties (such as pH, organic matter content, texture), weather conditions, time of year, general crop health and crop species. For best results, follow soil test or plant analysis recommendation.
- DO NOT MIX WITH PESTICIDE SPRAYS ON FRUIT.  
Always jar test new combinations for compatibility prior to field mixing.

KEEP OUT OF REACH OF CHILDREN



ACTON CORPORATION  
1451 S. Elm Eugene St,  
Greensboro, NC 27406  
Phone: (336) 230-6831  
[info@actonfertilizers.com](mailto:info@actonfertilizers.com)  
[www.actonfertilizers.com](http://www.actonfertilizers.com)



Net Weight:		
1 LB	5 LBS	55 LBS
(0.45Kg)	(2.25Kg)	(25Kg)

LOT #  
MFG. DATE  
EXP. DATE