

SEAWAVE

Guaranteed Analysis: (0-0-3.5)

Soluble Potash (K2O) 3.59

Derived from: Seaweed extract (Ascophyllum nodosum) and Potassium Hydroxide.

Information regarding the contents and levels of metals in this product is available on the internet at: http://www.aapfco.org/metals.html

ONLY A LICENSED AND EXPERIENCED APPLICATOR SHOULD APPLY THIS PRODUCT.

Limited Warranty: Manufacturer warrants that this product conforms to the guaranteed analysis for the growing season in which purchased. Manufacturer MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTIES. IN NO SUCH (ASS SHALL MANUFACTURER OR THE SELLER BE LIABLE FOR CONSEQUENTIAL, INCIDENTAL, SPECIAL, EXEMPLARY OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. ANY AND ALL DAMAGES CLAIMED SHALL BE LIMITED TO THE AMOUNT PAID FOR THE PRODUCT.

Manufacturer offers this product and the Buyer/User accepts it subject to the forgoing Conditions of Sale and Limited Warranty.

Precautionary Statement

Prevention: Observe good industrial hygiene practices.

Product may stain cloth and equipment. Response: Wash hands after handling. Storage: Store away from incompatible materials.

Storage of bulk products is best suited for original manufacturing containers.

Harmful if swallowed. Avoid skin contact. Causes eye and skin irritations. In case of contact, wash the affected areas with water immediately and continue washing for fifteen minutes. Seek medical attention if irritation persists. Disposa: Dispose of contents in accordance with local regulations.

LIQUID SEAWEED EXTRACT FOR SOIL AND FOLIAR APPLICATION

A Seaweed Extract is suitable for use in any foliar, soil applied, or irrigation water application. It can be applied to seed, seed pieces, bulbs, tubers and cuttings. Apply in early morning or late afternoon for best results. Allow foliar product to dry on plant prior to overhead irrigation.

Directions and General Recommendations

FOLIAR APPLICATION RATES: Apply 2-4 quarts per acre per application. FERTIGATION APPLICATION RATES: Drip irrigation: Apply 2-4 quarts per acre per application. Micro-sprinklers: Apply 2-4 quarts per acre per application. Water-run(flood): Apply 2-4 quarts per acre per application.

Optimum rate of application will vary depending on soil properties (such as pH, organic matter content, texture), weather conditions, time of year, general crop health and crop species. For best results, follow soil test or plant analysis recommendation.

Always jar test new combinations for compatibility prior to field mixing. NOTICE: Perform a phytotoxicity test on a small plot before large-scale application. To lower the risk of phytotoxicity, applications should be made early morning or late evening.

KEEP OUT OF REACH OF CHILDREN SHAKE WELL BEFORE USE





ACTON CORPORATION 1451 S. Elm Eugene St, Greensboro, NC 27406 Phone: (336) 230-6831 info@actonfertilizers.com www.actonfertilizers.com

