

BACK PAIN MYTH BUSTERS

We're busting myths and reinforcing what the latest evidence says is best for your back

Myth

Moving will make my back pain worse

Fact:

Don't fear
twisting and
bending - it's essential
to keep moving.
Gradually increase
how much you
are doing, and
stay on the go.

Myth I should

exercise, especially weight training

Fact:

Back pain shouldn't stop you enjoying exercise or regular activities. In fact, studies found that continuing with these can help you get better sooner - including using weights where appropriate.

Myth

A scan will show me exactly what is wrong

Fact:

Sometimes it will, but most often it won't. Also, even people without back pain have changes in their spine so scans can cause fear that influences behaviour, making the problem worse.

Myth

Pain equals damage

Fact:

This was the established view but more recent research has changed our thinking. Modern physio takes a holistic approach that helps people understand why they are in pain.



BACK PAIN MYTH BUSTERS

We're busting myths and reinforcing what the latest evidence says is best for your back

Fight the fear – move more

Half of all people experience back pain at some point in their lives and it is one of the leading causes of sickness absence.

The good news: we are constantly learning more about it.

Physiotherapists

are the experts in treating - and preventing - back pain but hear many myths about what is causing it and what can make it better.

It's easy to understand why they are held - indeed some may have been the established view before the evidence base moved on.

Problem is these myths create fear, which causes people to stop doing many of the activities they need to do to address the problem.

So let's bust those myths and reinforce what the latest evidence says is best for your back.

Find out more about the myths and full references at:

www.csp.org.uk /mythbusters

If you have pain that lasts for more than six weeks, please consult a physio or your GP.