ADHIKAANSH ACADEMY (IITJEE NEET IX X XI XII)

RUN BY:

DEEPAK SAINI SIR

B.TECH, M.TECH (N.S.I.T. DELHI UNIVERSITY)

Ex. Faculty of

Resonance Kota, Career Point Kota

Aakash Institute Mumbai

BIOLOGY NOTES (CLASS 11TH)



FOR MORE FREE QUESTION BANKS AND SAMPLE PAPER WT'S UP ON: 7665186856

A Best Faculty Group in Meerut....

ADHIKAANSH X ACADEMY IITJEE NEET IX X XI XII

सही समय और सही दिशा में कड़ी मेहनत, तो सफलता पक्की!

So why to wait..

और $m{Class} ~ m{11}^{th}$ से ही शुरू करें, $m{Board}$ के साथ-साथ, $m{NEET}$ और $m{IITJEE}$ की तैयारी

Special Classes For Students Studying in std. 9th & 10th.

DIRECTOR
DEEPAK SAINI (DSA SIR)
B.TECH, M.TECH
NSIT DELHI UNIVERSITY

EX. FACULTY OF RESONANCE KOTA, AAKASH INSTITUTE MUMBAI

225/5, 1ST FLOOR, PANCHSHEEL COLONY, BEHIND PINNACLE TOWER, GARH ROAD, MEERUT. @ 8057870069

www.adhikanshiitjeemedical.com | Sadhikanshiitjeemedical@gmail.com

Take Dummy Admission in Class 11th & Prepare For..

IITJEE & NEET

For Free Counselling & Career Guidance contact on:

7665186856

CBSE Quick Revision Notes (Class-11 Biology) CHAPTER-12 MINERAL NUTRITION

Mineral nutrition is the study of source, mode of absorption, distribution and metabolism of various inorganic substances (minerals) by plants for their growth, development, structure, physiology and reproduction.


Methods to study the Mineral Requirement of Plants

- **Hydroponics** is the technique of growing plants in nutrient solution in complete absence of soil. This method is used to determine the nutrients essential for plants.
- In this method plant is cultured in soil-free, defined mineral solution. These methods require purified water and mineral nutrients.
- Essential elements are identified and their deficiency symptoms are discovered by hydroponics methods. It is also used to commercial

production of vegetables, like tomato and cucumber.

Essential mineral nutrients- About 65 elements are found in different plants. Following criteria is used to determine the essentiality of an element.

- a. Element must be absolutely necessary for the normal growth and reproduction to complete their life cycle.
- b. The requirement of element must be specific and not replaceable.
- c. Element must be directly involved in the metabolism of plants.
- Macronutrients are present in plants tissues in larger quantity. C,H and O is obtained from water and rest are absorbed from soil.
- Micronutrients or trace nutrients are required in vary small quantity.



Funnel for

adding water and nutrients Cotton


Aerating

tube

Nutrient

solution

On the basis of diverse functions, essential elements are divided into following categories-



Role of Macro and Micro nutrients-

- (i) Essential elements participate in various metabolic processes in plants such as permeability of cell membrane, maintenance of osmotic potential, ETS. Etc.
- (ii) Act as major constituents of macromolecules and co-enzymes.

Various forms and function of essential nutrients-

- a. **Nitrogen-** required by plants in greatest amount, it is absorbed by plants as NO₂-,NO₃-and NH₄+. It is one of the major constituent of proteins, nucleic acids and vitamins.
- b. **Phosphorus-** Absorbed by plants from soil in form of phosphate ions. It is the constituent of cell membrane, all nucleic acids and nucleotides require phosphorus.
- c. **Potassium** absorbed as potassium ions (K⁺). Help to maintain cation-anion balance in cells. It is involved in protein synthesis, opening and closing of stomata.
- d. **Calcium** absorbed by plants from soil in form of Calcium ions (Ca²⁺). Used in synthesis of cell wall. It activates certain enzymes.
- e. **Magnesium-** absorbed by plants in form of Mg²⁺ ions. It activates the enzymes for respiration, photosynthesis, and involved in DNA and RNA. Constituent of chlorophyll.
- f. **Sulphur-** plants obtain sulphur in form of sulphate (SO_4^{2-}) . Present in amino acids (cysteine, methionine) and is main constituent of coenzymes and vitamins.
- g. **Iron-** obtains in form of ferric iron (Fe³⁺). It is important constituents of protein involved in transport system.
- h. **Manganese-** absorbed in form of Mn²⁺ ions. Main function is splitting of water to liberate Hydrogen and Oxygen during photosynthesis.
- i. **Zinc-** obtained as Zn^{2+} ions. Activate enzymes like carboxylases. Needed in formation of Auxin.
- j. **Copper** absorbed as cupric ions(Cu²⁺). Involved in various metabolic activities and redox reactions.
- k. **Boron-** absorbed as BO_3^{3-} or $B_4O_7^{-}$ ions. Required for uptake of calcium, cell elongation and pollen germination.
- l. **Chlorine** it is absorbed in form of Cl⁻ ions. Determine the solute concentration and splitting of water during photosynthesis.

Deficiency Symptoms of Essential elements

- ➤ When supply of essential elements becomes limited, plant growth is retarded. The concentration of essential elements below which plant growth is retarded is called **critical concentration.**
- In absence of any particular element shows certain morphological changes. These morphological changes are called **deficiency symptoms**.
- The parts of plant that show deficiency symptoms depend upon mobility of elements in the plants. Elements that are actively mobilized (N,Mg,K) shows deficiency in older regions. On the other hand, symptoms appear first in young region if the elements are relatively immobile (Ca) and not transported out of mature tissues.

Kinds of deficiency syndrome are as follows-

Deficiency Disease	Symptoms	Deficient elements
Chlorosis	Loss of chlorophyll leading	N, K, Mg, S, Fe, MN, Zn, Mo
	to yellowing of leaves.	
Necrosis	Death of tissue (leaf).	
Stunted plant growth	:	Fe, K.
Premature fall of leaves and buds.	Falling of leaves and buds.	P, Mg, Cu


- ➤ Deficiency of any element may cause many symptoms or same symptoms may be caused by different elements. To identify the deficient elements various symptoms are compared with standard chart.
- **Toxicity of micronutrients-** in higher doses, micronutrients become toxic. Any tissue concentration which reduces dry weight of tissue by 10% is called toxic concentration. Critical toxic concentration is different for different elements.
- Mechanism of absorption of elements takes place in two phases. In first phase, rapid intake of ions occurs in free space or outer space of the cells, apoplast. In second phase, ions are taken slowly into inner space, the symplast of the cells.
- Passive movement of ions in apoplast occurs through ion channels and trans-membrane protein. On the other hand, movement of ions into symplast occurs by expenditure of energy by active process.
- The movement of ion is called **flux**. The inward movement is called **influx** and outward movement is called **efflux**.
- **Translocation of solutes** occur through xylem along with ascending stream of water

Soil as reservoir of essential elements- most of the nutrients required for growth and development is obtained from soil by roots. These minerals are formed by weathering or

rocks. Soil also harbor nitrogen fixing bacteria and other microbes, holds water and supplies air to roots. Deficiency of essential elements affects the crop yield so, fertilsers are used to supplement these elements.

Metabolism of Nitrogen

- Nitrogen is the most prevalent elements in living world along with C, H and O. It is the main constituent of proteins, nucleic acids, fats, hormones, enzymes etc.
- The process of conversion of nitrogen to ammonia is called **nitrogen fixation**. In nature lightening and ultraviolet radiation provide energy to convert atmospheric



- nitrogen into nitrogen oxide (N0, N02 and N20).
- Industrial combustion, forest fire and automobiles along with thermal power plants produce nitrogen oxides.
- The decomposition of organic nitrogen of dead plants and animals into ammonia is called **ammonification**.
- Ammonia is first oxidized to nitrite by bacteria *Nitrosomonas* or *Nitrococcus*. Which is further oxidized to nitrate with help of bacteria *Nitrobactor*. These process are called nitrification.

$$2NH_3 + 3O_2 \longrightarrow 2NO_2^- + 2H^+ + 2H_2O$$

$$2NO_2^- + O_2 \longrightarrow 2NO_3^-$$


- Nitrates formed is absorbed by plants and transported to leaves. Nitrates is converted into free nitrogen by the process called **denitrificaion** by bacteria *Pseudomonas* and *Thiobacilus*.
- Reduction of nitrogen to ammonia by living organism is called **Biological Nitrogen Fixation**. The enzyme nitrogenase, present in prokaryotic organism called nitrogen fixer.

$$N = N \xrightarrow{\text{Nitrogenase}} NH_3$$

- Nitrogen fixing microbes may be symbiotic (Rhizobium) or free living (Nostoc, Azotobactor, Anabaena).
- Symbiotic biological nitrogen fixation includes legume-bacteria relationship in which rod shaped Rhizobium lives with symbiotic relation with nodules of Leguminous plants.
- Central portion of nodule is pink or red due to presence of leguminous haemoglobin or leghaemoglobin.

Nodule formation involves sequence of interaction between root and Rhizobium as follows-

- Ahizobia increase in number and attach with epidermis of roots. Root hairs curls and bacteria invade it. An infection thread is formed that carrying the bacteria into cortex of root.
- Nodule formation starts in cortex of root. Bacteria released from thread to cells which leads to formation of specialized nitrogen fixing cells.
- Nodules establish direct vascular connection with host with for exchange of nutrients.
- ❖ Nodule contain all necessary biochemical components like enzyme nitrogenase and leghaemoglobin.




Enzyme nitrogense is a Mo-Fe protein and catalyzes the conversion of atmospheric nitrogen into ammonia.

The reaction is as follows-

$$N_2 + 8e^- + 8H^+ + 16ATP \longrightarrow 2NH_3 + H_2 + 16ADP + 16P_1$$

- The enzyme nitrogenase is highly sensitive to molecular oxygen and needs anaerobic condition. To protect this enzyme from oxygen, the nodules contain an oxygen scavenger called leg-haemoglobin.
- **❖** The ammonia synthesized by nitrogenase enzyme require large amount of energy (18ATP) for each NH₃ produced.



Fat of ammonia- at physiological pH, ammonia is converted into ammonium ions (NH₄⁺).It is toxic for plants in larger concentration and ammonium ion is converted into amino acids by two methods-

a. **Reductive animation**- in this process ammonia reacts with α -ketoglutaric acid to form glutamic acid.

$$\alpha$$
 - ketoglutaric acid + NH₄⁺ + NADPH $\xrightarrow{\text{Glutamate}}$ glutamate + H₂O + NADP

b. **Transamination**- it involves the transfer of amino group from amino acids to keto group of keto acid. Glutamic acid is the main amino acid from which transfer of NH₃ takes place and another amino acid is formed by transamination.

Two important amides asparagine and glutamine found in plants in proteins. They are formed from aspartic acid and glutamic acid by addition of another amino groups to it.

FOR MORE FREE QUESTION BANKS AND SAMPLE PAPER WT'S UP ON: 7665186856

A Best Faculty Group in Meerut....

ADHIKAANSH X ACADEMY

सही समय और सही दिशा में कड़ी मेहनत, तो सफलता पक्की!

So why to wait..

और $m{Class} ~ m{11}^{th}$ से ही शुरू करें, $m{Board}$ के साथ-साथ, $m{NEET}$ और $m{IITJEE}$ की तैयारी

Special Classes For Students Studying in std. 9th & 10th.

DIRECTOR
DEEPAK SAINI (DSA SIR)
B.TECH, M.TECH
NSIT DELHI UNIVERSITY

EX. FACULTY OF RESONANCE KOTA, AAKASH INSTITUTE MUMBAI

225/5, 1ST FLOOR, PANCHSHEEL COLONY, BEHIND PINNACLE TOWER, GARH ROAD, MEERUT.
8057870069

www.adhikanshiitjeemedical.com | adhikanshiitjeemedical@gmail.com

Take Dummy Admission in Class 11th & Prepare For..

IITJEE & NEET

For Free Counselling & Career Guidance contact on:

7665186856