

# **ADHIKAANSH ACADEMY (IITJEE NEET IX X XI XII)**

**RUN BY:**

**DEEPAK SAINI SIR**


**B.TECH, M.TECH (N.S.I.T. DELHI UNIVERSITY)**

**Ex. Faculty of**

**Resonance Kota, Career Point Kota**

**Aakash Institute Mumbai**

# **CHEMISTRY NOTES (CLASS 12<sup>TH</sup>)**



**FOR MORE FREE QUESTION  
BANKS AND SAMPLE PAPER  
WT'S UP ON : 7665186856**

**A Best Faculty Group in Meerut....**

**ADHIKAANSH ACADEMY**  
IITJEE NEET IX X XI XII

**सही समय और सही दिशा में कड़ी मेहनत, तो सफलता पक्की!**

*So why  
to wait...*



**DIRECTOR**  
**DEEPAK SAINI (DSA SIR)**  
B.TECH, M.TECH  
NSIT DELHI UNIVERSITY

**और Class 11<sup>th</sup> से ही शुरू करें,  
Board के साथ-साथ,  
NEET और IITJEE की तैयारी**

**Special Classes For Students Studying in std. 9th & 10th.**

**EX. FACULTY OF RESONANCE KOTA, AAKASH INSTITUTE MUMBAI**  
225/5, 1ST FLOOR, PANCHSHEEL COLONY, BEHIND PINNACLE TOWER, GARH ROAD, MEERUT. ☎ 8057870069  
www.adhikanshiitjeemedical.com | adhikanshiitjeemedical@gmail.com

**Take Dummy Admission  
in Class 11th & Prepare For..**

**IITJEE & NEET**

**For Free Counselling &  
Career Guidance contact on:**

**7665186856**

---

**CBSE Class-12 Chemistry Quick Revision Notes**  
**Chapter-04: Chemical Kinetics**

---

- **Chemical kinetics:**  
It is the branch of chemistry that deals with the study of reaction rates and their mechanisms.
- **Rate of reaction:**  
It is the change in concentration of reactant (or product) in unit time.
- The unit of rate of reaction is mol L<sup>-1</sup>s<sup>-1</sup>.
- $A + B \rightarrow C + D$

$$\text{Rate of disappearance of } A = \frac{-d[A]}{dt}$$

where d[A] is small change in conc. of 'A' and dt is small interval of time

$$\text{Rate of disappearance of } B = \frac{-d[B]}{dt}$$

Where d[B] is small change in conc. of 'B' and dt is small interval of time

$$\text{Rate of appearance of } C = \frac{+d[C]}{dt}$$

Where d[C] is small change in conc. of 'C' and dt is small interval of time

$$\text{Rate of appearance of } D = \frac{+d[D]}{dt}$$

Where d[D] is small change in conc. of 'D' and dt is small interval of time

$$\text{Rate} = \frac{-d[A]}{dt} = \frac{-d[B]}{dt} = \frac{+d[C]}{dt} = \frac{+d[D]}{dt}$$

- **Rate law or rate equation:**  
It is the expression which relates the rate of reaction with concentration of the reactants. The constant of proportionality 'k' is known as rate constant.
- **Average rate:**  
It is the rate of reaction measured over a long time interval.  
$$\text{Average rate} = \frac{\Delta x}{\Delta t}$$
  
where  $\Delta x$  is change in concentration and  $\Delta t$  is large interval of time.
- **Instantaneous rate:**  
It is the rate of reaction when the average rate is taken over a particular moment of time.  
$$\text{Instantaneous rate} = \frac{dx}{dt}$$
  
where dx is small change in conc. and dt is the smallest interval of time.

---

It is the expression which relates the rate of reaction with concentration of the reactants.

- Rate constant:

When the concentration of reactants is unity, then the rate of reaction is known as rate constant. It is also called specific reaction rate.

- The constant of proportionality 'k' is known as rate constant.

- Molecularity of a reaction:

The total number of atoms, ions or molecules of the reactants involved in the reaction is termed as its molecularity. It is always in whole number and is never more than three. It cannot be zero.

- Order of a reaction:

The sum of the exponents (power) of the concentration of reactants in the rate law is termed as order of the reaction. It can be in fraction. It can be zero also.

- If rate law expression for a reaction is

$$\text{Rate} = k [A]^x [B]^y$$

Then its order of reaction = x + y

- Order cannot be determined with a given balanced chemical equation. It can be experimentally determined.

- Integrated rate law for zero order reaction:



$$\frac{dx}{dt} = k[R]^0$$

$$k = \frac{[R_0] - [R]}{t}$$

- If we plot a graph between concentration of R vs time t, the graph is a straight line with slope equal to -k and intercept is equal to [R<sub>0</sub>].

- Half- life of a reaction:

The time taken for a reaction, when half of the starting material has reacted is called half- life of a reaction.

- For zero order reaction, the half-life time is

$$t_{1/2} = \frac{[R_0]}{2k}$$

- For first order reaction, the half-life time is

$$t_{1/2} = \frac{0.693}{k}$$

where 'k' is rate constant

It is independent of initial concentration for first order reaction.

- Rate law for first order reaction:



$$k = \frac{2.303}{t} \log \frac{[R_0]}{[R]}$$

where 'k' is rate constant or specific reaction rate, [Ro] is initial molar conc., [R] is final molar conc. after time 't'.

$$k = \frac{2.303}{t} \log \frac{a}{a-x}$$

where 'a' is initial conc. reacted in time 't' final conc., after time 't' is (a - x).


- If we plot a graph between ln[R] with time, we get a straight line whose slope = - k and intercept ln[Ro].
- To calculate rate constant for first order gas phase reaction of the type  
A (g) → B(g) + C(g)

$$k = \frac{2.303}{t} \log \frac{p_i}{(2p_i - p_t)}$$

Where pi is initial pressure of A, pt is total pressure of gaseous mixture containing A, B, C

$$p_t = p_A + p_B + p_C$$

- Pseudo first order reaction:  
The reaction which is bimolecular but order is one is called pseudo first order reaction. This happens when one of the reactants is in large excess.  
Example - Acidic hydrolysis of ester (ethyl acetate).



- Activation energy (E<sub>a</sub>):  
It is extra energy which must be possessed by reactant molecules so that collision between reactant molecules is effective and leads to the formation of product molecules.
- Arrhenius equation of reaction rate:  
It gives the relation between rate of reaction and temperature.

$$k = Ae^{-E_a/RT}$$

$$\ln k = \ln A - E_a / RT$$

$$\log k = \log A - \frac{E_a}{2.303RT}$$

where k = rate constant, A = frequency factor, E<sub>a</sub> = energy of activation R = gas constant, T = temperature in Kelvin,

$$\log \frac{k_2}{k_1} = \frac{E_a}{2.303R} \left( \frac{T_2 - T_1}{T_1 T_2} \right)$$

- Probability factor or Steric factor

$$Rate = PZ_{AB} \cdot e^{\frac{-E_a}{RT}}$$

Where Z<sub>AB</sub> represents the collision frequency of reactants, A and B,  $e^{\frac{-E_a}{RT}}$  represents the fraction of molecules with energies equal to or greater than E<sub>a</sub> and P is called the probability or steric factor.

- Mechanism of reaction:

---

It is the sequence of elementary processes leading to the overall stoichiometry of a chemical reaction.

- **Activated complex:**

It is an unstable intermediate formed between reacting molecules. Since, it is highly unstable and it readily changes into product.

- **Rate determining step:**

It is the slowest step in the reaction mechanism.

- **The number of collisions per second per unit volume of the reaction mixture is known as collision frequency ( $Z$ ).**

FOR MORE FREE QUESTION  
BANKS AND SAMPLE PAPER  
WT'S UP ON : 7665186856

A Best Faculty Group in Meerut....

**ADHIKAANSH**  **ACADEMY**  
IITJEE NEET IX X XI XII

सही समय और सही दिशा में कड़ी मेहनत, तो सफलता पक्की!

So why  
to wait...



**DIRECTOR**  
**DEEPAK SAINI (DSA SIR)**  
B.TECH, M.TECH  
NSIT DELHI UNIVERSITY

और **Class 11<sup>th</sup>** से ही शुरू करें,  
**Board** के साथ-साथ,  
**NEET** और **IITJEE** की तैयारी

Special Classes For Students Studying in std. 9th & 10th.

**EX. FACULTY OF RESONANCE KOTA, AAKASH INSTITUTE MUMBAI**  
225/5, 1ST FLOOR, PANCHSHEEL COLONY, BEHIND PINNACLE TOWER, GARH ROAD, MEERUT. ☎ 8057870069  
www.adhikanshiitjeemedical.com | adhikanshiitjeemedical@gmail.com

Take Dummy Admission  
in Class 11th & Prepare For..

**IITJEE & NEET**

For Free Counselling &  
Career Guidance contact on:

**7665186856**