# Class X (CBSE 2019) Mathematics Abroad (Set-3)

#### **General Instructions:**

(i) All questions are compulsory.

(ii) The question paper consists of **30** questions divided into four sections – **A**, **B**, **C** and **D**.

(iii) Section A comprises 6 questions of 1 mark each. Section B contains 6 questions of 2 marks each. Section C contains 10 questions of 3 marks each. Section D contains 8 questions of 4 marks each.

(iv) There is no overall choice. However, an internal choice has been provided in **two** questions of **1** mark, **two** questions of **2** marks, **four** questions of **3** marks each and **three** questions of **4** marks each. You have to attempt only **one** of the alternative in all such questions.

(v) Use of calculators is **not** permitted.

# Question 1

Which term of the A.P. -4, -1, 2, ... is 101?

# SOLUTION:

We have been given an arithmetic progression where a = -4, d = -1 - (-4) = 3 and  $a_n = 101$ We need to find which term of the given AP is 101 so, we need to find *n*. Using  $a_n = a + (n - 1)d$ Substituting the values in the formula we get 101 = -4 + (n - 1)3 101 + 7 = 3n 3n = 108 n = 36Therefore, 36th term of given A.P is 101.

# **Question 2**

Evaluate: tan 65° cot 25°

OR

Express (sin 67° + cos 75°) in terms of trigonometric ratios of the angle between 0° and 45°.

# SOLUTION:

 $\frac{\tan 65^{\circ}}{\cot 25^{\circ}}$   $= \frac{\tan(90^{\circ} - 25^{\circ})}{\cot 25^{\circ}} \quad (\because \tan (90^{\circ} - \theta) = \cot \theta)$   $= \frac{\cot 25^{\circ}}{\cot 25^{\circ}}$  = 1

OR

 $(\sin 67^\circ + \cos 75^\circ)$ 

= $(\sin(90^\circ - 23^\circ) + \cos(90^\circ - 25^\circ))$  (:  $\sin(90^\circ - \theta) = \cos\theta$  and  $\cos(90^\circ - \theta) = \sin\theta$ ) =  $(\cos 23^\circ + \sin 25^\circ)$ 

#### **Question 3**

Find the value of *k* for which the quadratic equation kx(x-2) + 6 = 0 has two equal roots.

## **SOLUTION:**

Given quadratic equation is: kx (x - 2) + 6 = 0  $\Rightarrow kx^2 - 2kx + 6 = 0$ For a quadratic equation to have equal roots,  $b^2 - 4ac = 0$ Comparing the given equation with general equation  $ax^2 + bx + c = 0$ We get a = k, b = -2k and c = 6  $(-2k)^2 - 4 (k) (6) = 0$   $\Rightarrow 4k^2 - 24k = 0$   $\Rightarrow 4k (k - 6) = 0$ Therefore, k = 0 and k = 6

#### **Question 4**

Find a rational number between  $\sqrt{2}$  and  $\sqrt{7}$ .

OR

Write the number of zeroes in the end of a number whose prime factorization is  $2^2 \times 5^3 \times 3^2 \times 17$ .

# **SOLUTION:**

We know  $\sqrt{2} = 1.414$  $\sqrt{7} = 1.732$ So, rational number between  $\sqrt{2}$  and  $\sqrt{7}$  will be  $1.5 = \frac{3}{2}$ .

OR

Given prime factorisation is  $2^2 \times 5^3 \times 3^2 \times 17$ . A number will have zero at the end when we have  $2 \times 5$ . In  $2^2 \times 5^3 \times 3^2 \times 17$  we will have 2 zeroes as  $(2^2 \times 5^2) \times 5 \times 3^2 \times 17$ .

# Question 5

Find the distance between the points (a, b) and (-a, -b).

# SOLUTION:

Using distance formula:  $d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$ Here,  $x_1 = a$ ,  $y_1 = b$ ,  $x_2 = -a$  and  $y_2 = -b$ On substituting the values in the formula we get  $\sqrt{(-a - a)^2 + (-b - b)^2}$   $= \sqrt{(-2a)^2 + (-2b)^2}$  $= \sqrt{4a^2 + 4b^2}$ 

$$= \sqrt{4a^2 + 4b^2}$$
$$= 2\sqrt{a^2 + b^2}$$

Therefore, the distance between (a, b) and (-a, -b) is  $2\sqrt{(a)^2 + (b)^2}$ 

# **Question 6**

Let  $\triangle$  ABC  $\sim \triangle$  DEF and their areas be respectively, 64 cm<sup>2</sup> and 121 cm<sup>2</sup>. If EF = 15.4 cm, find BC.

# **SOLUTION:**

#### Given: $\triangle ABC \sim \triangle DEF$

We know ratio of the areas of two similar triangles is equal to the square of the ratio of their corresponding sides.

$$\frac{\operatorname{ar} \triangle \operatorname{ABC}}{\operatorname{ar} \triangle \operatorname{DEF}} = \left(\frac{\operatorname{BC}}{\operatorname{EF}}\right)^{2}$$

$$\Rightarrow \frac{64}{121} = \left(\frac{\operatorname{BC}}{15.4}\right)^{2}$$

$$\Rightarrow \left(\frac{8}{11}\right)^{2} = \left(\frac{\operatorname{BC}}{15.4}\right)^{2}$$

$$\Rightarrow \frac{8}{11} = \frac{\operatorname{BC}}{15.4}$$

$$\Rightarrow \operatorname{BC} = \frac{8 \times 15.4}{11} = 11.2 \text{ cm}$$
Thus, BC = 11.2 cm.

## **Question 7**

Find the solution of the pair of equation :  $\frac{3}{x} + \frac{8}{y} = -1; \ \frac{1}{x} - \frac{2}{y} = 2, \ x, \ y \neq 0$ OR

Find the value(s) of k for which the pair of equations

$$\left\{egin{array}{l} kx+2y=3\ 3x+6y=10 \end{array}
ight.$$
 has a unique solution.

# SOLUTION:

The given equations are  $\frac{3}{x} + \frac{8}{y} = -1 \qquad \dots (1)$   $\frac{1}{x} - \frac{2}{y} = 2 \qquad \dots (2)$ Let  $\frac{1}{x} = u$  and  $\frac{1}{y} = v$ (1) and (2) will become  $3u + 8v = -1 \qquad \dots (3)$  $u - 2v = 2 \qquad \dots (4)$ 

Multiply (4) with 4  

$$4u - 8v = 8$$
 ..... (5)  
Adding (3) and (5) we get  
 $7u = 7$   
 $\Rightarrow u = 1$   
Putting this value in (4)  
 $1 - 2v = 2$   
 $\Rightarrow v = \frac{-1}{2}$ 

Now  $\frac{1}{x} = u$   $\Rightarrow \frac{1}{x} = 1$   $\Rightarrow x = 1$ And  $\frac{1}{y} = v$   $\Rightarrow \frac{1}{y} = \frac{-1}{2}$  $\Rightarrow y = -2$ 

OR

The given equations are kx + 2y = 3 3x + 6y = 10For a unique solution,  $\frac{a_1}{a_2} \neq \frac{b_1}{b_2}$ where  $a_1 = k$ ,  $a_2 = 3$ ,  $b_1 = 2$ ,  $b_2 = 6$   $\frac{k}{3} \neq \frac{2}{6}$   $\Rightarrow k \neq 1$ For all values of *k* except 1, the given linear equations will have unique solution.

## **Question 8**

Use Euclid's division algorithm to find the HCF of 255 and 867.

# SOLUTION:

The given numbers are 255 and 867.

Now 867 > 255. So, on applying Euclid's algorithm we get

867=255×3+102

Now the remainder is not 0 so, we repeat the process again on 255 and 102

255=102×2+51

The algorithm is applied again but this time on the numbers 102 and 51

102=51×2+0

Thus, the HCF obtained is 51.

## **Question 9**

The point *R* divides the line segment AB, where A (-4, 0) and B (0, 6) such that  $AR = \frac{3}{4}AB$ . Find the coordinates of *R*.

## **SOLUTION:**

We have given that R divides the line segment AB

AR+ RB= AB

$$\begin{split} \frac{3}{4}AB + RB &= AB \\ \Rightarrow RB = \frac{AB}{4} \\ \Rightarrow AR : RB = 3 : 1 \\ \text{Using section formula:} \\ x &= \left(\frac{m_1 x_2 + m_2 x_1}{m_1 + m_2}\right), \ y = \left(\frac{m_1 y_2 + m_2 y_1}{m_1 + m_2}\right) \\ m_1 &= 3, \ m_2 = 1 \\ x_1 &= -4, \ y_1 = 0 \\ x_2 &= 0, \ y_2 = 6 \\ \text{Plugging values in the formula we get} \\ x &= \frac{3 \times 0 + 1 \times (-4)}{3 + 1}, \ y = \frac{3 \times 6 + 1 \times 0}{3 + 1} \\ x &= \frac{-4}{4}, \ y = \frac{18}{4} \\ \Rightarrow x &= -1, \ y = \frac{9}{2} \\ \text{Therefore, the coordinates of } R\left(-1, \frac{9}{2}\right). \end{split}$$

# Question 10

How many multiples of 4 lie between 10 and 205?

Determine the A.P. whose third term is 16 and 7<sup>th</sup> term exceeds the 5<sup>th</sup> by 12.

# SOLUTION:

We need to find the number of multiples of 4 between 10 and 205. So, multiples of 4 gives the sequence 12, 16, ..., 204 a = 12, d = 4 and  $a_n = 204$ Using the formula  $a_n = a + (n-1)d$ Plugging values in the formula we get 204 = 12 + (n - 1)4 204 = 12 + 4n - 4 4n = 196 n = 49Thus, there are 49 multiples of 4 between 10 and 205.

-		
63	ы.	
ີ	π.	
-	••	

Given: 3rd term of the AP is 16.  $a_3 = 16$ 

a + (3 - 1)d = 16

a + 2d = 16 .....(1) Also, 7th term exceeds the 5th term by 12.

 $a_7 - a_5 = 12$ 

[a+(7-1)d] - [a+(5-1)d] = 12

(a + 6d) - (a + 4d) = 12

2d = 12

*d* = 6

From equation (1), we obtain

a + 2(6) = 16

*a* + 12 = 16

*a* = 4

Therefore, A.P. will be 4, 10, 16, 22, ...

#### **Question 11**

Three different coins are tossed simultaneously. Find the probability of getting exactly one head.

#### SOLUTION:

Total possible outcomes of tossing three coins simultaneously are {TTT,TTH,THT,THH,HTT,HTH,HHT,HHH}

that is 8

We have to find the probability of getting exactly one head.

Cases of exactly one head are {TTH,THT,HTT}

that is 3

Probability of getting exactly on head is  $\frac{3}{\alpha}$ .

# Question 12

A die is thrown once. Find the probability of getting. (a) a prime number. (b) an odd number

# SOLUTION:

Total outcomes of throwing a dice once are 1, 2, 3, 4, 5 and 6

(1)Probability of getting a prime number Prime numbers are 2, 3 and 5 in throwing a die once Probability of getting a prime number  $= \frac{3}{6} = \frac{1}{2}$ (2)Probability of getting an odd number odd numbers are those that are not divisible by 2. So, there three odd numbers in throwing a dice once which is 1, 3 and 5. Probability of getting an odd number  $= \frac{3}{6} = \frac{1}{2}$ 

## **Question 13**

In Figure 1, BL and CM are medians of a  $\triangle ABC$  right-angled at A. Prove that 4 (BL<sup>2</sup> + CM<sup>2</sup>) = 5 BC<sup>2</sup>.



OR

Prove that the sum of the squares of the sides of a rhombus is equal to the sum of the squares of its diagonals.

#### **SOLUTION:**

To prove:  $4 (BL^2 + CM^2) = 5 BC^2$ Proof: In  $\triangle CAB$ , Applying Pythagoras theorem,  $AB^2 + AC^2 = BC^2$  .....(1) In  $\triangle ABL$ ,  $AL^2 + AB^2 = BL^2$   $\Rightarrow (\frac{AC}{2})^2 + AB^2 = BL^2$   $\Rightarrow AC^2 + 4 AB^2 = 4 BL^2$  .....(2) In  $\triangle CAM$ ,  $CA^2 + MA^2 = CM^2$   $\Rightarrow (\frac{BA}{2})^2 + CA^2 = CM^2$  $\Rightarrow BA^2 + 4 CA^2 = 4 CM^2$  .....(3)

Adding (2) and (3)  

$$AC^{2} + 4AB^{2} + BA^{2} + 4CA^{2} = 4BL^{2} + 4CM^{2}$$

$$\Rightarrow 5AC^{2} + 5AB^{2} = 4(BL^{2} + CM^{2})$$

$$\Rightarrow 5(AC^{2} + AB^{2}) = 4(BL^{2} + CM^{2})$$

$$\Rightarrow 5(BC^{2}) = 4(BL^{2} + CM^{2}) \quad (From (1))$$
Hence Proved.

OR



In  $\triangle AOB$ ,  $\triangle BOC$ ,  $\triangle COD$ ,  $\triangle AOD$ ,

Applying Pythagoras theorem, we obtain

$AB^2 = AO^2 + OB^2$	(1)
$BC^2 = BO^2 + OC^2$	(2)
$CD^2 = CO^2 + OD^2$	(3)
$AD^2 = AO^2 + OD^2$	(4)

Adding all these equations, we obtain

$$AB^{2} + BC^{2} + CD^{2} + AD^{2} = 2(AO^{2} + OB^{2} + OC^{2} + OD^{2})$$

$$= 2\left[\left(\frac{AC}{2}\right)^2 + \left(\frac{BD}{2}\right)^2 + \left(\frac{AC}{2}\right)^2 + \left(\frac{BD}{2}\right)^2\right]$$

(Diagonals bisect each other)

$$=2\left(\frac{(AC)^{2}}{2} + \frac{(BD)^{2}}{2}\right)$$
$$= (AC)^{2} + (BD)^{2}$$

#### **Question 14**

In Figure 2, two concentric circles with centre O, have radii 21 cm and 42 cm. If  $\angle AOB = 60^{\circ}$ , find the area of the shaded region.



**SOLUTION:** 



Radius of inner circle, OC = 21 cm

Radius of outer circle, OA = 42 cm

Area of circle with radius  $R = \pi R^2 = \pi (42)^2$ Area of circle with radius  $r = \pi r^2 = \pi (21)^2$ Area of sector AOB =  $\frac{\theta}{360} \times \pi R^2 = \frac{60}{360} \times \pi (42)^2 = \frac{\pi (42)^2}{6}$ Area of sector COD =  $\frac{\theta}{360} \times \pi r^2 = \frac{60}{360} \times \pi (21)^2 = \frac{\pi (21)^2}{6}$ Area of shaded portion = Area of circle with radius R – Area of circle with

radius *r* – [Area of sector AOB – Area of sector COD]

$$= \pi (42)^2 - \pi (21)^2 - \left[\frac{\pi (42)^2}{6} - \frac{\pi (21)^2}{6}\right]$$
  
=  $\pi \left[ (42)^2 - (21)^2 - \frac{1}{6} \left[ (42)^2 - (21)^2 \right] \right]$   
=  $\pi \left[ \left( (42)^2 - (21)^2 \right) \left( 1 - \frac{1}{6} \right) \right]$   
=  $\pi \left[ (42 - 21) \left( 42 + 21 \right) \frac{5}{6} \right]$   
=  $\frac{22}{7} \times \frac{5}{6} \times 21 \times 63$   
=  $3465 \text{ cm}^2$ 

#### **Question 15**

A cone of height 24 cm and radius of base 6 cm is made up of modelling clay. A child reshapes it in the form of a sphere. Find the radius of the sphere and hence find the surface area of this sphere.

#### OR

A farmer connects a pipe of internal diameter 20 cm from a canal into a cylindrical tank in his field which is 10 m in diameter and 2 m deep. If water flows through the pipe at the rate of 3 km/hr, how much time will the tank be filled ?

#### **SOLUTION:**

A cone has been reshaped in sphere Height of cone is 24 cm and radius of base is 6 cm Volume of sphere = volume of cone Volume of cone =  $\frac{1}{3}\pi r^2 h$ Plugging the values in the formula we get volume of cone =  $\frac{1}{3}\pi (6)^2 24$ =  $288\pi \text{ cm}^3$ Let the radius of sphere be *r* Volume of sphere =  $\frac{4}{3}\pi r^3$  Since, volume of cone = volume of sphere Volume of sphere =  $288\pi \text{ cm}^3$ So,  $288\pi = \frac{4}{3}\pi r^3$   $\Rightarrow 288 = \frac{4}{3}r^3$   $\Rightarrow r^3 = 216$   $\Rightarrow r = 6 \text{ cm}$ Hence, radius of reshaped sphere is 6 cm Now, surface area of sphere =  $4\pi r^2$   $= 4\pi (6)^2$   $= 144 \times \frac{22}{7}$   $= 452.5 \text{ cm}^2$ Therefore, surface area of sphere is 452.57 cm<sup>2</sup>.

#### OR



Consider an area of cross-section of pipe as shown in the figure.

Radius (*r*<sub>1</sub>) of circular end of pipe = 
$$\frac{20}{200} = 0.1 \text{ m}$$

Area of cross-section =  $\pi \times r_1^2 = \pi \times (0.1)^2 = 0.01\pi \text{ m}^2$ 

Speed of water = 3 km/h = 
$$\frac{3000}{60}$$
 = 50 metre/min

Volume of water that flows in 1 minute from pipe =  $50 \times 0.01\pi = 0.5\pi \text{ m}^3$ 

Volume of water that flows in *t* minutes from pipe =  $t \times 0.5\pi$  m<sup>3</sup>



Radius (*r*<sub>2</sub>) of circular end of cylindrical tank  $=\frac{10}{2}=5$ m

Depth  $(h_2)$  of cylindrical tank = 2 m

Let the tank be filled completely in *t* minutes.

Volume of water filled in tank in t minutes is equal to the volume of water flowed in *t* minutes from the pipe.

Volume of water that flows in *t* minutes from pipe = Volume of water in tank

 $t \times 0.5\pi = \pi \times (r_2)^2 \times h_2$ 

 $t \ge 0.5 = 5^2 \ge 2$ 

t = 100

Therefore, the cylindrical tank will be filled in 100 minutes.

## **Question 16**

Calculate the mode of the following distribution :

Class :	10 - 15	15 – 20	20 - 25	25 - 30	30 - 35
Frequency :	4	7	20	8	1

# **SOLUTION:**

Modal class is the class with highest frequency modal class is 20 - 25 lower limit of modal class i.e l = 20class size i.e h = 5frequency of modal class  $f_1=20$ frequency of preceding class  $f_0=7$ frequency of succeeding class  $f_2=8$ Using the formula  $\mathrm{mode} = l + \left(rac{f_1 - f_0}{2f_1 - f_0 - f_0}
ight) imes h$ 

Plugging the values in the formula we get

mode =  $20 + \left(\frac{20-7}{2 \times 20-7-8}\right) \times 5$ mode =  $20 + \left(\frac{13}{25}\right) \times 5$ mode =  $20 + \frac{13}{5}$ mode =  $\frac{113}{5} = 22.6$ Question 17

Show that  $\frac{2+3\sqrt{2}}{7}$  is not a rational number, given that  $\sqrt{2}$  is an irrational number.

# SOLUTION:

To prove:  $\frac{2+3\sqrt{2}}{7}$  is irrational, let us assume that  $\frac{2+3\sqrt{2}}{7}$  is rational.  $\frac{2+3\sqrt{2}}{7} = \frac{a}{b}$ ;  $b \neq 0$  and a and b are integers.  $\Rightarrow 2b + 3\sqrt{2}b = 7a$   $\Rightarrow 3\sqrt{2}b = 7a - 2b$   $\Rightarrow \sqrt{2} = \frac{7a - 2b}{3b}$ Since a and b are integers so, 7a - 2b will also be an integer. So,  $\frac{7a - 2b}{3b}$  will be rational which means  $\sqrt{2}$  is also rational. But we know  $\sqrt{2}$  is irrational(given). Thus, a contradiction has risen because of incorrect assumption. Thus,  $\frac{2+3\sqrt{2}}{7}$  is irrational.

## **Question 18**

Obtain all the zeroes of the polynomial  $2x^4 - 5x^3 - 11x^2 + 20x + 12$  when 2 and - 2 are two zeroes of the above polynomial

## **SOLUTION:**

We know that if  $x = \alpha$  is a zero of a polynomial, and then  $x - \alpha$  is a factor of f(x).

Since 2 and -2 are zeros of f(x).

Therefore

 $(x-2)(x+2) = x^2 - 4$  $(x^2 - 4)$  is a factor of f(x). Now, we divide  $2x^4 - 5x^3 - 11x^2 + 20x + 12$  by  $g(x) = (x^2 - 4)$  to find the zero of f(x).

$$\begin{array}{r} 2x^2 - 5x - 3 \\
x^2 - 4 \overline{\smash{\big)}2x^4 - 5x^3 - 11x^2 + 20x + 12} \\
\underline{2x^4 + 0x^3 - 8x^2} \\
\underline{- - +} \\
- 5x^3 - 3x^2 + 20x + 12 \\
\underline{- 5x^3 + 0x^2 + 20x} \\
\underline{+ - -} \\
- 3x^2 + 12 \\
\underline{- 3x^2 + 12} \\
\underline{- 3x^2 + 12} \\
0
\end{array}$$

By using division algorithm we have  $f(x) = g(x) \times q(x) - r(x)$ 

$$egin{aligned} &2x^4-5x^3-11x^2+20x+12=ig(x^2-4ig)ig(2x^2-5x-3ig)\ &=(x-2)\,(x+2)\,[2x\,(x-3)+(x-3)]\ &=(x-2)\,(x+2)\,(x-3)\,(2x+1) \end{aligned}$$

Hence, the zeros of the given polynomial are 2,  $-2,\ 3,\ rac{-1}{2}$  .

#### **Question 19**

A motorboat whose speed is 18 km/hr in still water takes on hour more to go 24 km upstream than to return downstream to the same spot. Find the speed of the stream.

## **SOLUTION:**

Speed of the motorboat is 18 km/h Let us assume the speed of th stream be x km/h speed of motorboat upstream is (18 - x) km / hspeed of motorboat downstream is (18 + x) km / hTime taken to go downstream is  $\left(\frac{24}{18+x}\right) \text{ hr}$ Time taken to go upstream is  $\left(\frac{24}{18-x}\right) \text{ hr}$ Equation becomes:  $\frac{24}{18-x} - \frac{24}{18+x} = 1$ Solving the above equation  $\frac{24(18+x)-24(18-x)}{(18-x)(18+x)} = 1$  $\frac{432+24x-432+24x}{324-x^2} = 1$ 

$$48x = 324 - x^{2}$$

$$x^{2} + 48x - 324 = 0$$
Solving for the value of x  
using  $x = \frac{-b \pm \sqrt{b^{2} - 4ac}}{2a}$ 

$$x = \frac{-48 \pm \sqrt{(48)^{2} - 4(1)(-324)}}{2}$$

$$x = \frac{-48 \pm \sqrt{2304 + 1296}}{2}$$

$$x = \frac{-48 \pm \sqrt{3600}}{2}$$

$$x = \frac{-48 \pm 60}{2}$$

$$x = \frac{12}{2}, \frac{-108}{2}$$

$$x = 6, -54$$

Since, speed can not be negative So, the speed of the stream is 6 km/h.

#### Question 20

Prove that:

 $(\sin \theta + 1 + \cos \theta) (\sin \theta - 1 + \cos \theta)$ . sec  $\theta$  cosec  $\theta = 2$ 

OR

Prove that :

$$\sqrt{\frac{\sec \,\theta - 1}{\sec \,\theta + 1}} + \sqrt{\frac{\sec \,\theta + 1}{\sec \,\theta - 1}} = 2 \operatorname{cosec} \,\theta$$

#### SOLUTION:

$$\begin{split} \mathrm{LHS} &= (\sin\theta + 1 + \cos\theta) \left(\sin\theta - 1 + \cos\theta\right) . \sec\theta \csc\theta \\ &= \left[\sin^2\theta - \sin\theta + \sin\theta \cos\theta + \sin\theta - 1 + \cos\theta + \sin\theta \cos\theta - \cos\theta + \cos^2\theta\right] \frac{1}{\cos\theta} \frac{1}{\sin\theta} \quad \left(\because \sec\theta = \frac{1}{\cos\theta} \text{ and } \csc\theta = \frac{1}{\sin\theta}\right) \\ &= \left[1 + 2\sin\theta \cos\theta - 1\right] \frac{1}{\cos\theta} \frac{1}{\sin\theta} \\ &= \left[2\sin\theta \cos\theta\right] \frac{1}{\cos\theta} \frac{1}{\sin\theta} \\ &= 2 = \mathrm{RHS} \\ \mathrm{Hence \ proved} \end{split}$$

$$\sqrt{\frac{\sec \theta - 1}{\sec \theta + 1}} + \sqrt{\frac{\sec \theta + 1}{\sec \theta - 1}} = \frac{\sqrt{\sec \theta - 1}}{\sqrt{\sec \theta + 1}} + \frac{\sqrt{\sec \theta + 1}}{\sqrt{\sec \theta - 1}}$$
$$= \frac{\sqrt{\sec \theta - 1}\sqrt{\sec \theta - 1} + \sqrt{\sec \theta + 1}}{\sqrt{\sec \theta - 1}}$$
$$= \frac{(\sqrt{\sec \theta - 1})^2 + (\sqrt{\sec \theta + 1})^2}{\sqrt{(\sec \theta - 1)(\sec \theta + 1)}}$$
$$= \frac{\sec \theta - 1 + \sec \theta + 1}{\sqrt{\sec^2 \theta - 1}}$$
$$= \frac{2 \sec \theta}{\sqrt{\tan^2 \theta}}$$
$$= \frac{2 \sec \theta}{\tan \theta}$$
$$= \frac{2 \frac{1}{\cos \theta}}{\cos \theta}$$
$$= 2 \frac{1}{\sin \theta}$$
$$= 2 \cos ec\theta$$

#### **Question 21**

In what ratio does the point P(-4, y) divide the line segment joining the points A(-6, 10) and B(3, -8) ? Hence find the value of y.

OR

Find the value of p for which the points (-5, 1), (1, p) and (4, -2) are collinear.

#### **SOLUTION:**

Let P divides the line segment AB in the ratio k: 1 Using section formula

 $egin{aligned} x &= rac{m_1 x_2 + m_2 x_1}{m_1 + m_2}, y = rac{m_1 y_2 + m_2 y_1}{m_1 + m_2} \ \mathrm{A} \left(-6, \ 10 
ight) ext{ and } \mathrm{B} \left(3, \ -8 
ight) \ m_1 \ \colon m_2 \ = \ k \ \colon 1 \end{aligned}$ 

plugging values in the formula we get

$$-4 = \frac{k \times 3+1 \times (-6)}{k+1}, \ y = \frac{k \times (-8)+1 \times 10}{k+1}$$

$$-4 = \frac{3k-6}{k+1}, \ y = \frac{-8k+10}{k+1}$$
Considering only *x* coordinate to find the value of *k*  

$$-4k-4 = 3k-6$$

$$-7k = -2$$

$$k = \frac{2}{7}$$

$$k : 1 = 2 : 7$$
Now, we have to find the value of *y*  
so, we will use section formula only in *y* coordinate to find the value of *y*  

$$y = \frac{2 \times (-8)+7 \times 10}{2+7}$$

$$y = \frac{-16+70}{9}$$

$$y = 6$$
Therefore, P divides the line segment AB in 2 : 7 ratio  
And value of *y* is 6.

OR

Points are collinear means the area of triangle formed by the collinear points is 0. Using

area of triangle = 
$$\frac{1}{2} [x_1 (y_2 - y_3) + x_2 (y_3 - y_1) + x_3 (y_1 - y_2)]$$
  
=  $\frac{1}{2} [-5 (p - (-2)) + 1 (-2 - 1) + 4 (1 - p)]$   
=  $\frac{1}{2} [-5 (p + 2) + 1 (-3) + 4 (1 - p)]$   
=  $\frac{1}{2} [-5p - 10 - 3 + 4 - 4p]$   
=  $\frac{1}{2} [-5p - 9 - 4p]$   
Area of triangle will be zero points being collinear  
 $\frac{1}{2} [-5p - 4p - 9] = 0$   
 $\frac{1}{2} [-9p - 9] = 0$   
 $9p + 9 = 0$   
 $p = -1$   
Therefore, the value of  $p = -1$ .

#### **Question 22**

ABC is a right triangle in which  $\angle B = 90^{\circ}$ . If AB = 8 cm and BC = 6 cm, find the diameter of the circle inscribed in the triangle.

#### **SOLUTION:**



#### **Question 23**

In an A.P., the first term is -4, the last term is 29 and the sum of all its terms is 150. Find its common difference

# SOLUTION:

a = -4, l = 29 and  $S_n = 150$ Using  $S_n = \frac{n}{2} [a + l]$   $150 = \frac{n}{2} [-4 + 29]$  300 = 25n n = 12Now, we will find dusing  $a_n = a + (n - 1)d$ Plugging the values in the formula we get 29 = -4 + (12 - 1)d 33 = 11d d = 3Therefore, the common difference is 3.

## **Question 24**

Draw a circle of radius 4 cm. From a point 6 cm away from its centre, construct a pair of tangents to the circle and measure their lengths

# **SOLUTION:**



## Step of construction

**Step: I-** First of all we draw a circle of radius AB = 4 cm.

Step: II- Mark a point P from the centre at a distance of 6 cm from the point O.

Step: III -Draw a perpendicular bisector of OP, intersecting OP at Q.

**Step:** IV- Taking Q as centre and radius OQ = PQ, draw a circle to intersect the given circle at T and T'.

Step: V- Join PT and PT' to obtain the required tangents.

Thus, PT and PT' are the required tangents.

The length of  $PT = PT' \approx 4.5 \text{ cm}$ 

# Question 26 Solve for $x:\frac{1}{2a+b+2x} = \frac{1}{2a} + \frac{1}{b} + \frac{1}{2x}; \ x \neq 0, \ x \neq \frac{-2a-b}{2}, \ a, \ b \neq 0$

The sum of the areas of two squares is  $640 \text{ m}^2$ . If the difference of their perimeters is 64 m, find the sides of the square.

#### **SOLUTION:**

$$\frac{1}{2a+b+2x} = \frac{1}{2a} + \frac{1}{b} + \frac{1}{2x}$$

$$\frac{1}{2a+b+2x} - \frac{1}{2x} = \frac{1}{2a} + \frac{1}{b}$$

$$\frac{2x-2a-b-2x}{4ax+2bx+4x^2} = \frac{b+2a}{2ab}$$

$$(-2a-b)(2ab) = (b+2a)(4ax+2bx+4x^2)$$

$$\frac{-(b+2a)(2ab)}{(b+2a)} = (4ax+2bx+4x^2)$$

$$-(2ab) = (4ax+2bx+4x^2)$$

$$4x^2 + 2bx + 4ax + 2ab = 0$$

$$2x(2x+b) + 2a(2x+b) = 0$$

$$2x(2x+b) + 2a(2x+b) = 0$$

$$\Rightarrow (2x+2a)(2x+b) = 0$$

$$\Rightarrow x = -a$$
or
$$(2x+b) = 0$$

$$\Rightarrow x = \frac{-b}{2}$$

Therefore, values of x are -a and  $\frac{-b}{2}$ 

Let the side of one square be x And the side of other square be y Sum of area of two square is 640 Equation becomes  $x^2 + y^2 = 640$ (: area of square is  $side^2$ ) .....(1) Now, difference of their perimeters is 64 Equation becomes 4x - 4y = 64 (: perimeter of square is  $4 \times \text{side}$ ) x - y = 16....(2) $\Rightarrow x = y + 16$ Solving the two equation by substitution method Substitute (2) in (1) $(16+y)^2 + y^2 = 640$  $256 + y^2 + 32y + y^2 = 640$  $2y^2 + 32y - 384 = 0$  $y^2 + 16y - 192 = 0$ Using  $y=rac{-b\pm\sqrt{b^2-4ac}}{2a}$ Plugging the values in the formula we get  $y = rac{-16 \pm \sqrt{256 - 4(-192)}}{2}$  $y = \frac{-16 \pm \sqrt{1024}}{2}$  $y = \frac{-16 \pm 32}{2}$  $y = \frac{-48}{2}, \frac{16}{2}$ y = -24, 8Since, sides can not be negative Therefore, y = 8Put y = 8 in (2) *x* = 8+ 16 x = 24Therefore, the sides of square are 24 m and 8 m.

**Question 27** 





#### SOLUTION:

Applying Pythagoras theorem in  $\triangle$ ADB, we obtain

 $AD^2 + DB^2 = AB^2$ 

 $\Rightarrow AD^2 = AB^2 - DB^2 \qquad \dots \dots (1)$ 

Applying Pythagoras theorem in  $\triangle$ ADC, we obtain

 $AD^{2} + DC^{2} = AC^{2}$   $AB^{2} - BD^{2} + DC^{2} = AC^{2} [Using equation (1)]$   $AB^{2} - BD^{2} + (BC - BD)^{2} = AC^{2}$   $AC^{2} = AB^{2} - BD^{2} + BC^{2} + BD^{2} - 2BC \times BD$   $AC^{2} = AB^{2} + BC^{2} - 2BC \times BD$ 

#### **Question 28**

A moving boat is observed from the top of a 150 m high cliff moving away from the cliff. The angle of depression of the boat changes from 60° to 45° in 2 minutes. Find the speed of the boat in m/min.

#### OR

There are two poles, one each on either bank of a river just opposite to each other. One pole is 60 m high. From the top of this pole, the angle of depression of the top and foot of the other pole are 30° and 60° respectively. Find the width of the river and height of the other pole.

**SOLUTION:** 



Let AO be the cliff of height 150 m. Let the speed of boat be x metres per minute. And BC be the distance which man travelled. So, BC = 2x [: Distance = Speed × Time]  $\tan (60^{\circ}) = \frac{AO}{OB}$   $\sqrt{3} = \frac{150}{OB}$   $\Rightarrow OB = \frac{150\sqrt{3}}{3} = 50\sqrt{3}$   $\tan (45^{\circ}) = \frac{AO}{OC}$   $\Rightarrow 1 = \frac{150}{OC}$   $\Rightarrow OC = 150$ Now OC = OB + BC  $\Rightarrow 150 = 50\sqrt{3} + 2x$   $\Rightarrow x = \frac{150-50\sqrt{3}}{2}$  $\Rightarrow x = 75 - 25\sqrt{3}$ 

Using  $\sqrt{3} = 1.73$  $x = 75 - 25 \times 1.732 \approx 32 \text{ m/min}$ Hence, the speed of the boat is 32 metres per minute.



Let the width of the river be w. In  $\triangle ABC$ ,

$$\tan 60^{\circ} = \frac{AB}{BC}$$

$$\Rightarrow \sqrt{3} = \frac{60}{w}$$

$$\Rightarrow w = \frac{60}{\sqrt{3}} = \frac{60\sqrt{3}}{3} = 20\sqrt{3}$$

$$\ln \triangle AED,$$

$$\tan 30^{\circ} = \frac{AE}{ED}$$

$$\Rightarrow \frac{1}{\sqrt{3}} = \frac{AE}{w}$$

$$\Rightarrow \frac{1}{\sqrt{3}} = \frac{AE}{20\sqrt{3}}$$

$$\Rightarrow AE = 20$$
Height of pole CD = AB - AE  

$$= 60 - 20 = 40 \text{ m}$$
Thus, width of river is  $20\sqrt{3} = 20 \times 1.732 = 34.64 \text{ m}$ 
Height of pole = 40 m

## **Question 29**

Calculate the mean of the following frequency distribution :

Class :	10-30	30-50	50-70	70-90	90-110	110-130
Frequency :	5	8	12	20	3	2

OR

The following table gives production yield in kg per hectare of wheat of 100 farms of a village :

Production yield (kg/hectare) :	40-45	45-50	50-55	55-60	60-65	65-70
Number of farms	4	6	16	20	30	24

Change the distribution to a 'more than type' distribution, and draw its ogive.

## **SOLUTION:**

Class	frequency $(f_i)$	Class mark $(x_i)$	$f_i x_i$
10-30	5	$\frac{10+30}{2} = 20$	100
30-50	8	$\frac{30+50}{2} = 40$	320
50-70	12	$\frac{50+70}{2} = 60$	720
70-90	20	$\frac{70+90}{2} = 80$	1600
90-110	3	$\frac{90+110}{2} = 100$	300
110-130	2	$\frac{110+130}{2} = 120$	240
	$\sum f_i = 50$		$\sum f_i x_i = 3280$

Using: mean =  $\frac{\sum f_t x_t}{\sum f_t}$ substituting the values in the formula mean=  $\frac{3280}{50} = 65.6$ 

#### OR

Production yield	Cumulative frequency
more than 40	100
more than 45	96
more than 50	90
more than 55	74
more than 60	54
more than 65	24



## **Question 30**

A container opened at the top and made up of a metal sheet, is in the form of a frustum of a cone of height 16 cm with radii of its lower and upper ends as 8 cm and 20 cm respectively. Find the cost of milk which can completely fill the container, at the rate of ₹ 50 per litre. Also find the cost of metal sheet used to make the container, if it costs ₹ 10 per 100 cm<sup>2</sup>. (Take  $\pi$  = 3.14)

## **SOLUTION:**

We have to find the cost of milk which can completely fill the container Volume of container = Volume of frustum

$$=rac{1}{3}\pi h\left(r_{1}^{2}+r_{2}^{2}+r_{1}r_{2}
ight)$$

Here, height = 16 cm radius of upper end = 20 cm

And radius of lower end = 8 cm Plugging the values in the formula we get Volume of container  $= \frac{1}{3} \times 3.14 \times 16 ((20)^2 + (8)^2 + 20 \times 8)$ =  $\frac{1}{3} \times 50.24 (400 + 64 + 160)$ =  $\frac{1}{3} \times 50.24 (624)$ = 10449.92 cm<sup>3</sup> = 10.449 litre (: 1 litre = 1000 cm<sup>3</sup>)

Cost of 1 litre milk is Rs 50 Cost of 10.449 litre milk = 50 × 10. 449 = Rs 522.45 We will find the cost of metal sheet to make the container Firstly, we will find the area of container Area of container = Curved surface area of the frustum + area of bottom circle (:: container is closed from bottom) Area of container =  $\pi (r_1 + r_2)l + \pi r^2$ Now, we will find /  $l = \sqrt{h^2 + (r_1 - r_2)^2}$   $l = \sqrt{(16)^2 + (20 - 8)^2}$   $l = \sqrt{(16)^2 + (12)^2}$   $l = \sqrt{256 + 144}$   $l = \sqrt{400}$ l = 20 cm

Area of frustum =  $3.14 \times 20(20+8)$ = 1758.4 cm<sup>2</sup>

Area of bottom circle =  $3.14 \times 8^2 = 200.96 \text{ cm}^2$ Area of container = 1758.4 + 200.96=  $1959.36 \text{ cm}^2$ 

Cost of making 100 cm<sup>2</sup> = Rs 10 Cost of making 1 cm<sup>2</sup> =  $\frac{10}{100}$  = Rs  $\frac{1}{10}$ Cost of making 1959. 36 cm<sup>2</sup> =  $\frac{1}{10} \times 1959.36$  = 195. 936

Hence, cost of milk is Rs 522.45 And cost of metal sheet is Rs 195.936