Matrix

Acid Crop Mix

Guaranteed Analysis

Zinc (Zn)	 	4%
Manganese (Mn)		
Copper (Cu)		
Boron (B)		0.02%

Derived from Zinc Chloride, Manganese Chloride, Copper Chloride, Boric Acid, and Citric Acid

CAUTION: KEEP OUT OF REACH OF CHILDREN. HARMFUL IF SWALLOWED.

Avoid contact with skin, eyes, and clothing. In case of contact, immediately flush eyes or skin with water for 15 minutes.

General Information

MATRIX CROP MIX can be used for economical correction of zinc, Manganese, Copper, deficiency of vegetables, row crops, small grains, deciduous trees and vines, citrus, and turf. MATRIX CROP MIX is specially designed for both soil applications and foliar applications.

Directions for Use: This product is recommended to correct specific micronutrient deficiencies. This product should be used on the basis of a soil and/or tissue analysis. For fertilizer application rates suitable for your geographical area or the maximum allowable application rate per acre, consult a certified drop advisor or a trained agronomist. For specific crop recommendations in your area, consult your state's recommendations.

MATRIX CROP MIX is designed to be more effective when used with polyphosphates and nitrogen solutions as a soil application for all crops requiring zinc, manganese, copper, and magnesium.

Row or Band Application: apply 1-2 qts per acre just before or at planting time.

Broadcast Application: apply 2-5 qts per acre just before planting and work into the top 3-5 inches of soil. Foliar Application: for foliar application this product may be used at the rate of 1-2 pints per acre with a sufficient amount of water to insure good coverage. Do not exceed one quart per gallon of water. Optimum rate of application will vary between fields depending on soil pH and organic matter content. For best results follow soil test or plant analysis recommendations. To insure even distribution on the leaf surfaces and increase the effectiveness of foliar application a small amount of wetting agent or surfactant may be added. Spray to thoroughly cover foliage. Rates and timing of applications are dependent on local conditions and practices. Apply 4-6 weeks after planting.

Liquid Fertilizer: Mix MATRIX CROP MIX with liquid fertilizer agitating to thoroughly disperse the trace elements. This can be achieved by adding the product to the mixing batch tank or inducting through a pump. When mixing in batch tank, add product to waste nitrogen solution before adding phosphate material. Matric Chemicals Inc. also manufactures a complete line of chelates for high-analysis fertilizer.

MATRIX CROP MIX is compatible with most insecticides, fungicides, foliar nutrients and herbicides. It can be applied in existing spray programs. Use 1 quart per acre in banded application and 2 quarts per acre in broadcast application. An ammonium Chloride source is recommended when tank mixing with glyphosate and some other herbicides.

MATRIX CROP MIX is manufactured to rigid controls at the highest possible concentration for effective usage. Any residue in this container is water soluble. Rinse the container with water and add solution to spray tank.

NET CONTENT AND WEIGHT: 275 gls (1,041 L) - 10.6 lbs./gl.

