

Please join us for a Speaker Program Event

tivdak
tisotumab vedotin-tftv
for injection 40 mg

Learn more about Tivdak, including a review of:



Phase 3 trial results



Safety profile



Case studies



Patient care and more

PLEASE JOIN PFIZER AND GENMAB FOR THIS LIVE EVENT!

This in-depth presentation is a great opportunity to learn more about Tivdak for your patients with previously treated recurrent or metastatic cervical cancer.

Healthcare providers, including medical oncologists, gynecologic oncologists, PAs/NPs, RNs, Pharm Ds, eye care providers, infusion nurses, and additional members of patient care teams, are all encouraged to participate.

SPEAKER

Scott Jordan, MD

Broward Health

DATE & TIME

Thursday, December 18, 2025

5:30 PM

LOCATION

36 Restaurant & Bar
36 North Main Street
Cape Girardeau, Missouri 63701
(573) 803-2333
Main Street

RSVP

Conni Stark: (314) 753-4856, csta@genmab.com

Kindly provide your full name, degree, specialty, address (including state), and email address when registering so you can be confirmed. Please also provide the Meeting ID# INT-0001951 when you RSVP.

To RSVP for this program, or for information regarding Tivdak, contact your local Account Manager or click here, <https://genmabprograms.my.salesforce-sites.com/Invitation/Register/INT-0001951>

INDICATION

TIVDAK is indicated for the treatment of adult patients with recurrent or metastatic cervical cancer (r/mCC) with disease progression on or after chemotherapy.

IMPORTANT SAFETY INFORMATION

BOXED WARNING: OCULAR TOXICITY

TIVDAK can cause severe ocular toxicities resulting in changes in vision, including severe vision loss, and corneal ulceration. Conduct an ophthalmic exam, including an assessment of ocular symptoms, visual acuity, and slit lamp exam of the anterior segment of the eye prior to initiation of TIVDAK, prior to every cycle for the first nine cycles, and as clinically indicated. Adhere to the required premedication and eye care before, during, and after infusion. Withhold TIVDAK until improvement and resume, reduce the dose, or permanently discontinue, based on severity.

Please see additional Important Safety Information on following pages and click for [full prescribing information](#), including **BOXED WARNING** for TIVDAK.

IMPORTANT SAFETY INFORMATION (continued)

Warnings and Precautions

Ocular adverse reactions: TIVDAK can cause severe ocular adverse reactions, including conjunctivitis, keratopathy (keratitis, punctate keratitis, and ulcerative keratitis), and dry eye (increased lacrimation, eye pain, eye discharge, pruritus, irritation, and foreign body sensation), that may lead to changes in vision and/or corneal ulceration.

Ocular adverse reactions occurred in 55% of patients with cervical cancer treated with TIVDAK across clinical trials. The most common were conjunctivitis (32%), dry eye (24%), keratopathy (17%), and blepharitis (5%). Grade 3 ocular adverse reactions occurred in 3.3% of patients, including severe ulcerative keratitis in 1.2% of patients. Nine patients (2.1%) experienced ulcerative keratitis (including one with perforation requiring corneal transplantation), six (1.4%) conjunctival ulcer, four (0.9%) corneal erosion, two (0.5%) conjunctival erosion, and two (0.5%) symblepharon.

In innovaTV 301, 8 patients (3.2%) experienced delayed ocular adverse reactions occurring more than 30 days after discontinuation of TIVDAK. These adverse reactions included 3 patients with ulcerative keratitis, and one patient (each) with keratitis, punctate keratitis and corneal erosion, blepharitis and conjunctival hyperemia, conjunctival scar, and conjunctivitis and xerophthalmia.

Refer patients to an eye care provider to conduct an ophthalmic exam prior to initiation of TIVDAK, prior to every cycle for the first nine cycles, and as clinically indicated. The exam should include visual acuity, slit lamp exam of the anterior segment of the eye, and an assessment of normal eye movement and ocular signs or symptoms which include dry or irritated eyes, eye secretions, or blurry vision.

Adhere to the required premedication and eye care before, during, and after infusion to reduce the risk of ocular adverse reactions. Monitor for ocular toxicity and promptly refer patients to an eye care provider for any new or worsening ocular signs and symptoms. Withhold, reduce, or permanently discontinue TIVDAK based on the severity or persistence of the ocular adverse reaction.

Peripheral neuropathy (PN) occurred in 39% of cervical cancer patients treated with TIVDAK across clinical trials; 6% of patients experienced Grade 3 PN. PN adverse reactions included peripheral sensory neuropathy (23%), PN (5%), paresthesia (3.8%), peripheral sensorimotor neuropathy (3.3%), muscular weakness (2.8%), and peripheral motor neuropathy (2.4%). One patient with another tumor type treated with TIVDAK at the recommended dose developed Guillain-Barre syndrome.

Monitor patients for signs and symptoms of neuropathy such as paresthesia, tingling or a burning sensation, neuropathic pain, muscle weakness, or dysesthesia. For new or worsening

PN, withhold, then dose reduce, or permanently discontinue TIVDAK based on the severity of PN.

Hemorrhage occurred in 51% of cervical cancer patients treated with TIVDAK across clinical trials. The most common all grade hemorrhage adverse reaction was epistaxis (33%). Grade 3 hemorrhage occurred in 4% of patients.

Monitor patients for signs and symptoms of hemorrhage. For patients experiencing pulmonary or central nervous system hemorrhage, permanently discontinue TIVDAK. For Grade ≥ 2 hemorrhage in any other location, withhold until bleeding has resolved, blood hemoglobin is stable, there is no bleeding diathesis that could increase the risk of continuing therapy, and there is no anatomical or pathologic condition that can increase the risk of hemorrhage recurrence. After resolution, either resume treatment or permanently discontinue TIVDAK.

Pneumonitis that is severe, life-threatening, or fatal can occur in patients treated with antibody-drug conjugates containing vedotin, including TIVDAK. Among cervical cancer patients treated with TIVDAK across clinical trials, 4 patients (0.9%) experienced pneumonitis, including 1 patient who had a fatal outcome.

Monitor patients for pulmonary symptoms of pneumonitis. Symptoms may include hypoxia, cough, dyspnea or interstitial infiltrates on radiologic exams. Infectious, neoplastic, and other causes for such symptoms should be excluded through appropriate investigations. Withhold TIVDAK for patients who develop persistent or recurrent Grade 2 pneumonitis and consider dose reduction. Permanently discontinue TIVDAK in all patients with Grade 3 or 4 pneumonitis.

Severe cutaneous adverse reactions (SCAR), including events of fatal or life-threatening Stevens-Johnson syndrome (SJS), can occur in patients treated with TIVDAK. SCAR occurred in 1.6% of cervical cancer patients treated with TIVDAK across clinical trials. Grade ≥ 3 SCAR occurred in 0.5% of patients, including 1 patient who had a fatal outcome.

Monitor patients for signs or symptoms of SCAR, which include target lesions, worsening skin reactions, blistering or peeling of the skin, painful sores in mouth, nose, throat, or genital area, fever or flu-like symptoms, and swollen lymph nodes. If signs or symptoms of SCAR occur, withhold TIVDAK until the etiology of the reaction has been determined. Early consultation with a specialist is recommended to ensure greater diagnostic accuracy and appropriate management. Permanently discontinue TIVDAK for confirmed Grade 3 or 4 SCAR, including SJS.

Embryo-fetal toxicity: TIVDAK can cause fetal harm when administered to a pregnant woman. Advise patients of the potential risk to a fetus. Advise females of reproductive potential to use effective contraception during treatment with TIVDAK and for 2 months after the last dose. Advise male

IMPORTANT SAFETY INFORMATION (continued)

Embryo-fetal toxicity (continued)

patients with female partners of reproductive potential to use effective contraception during treatment with TIVDAK and for 4 months after the last dose.

Adverse Reactions

Across clinical trials of TIVDAK in 425 patients with r/mCC, the most common ($\geq 25\%$) adverse reactions, including laboratory abnormalities, were hemoglobin decreased (45%), PN (39%), conjunctival adverse reactions (38%), nausea (37%), fatigue (36%), aspartate aminotransferase increased (33%), epistaxis (33%), alopecia (31%), alanine aminotransferase increased (30%), and hemorrhage (28%).

innovaTV 301 Study: 250 patients with r/mCC with disease progression on or after systemic therapy

Serious adverse reactions occurred in 33% of patients receiving TIVDAK; the most common ($\geq 2\%$) were urinary tract infection (4.8%), small intestinal obstruction (2.4%), sepsis, abdominal pain, and hemorrhage (each 2%). **Fatal adverse reactions** occurred in 1.6% of patients who received TIVDAK, including acute kidney injury, pneumonia, sepsis, and SJS (each 0.4%).

Adverse reactions leading to permanent discontinuation occurred in 15% of patients receiving TIVDAK; the most common ($\geq 3\%$) were PN and ocular adverse reactions (each 6%). **Adverse reactions leading to dose interruption** occurred in 39% of patients receiving TIVDAK; the most common ($\geq 3\%$) were ocular adverse reactions (16%) and PN (6%). **Adverse reactions leading to dose reduction** occurred in 30% of patients receiving TIVDAK; the most common ($\geq 3\%$) were PN and ocular adverse reactions (each 10%). The ocular adverse reactions included conjunctival disorders (4.8%), keratopathy (4%), and dry eye (0.8%).

innovaTV 204 Study: 101 patients with r/mCC with disease progression on or after chemotherapy

Serious adverse reactions occurred in 43% of patients; the most common ($\geq 3\%$) were ileus (6%), hemorrhage (5%), pneumonia (4%), PN, sepsis, constipation, and pyrexia (each 3%). **Fatal adverse reactions** occurred in 4% of patients who received TIVDAK, including septic shock, pneumonitis, sudden death, and multisystem organ failure (each 1%).

Adverse reactions leading to permanent discontinuation occurred in 13% of patients receiving TIVDAK; the most common ($\geq 3\%$) were PN (5%) and corneal adverse reactions (4%).

Adverse reactions leading to dose interruption occurred in 47% of patients; the most common ($\geq 3\%$) were PN (8%), conjunctival adverse reactions, and hemorrhage (each 4%).

Adverse reactions leading to dose reduction occurred in 23% of patients; the most common ($\geq 3\%$) were conjunctival adverse reactions (9%) and corneal adverse reactions (8%).

Drug Interactions

Strong CYP3A4 inhibitors: Concomitant use with strong CYP3A4 inhibitors may increase unconjugated monomethyl auristatin E (MMAE) exposure, which may increase the risk of TIVDAK adverse reactions. Closely monitor patients for TIVDAK adverse reactions.

Use in Specific Populations

Moderate or severe hepatic impairment: MMAE exposure and adverse reactions are increased. Avoid use.

Lactation: Advise lactating women not to breastfeed during TIVDAK treatment and for at least 3 weeks after the last dose.

Please see full Prescribing Information, including BOXED WARNING for TIVDAK.

Because late arrivals may be prohibited from participating in the program, please make every effort to arrive by the designated start time. You are expected to remain for the duration of the program.

Notice: Pfizer and Genmab are committed to the appropriate marketing of their products to ensure that interactions with HCPs are professional exchanges designed to benefit patients and enhance the practice of medicine. All Pfizer- and Genmab-sponsored programs are conducted in accordance with PhRMA Code principles to address a bona fide educational need with appropriate attendees in a venue and manner conducive to informational exchange.

State and Federal Employees: State and federal laws and regulations may carry additional restrictions. By attending this event, you confirm that you have obtained any necessary approvals from your employer.

Public Disclosures: Pfizer and Genmab report payment and transfer of value data as required by federal and state transparency laws.

State Law Restrictions:

Minnesota: Regardless of where you practice or reside, if you are a Minnesota-licensed practitioner with prescribing privileges, then you may not consume any food or beverage associated with this event.

Vermont: Regardless of where you practice or reside, if you are a Vermont-licensed HCP, you may not consume any food or beverage associated with this event. Additionally, if you are an employee/agent of a Vermont HCP (e.g., PAs, non-prescribing nurses, etc.), regardless of where you practice or reside, you may not consume any food or beverage associated with this event.

Pfizer and Genmab do not provide or pay for alcohol in connection with any program we sponsor.