

YOU ARE CORDIALLY INVITED TO ATTEND A SPEAKER PROGRAM

LOWER TOGETHER

PRESENTED BY

John Stein, MD

Chesterfield, MO

WHEN

6:00 PM Central

Thursday, January 17, 2019

WHERE

Celebrations Downtown

19 South Spanish Street
Cape Girardeau, MO 63703
(573) 334-8330

RSVP WITH YOUR AMGEN REPRESENTATIVE

Barbara Bergh

bergh@amgen.com
(314) 378-7922

INDICATIONS

Prevention of Cardiovascular Events: In adults with established cardiovascular disease, Repatha[®] is indicated to reduce the risk of myocardial infarction, stroke, and coronary revascularization.

IMPORTANT SAFETY INFORMATION

Contraindication: Repatha[®] is contraindicated in patients with a history of a serious hypersensitivity reaction to Repatha[®]. Serious hypersensitivity reactions including angioedema have occurred in patients treated with Repatha[®].

Please see additional Important Safety Information on the next page.

Notice: This event is conducted in accordance with the PhRMA Code on Interaction with Healthcare Professionals and is limited to invited healthcare professionals. Attendance by guests or spouse is not appropriate. Government employees are subject to state and federal laws and ethics rules that may limit your ability to receive any gifts, including meals, from pharmaceutical companies. If you are a state or federal employee, it is your responsibility to seek guidance and prior approval from your employer or site ethics counselor to attend this event.

State Laws: To comply with law and Amgen policies, Amgen is unable to offer food and beverages to (1) individuals with prescribing authority in Vermont and Minnesota; and (2) individuals employed by prescribers in Vermont who support the provision of healthcare. Please note that Amgen exercises diligence in reviewing the licensure of attendees and asks that you cooperate by disclosing all licensures in the sign in/registration process. We appreciate your understanding and support.

Disclosure by Amgen: Amgen reports payments and transfers of value made to healthcare professionals and other healthcare related entities in accordance with federal and state laws, regulations and other transparency obligations. Any items of value provided by Amgen at this event (including the provision of meals and refreshments) may be subject to public disclosure. If you have questions regarding this matter please contact Amgen at 805-447-7422 or HCCSpendInquiry@amgen.com.

IMPORTANT SAFETY INFORMATION

Contraindication: Repatha® is contraindicated in patients with a history of a serious hypersensitivity reaction to Repatha®. Serious hypersensitivity reactions including angioedema have occurred in patients treated with Repatha®.

Allergic Reactions: Hypersensitivity reactions (e.g. angioedema, rash, urticaria) have been reported in patients treated with Repatha®, including some that led to discontinuation of therapy. If signs or symptoms of serious allergic reactions occur, discontinue treatment with Repatha®, treat according to the standard of care, and monitor until signs and symptoms resolve.

Adverse Reactions in Primary Hyperlipidemia (including HeFH): The most common adverse reactions (>5% of patients treated with Repatha® and occurring more frequently than placebo) were: nasopharyngitis, upper respiratory tract infection, influenza, back pain, and injection site reactions. From a pool of the 52-week trial and seven 12-week trials: Local injection site reactions occurred in 3.2% and 3.0% of Repatha®-treated and placebo-treated patients, respectively. The most common injection site reactions were erythema, pain, and bruising.

Allergic reactions occurred in 5.1% and 4.7% of Repatha®-treated and placebo-treated patients, respectively. The most common allergic reactions were rash (1.0% versus 0.5% for Repatha® and placebo, respectively), eczema (0.4% versus 0.2%), erythema (0.4% versus 0.2%), and urticaria (0.4% versus 0.1%).

Adverse Reactions in the Cardiovascular Outcomes Trial: The most common adverse reactions (>5% of patients treated with Repatha® and occurring more frequently than placebo) were: diabetes mellitus (8.8% Repatha®, 8.2% placebo), nasopharyngitis (7.8% Repatha®, 7.4% placebo), and upper respiratory tract infection (5.1% Repatha®, 4.8% placebo).

Among the 16,676 patients without diabetes mellitus at baseline, the incidence of new-onset diabetes mellitus during the trial was 8.1% in patients assigned to Repatha® compared with 7.7% in those assigned to placebo.

Immunogenicity: Repatha® is a human monoclonal antibody. As with all therapeutic proteins, there is potential for immunogenicity with Repatha®.

Please see full Prescribing Information.

INDICATIONS

Prevention of Cardiovascular Events: In adults with established cardiovascular disease, Repatha® is indicated to reduce the risk of myocardial infarction, stroke, and coronary revascularization.

Primary Hyperlipidemia (including Heterozygous Familial Hypercholesterolemia): Repatha® is indicated as an adjunct to diet, alone or in combination with other lipid-lowering therapies (e.g., statins, ezetimibe), for the treatment of adults with primary hyperlipidemia to reduce low-density lipoprotein cholesterol (LDL-C).