

CROHN'S & ULCERATIVE COLITIS REFERRAL FORM

LOTUS HEALTH INFUSIONS
Fax Referral To: 877-643-0993
Phone: 832-284-4452



Date: _____

Needs by Date: _____ Ship to Patient's Home Prescriber 1st Order Only Prescriber All Orders

PATIENT INFORMATION

Patient Name: _____
Address: _____
City, State, Zip: _____
Home Phone: _____
Cell Phone: _____
Date of Birth: _____ Gender: M F

PRESCRIBER INFORMATION

Prescriber Name: _____
Address: _____
City, State, Zip: _____
Phone: _____ Fax: _____
DEA#: _____ NPI#: _____
Contact Person: _____

INSURANCE INFORMATION (Please attach the front and back of insurance and prescription drug card)

Primary Insurance: _____ ID#: _____ Group: _____
Secondary Insurance: _____ ID#: _____ Group: _____
Prescription Card: _____ ID#: _____ BIN: _____ PCN: _____ Group: _____

DIAGNOSIS & CLINICAL ASSESSMENT (Fill in below or attach lab work)

New to Therapy Currently on Therapy | Start Date: _____ Physician Provides Injection Training | Injection Date: _____
Primary Diagnosis Code & Condition: _____ **Date of Diagnosis:** _____
TB Test Results & Date: _____ **Current Weight:** _____ **Date:** _____ **Allergies:** _____
 New Therapy Induction Therapy Change Remicade Therapy Continuation, Weeks Completed: 0 2 4 6 Date: _____
 Inadequate Response to Methotrexate (Dose: _____) Unresponsive to Conventional Treatment, Other Therapies: _____

Cimzia® (certolizumab pegol)

Starter Kit (6) 200mg Prefilled Syringes
 2 x 200mg Vials
 2 x 200mg Prefilled Syringes
Dose / Directions / Frequency:
 Induction Dose: 2 x 200mg injections at Week 0, 2 and 4
 Maintenance Dose: 400 mg s-c monthly
 Other: _____
QTY: _____ Refill: _____

Entyvio® (vedolizumab)

300 mg Vial
Dose / Directions / Frequency:
 Induction Dose: 300mg IV at wk 0, 2 & 6
 Maintenance Dose: 300mg IV every 8 wks
 Other: _____
QTY: _____ Refill: _____

Humira® (adalimumab)

Crohn's Starter Kit, 6 x 40mg pens
 Pediatric Crohn's Starter Kit, 3 x 40mg PFS
 40mg Pens 40 mg PFS
 20mg pediatric PFS 10mg pediatric PFS
Dose / Directions / Frequency:
 Induction Dose: Adults & Children >= 88lbs; 160mg (4 x 40mg injections in one day or 2 x 40mg injections per day for two consecutive days); Second dose two weeks later (Day 15) 80mg
 Induction dose: Children < 88lbs; 80mg (2 x 40mg injections in one day) Second dose two weeks later (Day 15) 40mg
 Maintenance: _____mg every other week
 Other: _____
QTY: _____ Refill: _____
Step Therapies: Therapy tried and failed
Therapy: _____ Date: _____
Therapy: _____ Date: _____
Therapy: _____ Date: _____

Stelara® (ustekinumab)

2 x 130mg/26mL 3 x 130mg/26mL
 4 x 130mg/26mL 1 x 90mg/mL PFS
Dose / Directions / Frequency:
 Infuse 260mg intravenously over no less than one hour (<55kg)
 Infuse 390mg intravenously over no less than one hour (55kg to 85kg)
 Infuse 520mg intravenously over no less than one hour (>85kg)
 Inject 90mg SQ 8 weeks post-initial IV dose, then q 8 weeks thereafter
QTY: _____ Refill: _____

Simponi®

Auto Injection: _____ 50mg _____ 100mg
 Syringe: _____ 50mg _____ 100mg

Dose / Directions / Frequency:
 Induction Dose: 200mg s-c initially then 100mg 2 weeks later
 Maintenance Dose: 100mg s-c Q 4 wks
 Other: _____
QTY: _____ Refill: _____

Remicade® (infliximab)

100 mg Vial **SIG:** _____
QTY: _____ Refill: _____

Other/Notes: _____

Prescriber Signature: _____ **DAW (Dispense as Written)** Y N **Date:** _____

The information contained in this facsimile may be confidential and is intended solely for the use of the named recipient(s). Access, copying or re-use of the facsimile or any information contained therein by any other person is not authorized. If you are not the intend recipient, please notify us immediately by faxing back to the originator.