IGIV and General Immune Disorders Enrollment Form Date:	Enrollment Form Fax Refer		_	
PATIENT INFORMATION Patient Name: Address: City, State, Zip: Home Phone: Cell Phone: Alternate Phone: Date of Birth:		Prescriber Name: Address: City, State, Zip: Phone: Fax: DEA#:		RIBER INFORMATION
				uranaa and proceedintion drug aard)
Primary Insurance:	#:	ID#:		urance and prescription drug card)Group: Group:Group:
Neurological G61.81 Chronic Inflammatory Demyelinating Per G61.82 Multifocal Motor Neuropathy (MMN) G61.0 Guillain-Barre G35 Multiple Sclerosis G70.01 Myasthenia Gravis w/Exacerbation Other:	Immunological Primary Immune Deficiency – Please specify ICD-10 Code: D80.9 Deficiency of Humoral Immunity D83.9 Common Variable Immunodeficiency D89.9 Immune Mechanism Disorder D81.9 Immune Deficiency NOS D69.3 Idiopathic Thrombocytopenia D80.1 Hypogammaglobulinemia Other:			
CLINICAL INFORMATION (Please attach all clinical information, lab results, and other medical history documents) Patient Weight: Kg/Lbs Height: Inches/CM Allergies: Inches/CM Height: Meds by Date: Inches/CM Has patient previously received IVIG Yes No Line Access: PIV PICC PORT Needs by Date:				
Intravenous □ Gammagard®Liq.10% □ Privigen® 10% □ Gamunex-C®10% □ Octagam® 5% □ Gammaked®10% □ Octagam® 10% □ Bivigam® □	(Pharmacy to round	Rday(s); Ever		Directions Infuse total dose of Immunoglobulin intravenously based on manufacturer recommend infusion rate as tolerated. Infuse via:
□ Gammagard® S/D □ Other: □ Panzyga® 10%	□ 1 month □ 3 months □ 6 months □ 12 months □ Other			Infusion Pump Gravity Excludes Medicare D
Medication Subcutaneous Gammagard® Liq. 10%	Dose grams ORgram(s) per kg (Pharmacy to round to nearest vial size) Infuse total dose OVERday(s); Every week(s) for: 1 month 3 months 6 months 12 months Other		ery	Directions Infuse total dose of Immunoglobulin subcutaneously in one or more infusion sites via infusion pump based on manufacturer recommend infusion rate as tolerated. Other:
Premedication to be given 30 minutes prior to infusion: □ Diphenhydramine 25-50 mg po – 25mg #2 per dose □ Diphenhydramine 25-50 Slow IV-Push – 50mg vial #1 per dose □ Acetaminophen 325-650 mg po – 325mg #2 per dose □ Ketorolac 30mg Slow IV-Push – 30mg/ml vial #1 per dose □ Other:		IV Access Flush Order / EpiPen® Order: (Infusion supplies per pharmacy protocol) NaCl 0.9% 5-10ml IV before and after infusion Heparin 10 units/ml 3-5ml IV after infusion for peripheral access and PRN Heparin 100 units/ml 3-5ml IV after infusion for central IV access and PRN All infusion supplies necessary to administer the medication EpiPen® 0.3mg auto-injector for severe anaphylactic reaction for patient weighing ≥ 30kg. EpiPen Jr. ® 0.15mg for patients weighing under 30kg		
By signing below, I certify that above therapy is medically necessary. Prescriber's Signature (SIGN BELOW)				
Dispense as Written Da	ite	Substitutio	onAllowed	Date

The information contained in this facsimile may be confidential and is intended solely for the use of the named recipient(s). Access, copying or re-use of the facsimile or any information contained therein by any other person is not authorized. If you are not the intend recipient, please notify us immediately.