## 8. RECEIPT BY PANJRAPOLE (INFIRMARY) RECEIVING ANIMALS

Police Station:			Po	lice Station				
CR No		Da	ited					
Total	slaughter police we	house, ere taker of the ur	were loan to	nded in a t	truck No Panjai	capole and	were kep	llegally to a and through t therein in als has been
Sr. No.	Tag No. of the Animal affixed on the ear.	Sex of the animal	Age of the animal Yrs.	Colour of the animal	Infection etc	Condition	Primary treat- ment	Medical treat- ment
1	MP1	Male	1 1/2	Whitish	Skin bruished	Front and rear legs are tired with rope in a manner that neck of all the calves remain in between.	FMD	P.C. Genticine
2	MP2	Male	2 1/4	Black	Injury on leg.			Antict onetice
3	MP3	Male	1 1/4	White	Aafaro i.e. windness in stomach		1	Avil
4	MP4	Male	1 1/4	White	Legs bruished			P.P.
5	MP5	Male	1 ½	Whitish	Aafaro			Templine old
6	MP6	Male	2	Whitish	Wound on the			Jink Oxide powder

leg

7	MP7	Male	1 1/2	Black	Blood ozzing from no strails		Pexona
8	MP8	Male	1	Black	Skin bruished		
9	MP9	Male	1 ½	White	Blood coming from mouth		FMD Vaccin- ation
10	MP 10	Male	1 ½	Reddish	One eye lost		
11	MP 11	Male	2	Reddish	Blood from urine		
12	MP 12	Male	1	Black	Skin Bruished		
13	MP 13	Male	2	Black	Wound on the leg		
14	MP 14	Male	1	Whitish	Skin bruished		
15	MP 15	Male	2	Whitish	Aafaro		
16	MP 16	Male	1	Whitish	Loose motion		
17	MP 17	Male	1 ½	Black	Loose motion		
18	MP 18	Male	1	Whitish	Blood from No strials		
19	MP 19	Male	2	Reddish	Skin bruished.		

## **Over** and above that:-

- (1) Aforesaid calves were found tied tightly with the rope.
- (2) The age of the aforesaid calves is found between 1 to 2 years.
- (3) The aforesaid calves can be used for agricultural operations after about 2 years.
- (4) The physical strength of the aforesaid calves is found to be very weak. Majority of them are found having skin infection.
- (5) The symptoms of FMD (Khar Movasa) are found in aforesaid animals.
- (6) No arrangements worth the name of sand, water, fodder for the animals were found in the said truck.