



## SculpSure

### Contraindications

As with any procedure, it is the physician's responsibility to determine whether SculpSure is an appropriate treatment option for their clients.

Treatment using the SculpSure system is contraindicated in any client who has:

- **Open lesions and wounds. Treatment should only be applied to intact, healthy skin, with no evidence of compromised wound healing.**
- **An unrepaired abdominal hernia.**
- **Had prior treatment with parenteral gold therapy (gold sodium thiomalate).**
- **Is pregnant and/or breastfeeding.**

### Cautionary Criteria

Treatment using the SculpSure system is cautioned in any client who:

- **Has a history of immunosuppression / immune deficiency disorders, including HIV infection or AIDS, or use of immunosuppressive medications, including Advil and Aleve.**
- **Is currently taking antiplatelets, thrombolytics, anti-inflammatories or anticoagulants- including but not limited to the heavy use of aspirin, i.e., greater than 81 mg per day.**
- **Has a coagulation disorder.**
- **Has had a previous procedure or surgery in the treatment area that may have resulted in decreased sensation, scar/fibrotic tissue or large scars that may interfere with the treatment.**
- **Has an active tan, or the need to be exposed to artificial tanning devices or excessive sunlight a week (7 days) before or after treatment.**
- **Has a tattoo in the treatment area.**
- **Has a history of skin photosensitivity disorders, or is taking photo-sensitized medication.**
- **Has a neuropathic disorder, impaired skin sensation, or diabetic neuropathy.**
- **Has a history of keloid formation.**
- **Has a history or evidence of squamous cell carcinoma or melanoma.**