

When Do Children and Teens Need Vaccinations?

Age	HepB Hepatitis B	DTaP/Tdap Diphtheria, tetanus, pertussis (whooping cough)	Hib <i>Haemophilus influenzae</i> type b	IPV Polio	PCV Pneumo- coccal conjugate	RV Rotavirus	MMR Measles, mumps, rubella	Vari- cella Chickenpox	HepA Hepatitis A	COVID-19	Dengue	HPV Human papillomavirus	Men- ACWY		MenB	Influenza Flu
													Meningococcal			
at Birth (within 24 hours of birth)	✓															
2 months	✓	✓	✓	✓	✓	✓										
4 months	✓ ¹	✓	✓	✓	✓	✓										
6 months	✓ (6–18 mos)	✓	✓ ¹	✓ (6–18 mos)	✓	✓ ¹										✓ (6 mos and older)
12 months			✓ (12–15 mos)		✓ (12–15 mos)		✓ (12–15 mos)	✓ (12–15 mos)	✓✓ (2 doses given 6 months apart routinely at age 12–23 months)							
15 months		✓ ² (15–18 mos)														
18 months																
19–23 months									HepA vaccine (2 doses) is also recommended for children and teens not previously vaccinated							
4–6 years		✓		✓			✓	✓								
7–10 years										✓ ³ As of 5/1/22, COVID-19 vaccine is recom- mended for everyone age 5 yrs and older						
11–12 years		✓ (Tdap)									✓✓✓✓ ⁴ Dengue vaccine is recom- mended for certain children	✓✓ ^{5,6}	✓			
13–15 years																
16–18 years													✓	✓✓ ^{7,8}		Influenza vaccine is recommended every year for everyone age 6 months and older

One dose each fall or winter to all people ages 6 mos and older. Some children younger than age 9 years need 2 doses; ask your child's health-care provider if your child needs more than 1 dose.

- NOTES**
- 1 Your child may not need this dose depending on the brand of vaccine that your healthcare provider uses.
 - 2 This dose of DTaP may be given as early as age 12 months if it has been 6 months since the previous dose.
 - 3 The number of doses and possible booster is determined by the vaccine brand given.

- 4 Children ages 9 through 16 years living in Puerto Rico, American Samoa, U.S. Virgin Islands, Federated States of Micronesia, Republic of Marshall Islands, and the Republic of Palau, and have lab-confirmed previous dengue infection are recommended to receive a 3-dose series of Dengue vaccine.
- 5 HPV vaccine is routinely given at age 11 or 12 years but may be started at age 9.

- 6 Children with certain medical conditions will need a third dose.
- 7 This vaccine may be given to healthy teens. It is also recommended for adolescents with certain health conditions.
- 8 Your teen may need an additional dose depending on your healthcare provider's recommendation.