



EMERGENCY MEDICATION

NAME: ADRENALINE (EPINEPHRINE) EPIPEN

How to USE: Must be injected into thigh muscle If necessary repeat doses.

When to USE: Used for emergency treatment of severe allergic reaction

NAME: CHLORPHENIRAMINE

HOW to USE: Swallow your chlorphenamine tablets with a drink of water.

WHEN to USE: It is used to relieve allergies, drug allergies, and allergic skin reactions, and to relieve itching caused by infections.

NAME: DEXTROGEL

How to USE: Following blood glucose monitor when the level is under 4. Refer to the glucose machine and range.

WHEN to USE: Squeeze the tube to insert the gel into the person's mouth between their teeth and their cheek. Rub the outside of their cheek gently to allow the gel to become absorbed. It is absorbed through the lining of the mouth and should help raise the blood glucose within 15 minutes.

NAME: DEXAMETHASONE

HOW to USE: Inject intramuscular 3.3mgs- every 30-60 minutes until controlled.

Initial dose repeat after 30 minutes.

WHEN to USE: Dexamethasone is a corticosteroid. It is used to treat inflammation of the skin, joints, lungs, and other organs. Common conditions treated include asthma, allergies.

NAME: DISPERSIBLE ASPIRIN

HOW to USE: 75mg tablets are indicated for the secondary prevention of thrombotic cerebrovascular or cardiovascular disease.

WHEN to USE Chest Pain - For oral administration. The tablets should be dispersed in water before taken.

NAME: GLYCERYL TRINITRATE

Use it when chest pain occurs (angina)

HOW to USE: 400-800 micrograms to be administered under the tongue (2 sprays) when chest pain occurs and then close mouth.

WHEN to USE: When chest pain is an acute episode

.NAME: GLUCAGEN HYPOKIT

HOW to USE: Following blood glucose monitor when the level is under 4. Refer to the glucose machine and range. Choose the injection site. Common injection sites for GlucaGen® are upper arms, thighs, or buttocks. With one hand, gently pinch the skin at the injection site. With your other hand, insert the needle into the skin and push the syringe plunger down until the syringe is empty.

WHEN to USE: GlucaGen® HypoKit is used to treat severe hypoglycaemia (extremely low blood sugar levels, or "hypos' ') in people using insulin or taking tablets to control their diabetes, who have become unconscious.

NAME: HYALURONIDASE

WHEN to USE:

- 1) Vascular occlusion
- 2) Blindness
- 3) Tyndall effect
- 4) Unacceptable cosmetic outcome
- 5) Delayed onset nodules.
- 6) Allergic or Immunogenic reaction to the HA dermal filler

HOW to USE:

Reconstitute the hyaluronidase 1500 IU with 1ml of normal saline or water for injection. Rotate the vial to ensure the powder is fully dissolved - inject carefully within the area affected

NAME: PARACETAMOL

HOW to USE: Take paracetamol with water. Paracetamol can be taken with or without food. The usual dose for adults is one or two 500mg tablets up to 4 times in 24 hours.

WHEN to USE: Paracetamol is a commonly used medicine that can help treat pain and reduce a high temperature (fever). It's typically used to relieve mild or moderate pain, such as headaches.

NAME: VENTOLIN

HOW to USE: Hold your breath for about 10 seconds after breathing in the medicine. Then breathe out fully.

WHEN to USE: Ventolin is used to treat or prevent bronchospasm, or narrowing of the airways in the lungs, in people with asthma or certain types of chronic obstructive pulmonary disease (COPD).

DEFRILLATOR MACHINE -outside of BIC RECEPTION