

RANCHER'S CHOICE®

COOKED MOLASSES PROTEIN TUB 30% PROTEIN

For Beef Cattle on Pasture or Roughage Rations

GUARANTEED ANALYSIS:

Crude Protein, Min.....	30.0%
(This includes not more than 15% equivalent crude protein from non-protein nitrogen.)	
Crude Fat, Min.....	4.0%
Crude Fiber, Max.....	1.5%
Calcium (Ca), Min.....	1.6%
Calcium (Ca), Max.....	2.1%
Phosphorus (P), Min.....	1.0%
Magnesium (Mg), Min.....	1.0%
Potassium (K), Min.....	3.0%
Cobalt (Co), Min.....	3 ppm
Copper (Cu), Min.....	300 ppm
Iodine (I), Min.....	50 ppm
Manganese (Mn), Min.....	1,200 ppm
Selenium (Se), Min.....	6.6 ppm
Zinc (Zn), Min.....	1,200 ppm
Vitamin A, Min.....	82,000 IU/lb
Vitamin D ₃ , Min.....	14,000 IU/lb
Vitamin E, Min.....	82 IU/lb

INGREDIENTS:

Molasses Products, Plant Protein Products, Urea, Processed Grain By-Products, Monocalcium Phosphate, Vegetable Oil, Calcium Carbonate, Magnesium Oxide, Manganese Sulfate, Zinc Sulfate, Copper Sulfate, Vitamin A Supplement, Vitamin E Supplement, Sodium Selenite, Ethylenediamine Dihydroiodide, Vitamin D₃ Supplement and Cobalt Carbonate.

FEEDING DIRECTIONS:

Feed RANCHER'S CHOICE® 30% Protein Cooked Molasses Tub free-choice to cattle on pasture or roughage rations. Make sure animals have adequate pasture or roughage available and are not starved for salt or minerals. Feed continuously. Recommended consumption of RANCHER'S CHOICE® 30% Protein Cooked Molasses Tub is 0.75 to 1.00 pounds per head per day. Management of the number and location of the tubs by the producer may be required to achieve target intakes. Increase the number of tubs and move tubs closer to water or loang areas to increase intake. Decrease the number of tubs or move farther from water or loang areas to decrease intake. Provide at least one tub for every 15 to 20 animals. Provide free access to fresh, clean water at all times.

CAUTION:

Contains copper. Do not feed to sheep. Feeding selenium at levels in excess of 0.3 ppm in the total diet is prohibited. This feed contains a source of non-protein nitrogen. The equivalent protein supplied by non-protein nitrogen should not exceed 1/3 of the crude protein of the total ration. Use only as directed.