

**MESOTHERAPY “LIPO ‘FAT’ DISSOLVE” TREATMENT
PATIENT PRE- & POST-TREATMENT INSTRUCTIONS**

A few simple guidelines before and after your treatment can make a difference

Pre-Treatment

1. If you have a special event or vacation coming up, please keep in mind that you may want to schedule your treatment at least two weeks in advance. Swelling will occur. It is the body's natural, healthy response to go through an inflammation process to remove the fat cells from the body.
2. It is recommended to discontinue the use of aspirin, NSAIDS (Ibuprofen/Advil), Aleve, Motrin, etc.), fish/flax oil or any other blood thinning supplements one week before treatment to minimize bruising or bleeding. Please consult with your primary care provider prior to discontinuing any prescribed blood thinning medications.
3. Avoid alcohol, caffeine, niacin supplement, high-sodium foods, high sugar foods, refined carbohydrates, and spicy foods 24-48 hours before and after your treatment. These items may contribute to increased swelling or irritation.
4. If you develop a cold/flu, infection, blemish, or rash, etc. in the area to be treated prior to your appointment, we recommend that you please reschedule your appointment until it resolves.
5. We will not perform mesotherapy treatments on patients with current or past history of dysphagia “trouble swallowing”, currently pregnant or breastfeeding, with presence of infection at or near the injection sites, or under the age of 18.
6. We will take careful consideration in performing mesotherapy treatments on patients with excessive skin laxity, prominent platysmal bands, enlarged thyroid, enlarged neck lymph nodes, pronounced submandibular glands, prior surgery or aesthetic treatment to the treatment area, presence of scar tissue in the treatment area, sunburned or irritated skin, and current use of blood thinning agents.
7. We will assist and advise in coordinating mesotherapy treatments when being treated with other aesthetic services to provide optimal results and efficacy of concurrent treatments.
8. On the morning of your treatment, wash the area to be treated, DO NOT apply any lotions, creams, oils or sprays on the area being treated. Eat a light, high-protein meal 2-4hrs before your appointment.

Post-Treatment

1. You can expect to have swelling, redness, itchiness in the treatment area for the first few days.
2. Place a cold compress or ice pack on the area for 20 minutes and remove the cold compress or ice pack for 20 minutes. Continue this pattern as needed for 24 hours.
3. Continue to use Arnica to help decrease bruising, swelling, and discomfort for another 2-3 days.
4. Take (over the counter) acetaminophen (i.e. Tylenol) if needed to decrease post treatment discomfort.
5. Sleep on your back and with head elevated for the next 3-5 days after treatment (if submental area treated).
6. Drink plenty of water (72oz/2L) after treatment and for several days post treatment to help stay hydrated and flush out by-products.
7. Avoid hot baths or showers for at least 48hrs post treatment.
8. Avoid vigorous exercise, sun and heat exposure for 3-5 days after treatment. Always use SPF 30 or higher afterwards to protect the treated area, especially in the first 2 weeks after your treatment.
9. Avoid steroids (prednisone), NSAIDS (Ibuprofen/Advil, Motrin, Aleve, etc), alcohol, caffeine, niacin supplement, high-sodium foods, high sugar foods, refined carbohydrates, and spicy foods 24-48 hours after your treatment.
10. Please contact us immediately if you develop an asymmetric smile or facial muscle weakness, skin ulceration in the treatment area, difficulty swallowing, or if any existing symptom worsens. These are very rare.

Please call our clinic should you have any questions or concerns regarding your treatment
or aftercare (705-522-2252)