Ending the threat of HIV in Papua New Guinea.

23rd November 2022

Percy Pokeya
HIV Team Lead USAID

Milestones of HIV in PNG!

- 1987 First Case of HIV diagnosed in PNG.
- 1997 National Aids Counsel Secretariat established by act of Parliament.
- 2005 Antiretroviral Treatment first introduced into PNG.
- 2017 First national HIV Drug Resistance Survey which showed that PNG had high rates of HIV drug resistance (5th Highest according WHO 2020 HIV drug resistance report)
- 2019 PNG introduced optimum ART regimen (Tenofovir/Lamivudine/Dolutegravir TLD once daily pill)
- We are all in this together.
- We have come a long Way.
- We have a long way to go.
- We need to accelerate.




2021 HIV statistics of concern for PNG

- 13800 new infections. (45% increase since 2010)
- 59000 Total PLHIV
- HIV Vertical Transmission 36%
- 16% HIV Testing at ANC


https://www.unaids.org/en/resources/documents/2022/in-danger-global-aids-update

National 95-95-95 Progress, 2021 Estimates



Source: National Spectrum Estimates, 2021, VL data – HPDB September 2022

National Capital District 95-95-95 Progress, September 2022



Opportunities across the Cascade.

HIV Testing:

- We need to focus on optimizing case finding partner testing.
- We need to increase HIV testing of more importance is in ANC and TB clinics.
- Advocacy for HIV testing and increased provision of HIV testing.
- How can I advocate to help increase HIV testing.

HIV Treatment:


- Need to improve treatment continuity.
- Strengthen programs that reduce the number of people dropping out of Care and interrupting treatment.
- We commenced 53,248 PLHIV on ART, currently we have only 38,376 on ART, what happened to the 14,872 PLHIV who are Lost to follow up?
- How can I help improve treatment continuity?

• We need to improve viral load testing coverage.

- <u>U=U</u> messaging (Undetectable = Untransmittable), ie, if your viral load is undetectable after 12 months of continuous treatment, and you remain adherent to treatment, you will not transmit HIV.
- Achieving viral suppression has 2 important outcomes
- Prevents Transmission of HIV and patient has good clinical outcome.
- Prevents HIV drug resistance
- How can I amplify the U=U messaging?

HIV viral load testing:

HIV Prevention Care and Treatment Continuum



 The goal of antiretroviral therapy (ART) is to achieve virological suppression – <u>U=U</u>

Enablers

Improved functional information systems.

- •Improved HIV data information systems will help
- •Monitor the HIV epidemic
- •Improve ART procurement and distribution
- Advocate for resource allocation.

Strengthen CLM

- •Strengthen Community-Led monitoring to provide opportunity for users of services to be involved in improving services and generate demand for services amongst the communities. Right now KPAC made of 7 groups is leading CLM activities.
- •How can I help support CLM?


Improved Economy


- •Improved Economy with increased allocation of resources to address the HIV situation in PNG.
- •Improved HIV services including Human resources.
- •Increase in employment opportunities.
- •Improved livelihood of citizens.

Normalize HIV

•Reducing stigma and discrimination is critical to achieving HIV epidemic control.

Prevention toolbox





Strategic Documents

National Health Plan 2021-2030

National HIV/STI Strategy 2023-2027


Elimination of Mother to Child transmission of HIV/Syphilis/Hepatitis B.

Viral Load testing road map.

Do we have what it takes to win the race?



We can do this!



Thank you for listening!