Please check the examination detail	s below	before ente	ring your can	didate information
Candidate surname			Other name	s
Pearson Edexcel Level 1/Level 2 GCSE (9–1)	Centre	Number		Candidate Number
Morning (Time: 1 hour 45 minutes)	Paper R	eference 1	CH0/1H
Chemistry Paper 1				
				Higher Tier
You must have: Calculator, ruler				Total Marks

Instructions

- Use **black** ink or ball-point pen.
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions.
- Answer the questions in the spaces provided
 there may be more space than you need.
- Calculators may be used.
- Any diagrams may NOT be accurately drawn, unless otherwise indicated.
- You must show all your working out with your answer clearly identified at the end of your solution.

Information

- The total mark for this paper is 100.
- The marks for each question are shown in brackets
 - use this as a guide as to how much time to spend on each question.
- In questions marked with an **asterisk** (*), marks will be awarded for your ability to structure your answer logically showing how the points that you make are related or follow on from each other where appropriate.
- There is a periodic table on the back cover of the paper.

Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.

Turn over ▶







(2)

Answer ALL questions. Write your answers in the spaces provided.

Some questions must be answered with a cross in a box \boxtimes . If you change your mind about an answer, put a line through the box \boxtimes and then mark your new answer with a cross \boxtimes .

- 1 Alloys of gold are often used to make jewellery. The purity of gold is measured in carats. Different alloys of gold have different carats.
 - (a) Figure 1 shows the percentage of different metals in two samples of gold.

	percentage of metal						
	gold	silver	copper				
18 carat gold	75.0	15.0	10.0				
24 carat gold	100.0	0.0	0.0				

Figure 1

Explain why 18 carat gold is stronger than 24 carat gold.

You may use diagrams to help your answer.

(b) Figure 2 shows the relationship between the purity of gold in carats and the percentage of gold in the alloy.

in carats




Figure 2

A necklace with a mass of 5.0 g was found to contain 2.9 g of gold.

Determine the purity of the gold necklace in carats. Show your working.

purity of the gold necklace = carats

(3)

(c) A gold ring contains 3.94g of gold.

Calculate the number of gold atoms in the ring. (relative atomic mass: Au = 197, Avogadro constant = 6.02×10^{23})

Show your working.

(2)

number of gold atoms =

(Total for Question 1 = 7 marks)

BLANK PAGE



2 Figure 3 shows the apparatus that can be used to electrolyse sodium sulfate solution using inert electrodes.




Figure 3

- (a) Hydrogen is produced at the negative electrode during electrolysis.
 - (i) Describe the test to show the gas is hydrogen.

(2)

(ii) What is the name of gas **X** that forms at the positive electrode?

(1)

- 🛛 A ammonia
- B oxygen
- C nitrogen
- D sulfur dioxide
- (iii) State what is meant by the term **electrolysis**.

(2)

(b) The sodium sulfate solution was made by dissolving 28.4 g of sodium sulfate in water to make 250 cm³ of solution.	
Calculate the concentration of this solution in $g dm^{-3}$.	
Give your answer to three significant figures.	(3)
concentration =	gdm ⁻³
(c) The ions present in sodium sulfate are	
sodium Na ⁺ sulfate SO ₄ ²⁻	
Write the formula of sodium sulfate using this information.	(1)
(Total for Question 2 = 9 r	marks)

3 The word equation for the reaction between copper carbonate and dilute sulfuric acid is

copper + sulfuric
$$\rightarrow$$
 copper + carbon
carbonate + acid \rightarrow sulfate + dioxide + wate

(a) (i) Complete the balanced equation for this reaction.

(2)

$$CuCO_3 + \dots + CO_2 + H_2O$$

(ii) Calculate the relative formula mass of copper carbonate, $CuCO_3$. (relative atomic masses: C = 12.0, O = 16.0, Cu = 63.5)

(2)

relative formula mass of CuCO₃ =

(iii) What is the chemical test to show that a gas is carbon dioxide?

(1)

- A bubble the gas through limewater, limewater turns cloudy
- **B** put damp blue litmus paper in the gas, litmus paper turns red
- C put a lighted splint into the gas, the splint is extinguished
- \square **D** measure the pH of the gas, pH = 4

(b) Figure 4 shows a conical flask containing dilute sulfuric acid. Copper carbonate is added to the acid in the flask. The copper carbonate is added one spatula measure at a time until the reaction has finished.




Figure 4

State **two** observations that would show the reaction has finished.

(2)

 																					••••			
(c)	Th	ne	ele	ect	trc	ni	ic	co	n	fig	ju	ra	tie	or	۱ (of	ca	arl	oc	n	is	2.	.4

The electronic configuration of oxygen is 2.6

Draw a dot and cross diagram for a molecule of carbon dioxide.

Show outer electrons only.

(2)

(Total for Question 3 = 9 marks)



BLANK PAGE

4 Potassium hydroxide reacts with hydrochloric acid to form potassium chloride and water.

potassium hydroxide + hydrochloric acid → potassium chloride + water

(a) A student carried out a titration to find the exact volume of dilute hydrochloric acid that reacted with 25.0 cm³ of potassium hydroxide solution.

There were five steps in the titration.

The steps shown are not in the correct order.

- **step J** pour the potassium hydroxide solution into a conical flask and add a few drops of indicator to this solution
- **step K** fill a burette with the dilute hydrochloric acid and record the initial reading from the burette
- **step L** use a measuring cylinder to obtain 25 cm³ of potassium hydroxide solution
- **step M** take a final reading from the burette and calculate the volume of the dilute hydrochloric acid reacted
- **step N** run the dilute hydrochloric acid from the burette into the conical flask until the indicator changes colour
- (i) Write the steps in the correct order.

Some of the steps have been completed for you.

(1)

first step				last step
К				М

(ii) Suggest an alternative piece of apparatus that could be used in step L to obtain exactly 25.0 cm³ of potassium hydroxide solution.

(1)

(b)	A student was then asked to produce a pure sample of solid potassium chloride.	
	After finding the volume of acid reacted in step M, the student added this volume of acid to a fresh 25.0 cm ³ sample of the potassium hydroxide solution. This mixture was then evaporated.	
	(i) Explain why this new mixture was evaporated rather than the original mixture from the titration, to produce a pure sample of solid potassium chloride.	(2)
	(ii) After evaporation, the mass of the potassium chloride was determined.	
	The theoretical yield of the experiment was 0.70 g. The actual yield was 0.84 g.	
	This gave a percentage yield greater than 100%.	
	Calculate the percentage yield of this experiment.	(2)
	percentage yield =	
	(iii) Suggest a reason why the actual yield was greater than the theoretical yield.	(1)



(iv) The equation for the reaction between potassium hydroxide solution and dilute hydrochloric acid is

$$KOH + HCl \rightarrow KCl + H_2O$$

Calculate the atom economy for the production of potassium chloride from potassium hydroxide and hydrochloric acid. (relative formula masses: KOH = 56.0, HCl = 36.5, KCl = 74.5, $H_2O = 18.0$)

Give your answer to one decimal place.

(4)

atom economy =%

(Total for Question 4 = 11 marks)

5 (a) A sample of rock salt contains a mixture of sodium chloride and some insoluble substances.

The rock salt is added to water and the mixture stirred.

The mixture is then filtered to obtain a filtrate of sodium chloride solution.

(i) Draw a labelled diagram of the apparatus used to filter the mixture and collect the sodium chloride solution.

(2)

(ii) Describe how a sample of pure, dry sodium chloride crystals can be obtained from the filtrate.

(3)



(b) Inks contain coloured dyes.

Samples of four inks, **W**, **X**, **Y** and **Z**, were separated using paper chromatography. Figure 5 shows the chromatogram obtained.




Figure 5

(i) In the experiment, the solvent front moved 6.5 cm. Calculate the $R_{\rm f}$ value of the dye that is present in both inks $\bf X$ and $\bf Z$.

(1)

$$R_c =$$

(ii) State what could be changed in the experiment to make the $R_{\scriptscriptstyle f}$ value more accurate.

(1)

(iii) In this experiment, ink sample **Y** did not move from the start line. Explain a change to the experiment that would be needed to separate the dyes in ink sample **Y**.

(2)

(Total for Question 5 = 9 marks)



BLANK PAGE

- **6** Titanium and iron are examples of transition metals.
 - (a) Figure 6 shows the percentage abundance of each isotope in a sample of titanium.




Figure 6

Calculate the relative atomic mass of titanium in this sample.

(3)

relative atomic mass =

- (b) Iron, when heated in air, reacts with oxygen to form iron oxide.
 - (i) This reaction is an example of

(1)

(3)

- A crystallisation
- **B** distillation
- C neutralisation
- **D** oxidation
- (ii) The equipment shown in Figure 7 can be used to find the mass of oxygen that combines with iron.




Figure 7

Describe how the equipment shown in Figure 7 could be used to find the mass of oxygen that combines with 0.500 g of iron wool in a crucible and lid of known mass.

(c)	2.24 g of iron combines with 0.96 g of oxygen to form an oxide of iron.							
	Determine the formula of this oxide of iron and use it to complete the balanced equation.							
	(relative atomic masses: Fe = 56.0, O = 16.0)							
	You must show your working.							
		(4)						
	balanced equation for the reaction is							

.....Fe + $O_2 \rightarrow$

(Total for Question 6 = 11 marks)

BLANK PAGE

7	(a)	Methane	reacts with	steam to	form	hydrogen	and carbon	dioxide.
---	-----	---------	-------------	----------	------	----------	------------	----------

The reaction takes place in two stages.

stage 1: $CH_4(g) + H_2O(g) \rightleftharpoons 3H_2(g) + CO(g)$

stage 2: $CO(g) + H_2O(g) \rightleftharpoons H_2(g) + CO_2(g)$

(i) Stage 1 takes in heat energy, it is endothermic.

Explain the effect of increasing the temperature on the yield of the products of stage 1.

(2)

(ii) The overall equation for the process is


$$CH_4(g) + 2H_2O(g) \rightarrow 4H_2(g) + CO_2(g)$$

0.40 g of methane were fully reacted with steam to form carbon dioxide and hydrogen.

Calculate the maximum volume of hydrogen in dm³, measured at room temperature and pressure, that could be made in this reaction.

(relative formula mass: $CH_4 = 16$, 1 mol of any gas at room temperature and pressure occupies 24 dm³)

(3)





*(b) Hydrogen-oxygen fuel cells can be used to provide electrical energy in a spacecraft.								
The reaction that takes place in the fuel cell is								
hydrogen + oxygen → water								
Evaluate the advantages and disadvantages of providing electrical energy in a spacecraft using hydrogen-oxygen fuel cells rather than chemical cells. (6)								

(Total for Question 7 = 11 marks)



- **8** (a) Calcium has an atomic number of 20. A calcium atom has a mass number of 40.
 - (i) Which row of the table shows the number of protons and number of neutrons in this atom of calcium?

(1)

		number of protons	number of neutrons
X	Α	20	20
X	В	40	20
X	C	20	60
X	D	60	20

(ii) Figure 8 shows the arrangement of electrons in an atom of calcium.




Figure 8

Explain, using the information in Figure 8, in which period of the periodic table calcium can be found.

(2)

(b) Ca	lcium and potassium react with water in similar ways.	
(i)	One similarity in the reactions is that hydrogen gas is produced.	
	State one other similarity in the products of the reactions of calcium and potassium with water.	
		(1)
(ii)	Potassium is higher in the reactivity series than calcium and reacts more vigorously with water than calcium reacts with water.	
	State why potassium is higher in the reactivity series and reacts more vigorously with water than calcium.	
		(1)

*(c) Calcium chloride can be prepared by the reaction of calcium with chlorine gas.

Figure 9 shows some properties of calcium, chlorine and calcium chloride.

substance	relative melting	ability to conduct electricity			
substance	point	when solid	when molten		
calcium	high	good	good		
chlorine	low	poor	poor		
calcium chloride	high	poor	good		

Figure 9

(6)

(Total for Question 8 = 11 marks)



- **9** (a) Dilute hydrochloric acid is a strong acid.
 - (i) Explain why dilute hydrochloric acid is described as a strong acid.

(2)

(ii) 1 cm³ of hydrochloric acid of pH 2 is made up to a volume of 10 cm³ with distilled water.

State the pH of the new solution.

(1)

pH =

(b) Magnesium oxide reacts with dilute hydrochloric acid to produce magnesium chloride solution and water.

$$MgO(s) + 2HCl(aq) \rightarrow MgCl_2(aq) + H_2O(l)$$

Write the ionic equation for this reaction.

(3)

(c) In an experiment magnesium hydroxide powder is added in 0.1 g portions to 25 cm³ of dilute hydrochloric acid until the magnesium hydroxide is just in excess.

Universal indicator paper can be used to test the pH of the solution after each addition of magnesium hydroxide.

(i) Give the name of an alternative piece of equipment that can be used to measure pH.

(1)

(ii)	State and explain how the pH changes as the magnesium hydroxide is added to the dilute hydrochloric acid.	(4)
	(Total for Question 9 = 11 ma	rks)

10 (a) Ammonia is manufactured by the Haber process.

The equation for the reaction is

$$N_2(g) + 3H_2(g) \rightleftharpoons 2NH_3(g)$$

The reaction is reversible and can reach equilibrium.

(i) An iron catalyst can be used in the reaction.

Which row of the table shows how adding the iron catalyst affects the rate of attainment of equilibrium and the equilibrium yield of ammonia?

(1)

		rate of attainment of equilibrium	equilibrium yield of ammonia
X	Α	increases	increases
X	В	decreases	does not change
X	C	decreases	increases
X	D	increases	does not change

(ii) Which of the following statements is correct when the reaction reaches equilibrium?

(1)

- A the reverse reaction starts to take place
- **B** the amounts of nitrogen, hydrogen and ammonia are equal
- ☑ C the amounts of nitrogen, hydrogen and ammonia become constant
- **D** the reaction stops

(b) Ammonium sulfate and ammonium nitrate are used as fertilisers as they both contain nitrogen, which will increase the yield of crops. (i) Suggest one other reason for using solid ammonium sulfate and solid ammonium nitrate as nitrogenous fertilisers. (ii) Ammonium nitrate can be made by the reaction of ammonia with nitric acid. Write the balanced equation for this reaction.		Explain what effect a pressure higher than 200 atmospheres would have on the rate of attainment of equilibrium and on the equilibrium yield of ammonia.	e
contain nitrogen, which will increase the yield of crops. (i) Suggest one other reason for using solid ammonium sulfate and solid ammonium nitrate as nitrogenous fertilisers. (ii) Ammonium nitrate can be made by the reaction of ammonia with nitric acid. Write the balanced equation for this reaction. (iii) Describe one similarity and one difference between the industrial production of ammonium sulfate and the laboratory preparation of ammonium sulfate.		rate of attainment of equilibrium and on the equilibrium yield of animonia.	(4)
contain nitrogen, which will increase the yield of crops. (i) Suggest one other reason for using solid ammonium sulfate and solid ammonium nitrate as nitrogenous fertilisers. (ii) Ammonium nitrate can be made by the reaction of ammonia with nitric acid. Write the balanced equation for this reaction. (iii) Describe one similarity and one difference between the industrial production of ammonium sulfate and the laboratory preparation of ammonium sulfate.			
contain nitrogen, which will increase the yield of crops. (i) Suggest one other reason for using solid ammonium sulfate and solid ammonium nitrate as nitrogenous fertilisers. (ii) Ammonium nitrate can be made by the reaction of ammonia with nitric acid. Write the balanced equation for this reaction. (iii) Describe one similarity and one difference between the industrial production of ammonium sulfate and the laboratory preparation of ammonium sulfate.			
contain nitrogen, which will increase the yield of crops. (i) Suggest one other reason for using solid ammonium sulfate and solid ammonium nitrate as nitrogenous fertilisers. (ii) Ammonium nitrate can be made by the reaction of ammonia with nitric acid. Write the balanced equation for this reaction. (iii) Describe one similarity and one difference between the industrial production of ammonium sulfate and the laboratory preparation of ammonium sulfate.			
contain nitrogen, which will increase the yield of crops. (i) Suggest one other reason for using solid ammonium sulfate and solid ammonium nitrate as nitrogenous fertilisers. (ii) Ammonium nitrate can be made by the reaction of ammonia with nitric acid. Write the balanced equation for this reaction. (iii) Describe one similarity and one difference between the industrial production of ammonium sulfate and the laboratory preparation of ammonium sulfate.			
(ii) Ammonium nitrate can be made by the reaction of ammonia with nitric acid. Write the balanced equation for this reaction. (iii) Describe one similarity and one difference between the industrial production of ammonium sulfate and the laboratory preparation of ammonium sulfate.		· · · · · · · · · · · · · · · · · · ·	
Write the balanced equation for this reaction. (iii) Describe one similarity and one difference between the industrial production of ammonium sulfate and the laboratory preparation of ammonium sulfate.	(i)		(1)
(iii) Describe one similarity and one difference between the industrial production of ammonium sulfate and the laboratory preparation of ammonium sulfate.	(ii) Ammonium nitrate can be made by the reaction of ammonia with nitric acid.	
of ammonium sulfate and the laboratory preparation of ammonium sulfate.		Write the balanced equation for this reaction.	(2)
•	(ii		(2)
ference	nilarity		
	fference	e	
(Total for Question 10 = 11 mark		(Total for Question 10 = 11 ma	rks)



The Periodic Table of the Elements

0 4 He helium 2	20 Ne	40 Ar argon 18	84 Kr krypton 36	131 Xe xenon 54	[222] Rn radon 86	t full y
7	19 F fluorine 9	35.5 CI chlorine 17	80 Br bromine 35	127 	[210] At astatine 85	orted but not
9	16 O oxygen 8	32 S sulfur 16	79 Se selenium 34	128 Te tellurium 52	[209] Po polonium 84	ave been rep
2	14 N nitrogen 7	31 P phosphorus 15	75 As arsenic 33	122 Sb antimony 51	209 Bi bismuth 83	s 112–116 ha authenticated
4	12 C carbon 6	28 Si silicon 14	73 Ge germanium 32	119 Sn tin 50	207 Pb lead 82	mic numbers
ဇ	11 B boron 5	27 Al aluminium 13	70 Ga gallium 31	115 In indium 49	204 TI thallium 81	Elements with atomic numbers 112–116 have been reported but not fully authenticated
			65 Zn zinc 30	112 Cd cadmium 48	201 Hg mercury 80	Elem
			63.5 Cu copper 29	108 Ag silver 47	197 Au gold 79	Rg roentgenium 111
			59 nickel 28	106 Pd palladium 46	195 Pt platinum 78	[271] Ds damstadtium 110
			59 Co cobalt 27	103 Rh rhodium 45	192 	[268] Mt meitnerium 109
T hydrogen			56 iron 26	101 Ru ruthenium 44	190 Os osmium 76	[277] Hs hassium 108
			55 Mn manganese 25	[98] Tc technetium 43	186 Re rhenium 75	[264] Bh bohnium 107
	mass ɔol ıumber		52 Cr	96 Mo molybdenum 42	184 W tungsten 74	[266] Sg seaborgium 106
Key	relative atomic mass atomic symbol name atomic (proton) number		51 V vanadium 23	93 Nb niobium 41	181 Ta tantalum 73	[262] Db dubnium 105
	relativ ato atomic		48 Ti tftanium 22	91 Zr zirconium 40	178 Hf hafnium 72	[261] Rf nutherfordium 104
			45 Sc scandium 21	89 Y yttrium 39	139 La* Ianthanum 57	[227] Ac* actinium 89
2	9 Be beryllium 4	24 Mg magnesium	40 Ca calcium 20	88 Sr strontium 38	137 Ba barium 56	[226] Ra radium 88
~	7 Li lifhium 3	23 Na sodium 11	39 K potassium 19	85 Rb rubidium 37	133 Cs caesium 55	[223] Fr francium 87

^{*} The lanthanoids (atomic numbers 58–71) and the actinoids (atomic numbers 90–103) have been omitted.

The relative atomic masses of copper and chlorine have not been rounded to the nearest whole number.