
Vasculitis and related disorders

Soheil S Dadras MD-PhD


October 16, 2023

Tissue reaction patterns




Atopic dermatitis

Spongiotic



Psoriasis


Psoriasiform



Lichen planus


Lichenoid

Tissue reaction patterns




Bullous pemphigoid

Vesiculobullous



Granuloma annulare


Granulomatous



Atrophie blanche

Vasculopathic


Cutaneous vascular system



Immunohistochemical markers for blood vessels and lymphatics

Marker	Lymphatic Vessels	Blood Vessels
Blood Vascular Specific		
CD34	-	+
CD44	-	+
PAL-E	-	+
Collagen type IV	-/(+)	+
Collagen type XVIII	-/(+)	+
Laminin	-/(+)	++
Neuropilin-1	-	+
Lymphatic Specific		
VEGFR-3	+	-
Podoplanin	+	-
SLC/CCL21	+	-
LYVE-1	+	-
Prox1	+	-
Panvascular		
CD31 (PECAM-1)	+	++
VEGFR-2	+	+
Factor VIII-related antigen	+	++

Dadras and Detmar, Hem. Onc. of North Ame. 2004




Diagnostic approach to vasculitis

- Is there vasculitis or vasculopathy?
- Is the process cutaneous or systemic?
- Is the vascular injury primary or secondary?
- What is the size and the type of vessel involved?
 - Small, medium or large?
- What is the composition of the inflammatory infiltrate?
 - Neutrophilic, lymphocytic, eosinophilic, or granulomatous

- What are the associated findings?
 - Infection (special stains and microbial cultures)?
 - Plasma cells (spirochete)?
 - Viral infection (HSV vascular changes)?
 - Suggestion of connective tissue disease (interface dermatitis or dermal mucin)
 - Edema (urticarial reaction)
 - Serologic studies: ANA, ANCAs, RF, cryoglobulins, DIF
 - Clinical context: demographics and presentation, other signs/symptoms

Diagnostic approach to make/confirm vasculitis or vascular damage

- The damage could involve blood vessels in the deep dermis or subcutis
- Histologic evaluation: Perform punch (NOT shave) skin biopsy to include deep reticular dermis and subcutaneous tissue
- Direct immunofluorescence
 - Vascular pattern of immunoglobulins and complements (C3)
- PAS and elastic special stains
- IHC: smooth muscle actin





Leukocytoclastic vasculitis: IgM in the luminal surface of blood vessels

Categorization of disease of blood vessels

Types and definitions of vasculitis

- Adopted by the 2012 Chapel Hill Consensus Conference
- List of clinical criteria and disease nomenclature
- Classified according to the type and size of the vessels
 - e.g., polyarteritis nodosa is under medium vessel vasculitis





Fibrinoid necrosis and Inflammation


Clinical and histologic definitions


Clinical

- Vascular damage:
 - Edema
 - Livedo reticularis (net-like pattern of red-blue skin discoloration)
 - Subcutaneous hemorrhage, or purpura
 - Petechiae, < 3-mm diameter
 - Ecchymoses, larger
 - Palpable purpura (raised, inflammatory infiltrate) [Punch Biopsy]
 - Severe
 - Vascular occlusion --> ischemia --> necrosis --> gangrene --> ulceration


Histologic


- Necrosis of vessel wall with deposition of fibrinogen material
- Leukocytoclasis (nuclear debris from infiltrating neutrophils)
- Karyorrhexis (nuclear dust)
- Extravasation of erythrocytes
- Endothelial cell swelling
- Luminal thrombosis
- Edema





Lymphocytic vasculitis: vascular damage + lymphocytes ± minimal fibrin






Numerous lymphocytes surround a blood vessel; no leukocytoclasis


Antineutrophil cytoplasmic antibodies (ANCA)-associated systemic vasculitis


- Serologic marker to detect specific autoantibody
- Indirect immunofluorescence (IIF) of normal peripheral blood neutrophils
- Followed by enzyme-linked immunosorbent assay (ELISA)
- IIF two patterns:
 - Cytoplasmic (c-ANCA); proteinase 3
 - Perinuclear (p-ANCA); myeloperoxidase



c-ANCA


Medium and large-seized vasculitis





Elastic van Gieson: disruption of internal elastic lamina

Neutrophilic dermatoses: Neutrophilic vascular reaction (not necrotizing vasculitis)



Vascular occlusive disease: vasculopathic reaction and pseudovasculitis

Cryoglobulinemia


Cholesterol embolism

Livedo reticularis

Atrophie blanche


Degos syndrome

Calciphylaxis




Atrophie blanche (livedoid vasculopathy) white atrophy


- Thrombogenic vasculopathy
- Decreased fibrinolytic activity (tissue plasminogen activator release defective)
- "Thrombogenic state" in several conditions
 - Lupus-type anticoagulant
 - Increased anticardiolipin antibody
- Painful, purpuric ulcers with reticular pattern on the lower extremities
- Histopathology:
 - Fibrinoid material
 - Minimal lymphocytic inflammation
 - Necrotic blood vessels
 - No Leukocytoclasis
 - Extravasation of erythrocytes
 - No edema



Unknown cases

Will be available in full on https://digitalskinpathology.com/







Smooth muscle actin


Polyarteritis nodosa

- 64/F LEFT ARM PUNCH with h/o NLD
- The pathology report pertaining to the prior biopsy from right leg/subcutaneous nodule is noted and reviewed (SGS-21-12351).
- The prior biopsy was sent to Dr. at UCSF Dermatopathology Laboratory for second opinion. His diagnosis was palisaded granulomatous dermatitis with focal vasculitis, probably necrobiosis lipoidica.

- LEUKOCYTOCLASTIC VASCULITIS INVOLVING MEDIUM-SIZED MUSCULAR ARTERY IN DEEP DERMIS.
- The current biopsy demonstrates a different histopathology than the prior in that there is no panniculitis. There is only pure leukocytoclastic vasculitis involving medium-sized artery, which can be seen in polyarteritis nodosa. Although necrobiosis lipoidica can demonstrate secondary lymphocytic vasculitis, the current findings are diagnostic for leukocytoclastic vasculitis and do not support necrobiosis lipoidica.











Urticarial Vasculitis

- 34/F RIGHT THIGH PUNCH BIOPSY
- L50.1; DIFFUSE URTICARIAL PAPULES, AREAS OF BRUISING, R/O URTICARIAL VASCULITIS
- COMPATIBLE WITH URTICARIAL VASCULITIS. Overall, the findings support the clinical impression of urticarial vasculitis, early phase. The differential diagnosis includes acute urticaria. **Diagnostic features of** established leukocytoclastic vasculitis are absent.









LCV in SLE: Lupus vasculitis

- L30.9; 41/F RIGHT THIGH H&E and DIF. PATIENT WITH A HISTORY OF SLE COMPLICATED BY NEPHROTIC SYNDROME AND RHEUMATOID ARTHRITIS WITH AN ACUTE ONSET RASH THAT INCLUDED ARCUATE PLAQUES, EDEMATOUS PAPULES AND PLAQUES ON THE CHEST AND BACK WITH NO MALAR RASH
- LEUKOCYTOCLASTIC VASCULITIS, INVOLVING SMALL-CALIBER BLOOD **VESSELS IN SUPERFICIAL-MID DERMIS.** The findings of leukocytoclastic vasculitis in this clinical context may represent vasculitis in systemic lupus erythematosus, which can be seen in roughly 11-30% of patients with systemic lupus erythematosus, supported by positive lupus band test and vascular reaction according to the results for DIF.


Digital Skin Pathology https://digitalskinpathology.com/

- Meet the challenges of the growing needs for dermatopathology knowledge
- Learn Dermatopathology based on actual reallife cases
- Dermatology PAs and NPs
- Primary MDs and general surgeons
- Residents of Dermatology and Pathology



Learn Histologic Diagnosis Case-By-Case

DERMATOPATHOLOGY: LEARN HOW TO DIAGNOSE SKIN DISEASES DERM PATH DIAGNOSTICS



Understand your patient's dermatopathology diagnostic report to provide better clinical care (how to diagnose skin diseases). derm path diagnostics

References

- Dermatopathology
 - Raymond L. Barnhill, editor-in-chief
 - A. Neil Crowson
 - Cynthia M Magro
 - Michael W Piepkorn


THIRD EDITION

Raymond L. Barnhill Editor-in-Chief


A. Neil Crowson Cynthia M. Magro Michael W. Piepkorn

Dermatopathology














Weedon's SKIN PATHOLOGY



References

- Weedon's skin pathology
- McKee's pathology of the skin