




LYMPH NODES AND IMMUNE SURVEILLANCE

WHERE LYMPHATIC FLOW MEETS THE IMMUNE SYSTEM

By: Laine Kristin Miller
 PTA, LMT, CLT

When we think about the lymphatic system, we often think about fluid: swelling, drainage, lymphatic vessels, and the movement of lymph back toward the bloodstream.

But lymph is more than fluid.

As lymph travels through the tissues, it carries information about what is happening in those tissues. Proteins, cellular debris, antigens, inflammatory mediators, immune cells, and material associated with microorganisms can all enter lymphatic vessels and travel toward regional lymph nodes.

This makes the lymphatic system not only a transportation system, but also an important communication network between peripheral tissues and the immune system.

What Is Immune Surveillance?

Immune surveillance is the continuous process through which the immune system monitors the body for signs that something is different, damaged, infected, or potentially dangerous.

Every moment, immune cells are moving through tissues, blood, lymphatic vessels, and lymphoid organs.

They are looking for information.
Is this cell normal?

Does this protein belong here?
Has a microorganism entered the tissue?
Has a cell been damaged?
Is there evidence of inflammation?
Has a cell become abnormal?

Most of the time, this surveillance happens without us ever knowing it.

Lymph nodes help make that enormous surveillance task possible by bringing together information arriving from the tissues with immune cells capable of recognizing and responding to it.

Lymph Nodes Are More Than Filters

Lymph nodes are frequently described as filters.

That description is correct, but incomplete. A lymph node does not simply strain unwanted material from lymph. It is a highly organized immune organ containing specialized anatomical regions where different immune cells gather, communicate, recognize antigens, and coordinate immune responses.

A lymph node can therefore be thought of as both a filtering station and an immune communication center.

Lymph arriving from a particular region of the body enters its draining lymph nodes through afferent lymphatic vessels. It first reaches the subcapsular sinus and then moves through a network of lymphatic sinuses before leaving through an efferent lymphatic vessel. During that journey, the contents of lymph are exposed to strategically positioned immune cells.

The lymph node is essentially sampling what is happening upstream.

The Journey Begins in the Tissues

Imagine a small area of tissue somewhere in the body.

Cells are constantly exchanging substances with the surrounding interstitial fluid. Proteins, metabolic products, signaling molecules and cellular material may be present within this extracellular environment.


If tissue becomes injured or infected, additional information appears.

There may be:

- fragments of damaged cells
- foreign antigens
- bacteria or viral material
- inflammatory cytokines and chemokines
- proteins
- activated immune cells
- antigen-presenting cells

Some of this material enters the initial lymphatic vessels.

From there, lymph moves into larger collecting vessels and toward the regional lymph nodes. In this way, lymphatic vessels continually carry a molecular and cellular sample of the tissues they drain.



Entering the Lymph Node

Afferent lymphatic vessels deliver lymph into a space just beneath the capsule of the lymph node called the subcapsular sinus.

This is an important first checkpoint.

Macrophages positioned along lymphatic sinuses are directly exposed to incoming lymph and can rapidly capture microorganisms and other lymph-borne material.

Smaller soluble molecules can also gain access to specialized conduit networks that rapidly deliver information deeper into the node. Larger particles and cells may be handled differently and remain accessible to immune cells positioned around the sinuses.

This means the lymph node does not randomly mix everything together.

Its architecture determines where information goes and which immune cells encounter it.

The Neighborhoods Inside a Lymph Node


A lymph node contains specialized regions populated by different immune cells.

The outer cortex contains B-cell follicles. Deeper within the node, the paracortex is particularly rich in T lymphocytes and dendritic cells.

The medulla contains lymphatic sinuses along with macrophages, plasma cells and other immune cells.

This organization is extremely important.

Instead of relying on the chance meeting of an antigen and one of the relatively small number of lymphocytes capable of recognizing it, the lymph node brings enormous numbers of immune cells through an organized environment designed to increase the probability that the correct cells will meet.



Dendritic Cells: Carrying the Message

Dendritic cells are among the most important antigen-presenting cells involved in initiating adaptive immune responses.

Some dendritic cells reside in peripheral tissues, where they continually sample their environment.

When they encounter potentially important material, particularly in the setting of inflammatory or danger signals, dendritic cells can process antigens and migrate into lymphatic vessels.

They then travel through afferent lymphatics toward the draining lymph node. Once inside the lymph node, chemokine signals help direct these cells toward T-cell-rich regions.

The dendritic cell is now carrying information from the tissue directly to immune cells capable of mounting a highly specific response.

Antigen Presentation: Showing the Immune System What Was Found

T cells do not simply react to everything floating through lymph.

They require a much more controlled process. Antigen-presenting cells such as dendritic cells process proteins into smaller fragments and display those fragments on molecules called the major histocompatibility complex, or MHC. This is antigen presentation.

A useful way to picture it is:

Tissue → antigen → dendritic cell → lymphatic vessel → lymph node → T cell
Inside the lymph node, T cells move through the T-cell zone interacting with antigen-presenting cells.


Most encounters do not produce an immune response.

But if a T cell possesses a receptor capable of recognizing the presented antigen—and the appropriate additional signals are present—that T cell can become activated. This is one of the remarkable features of lymph nodes.

A rare T cell capable of recognizing one particular antigen can encounter a dendritic cell carrying exactly that antigen because both have been brought into the same highly organized environment.

Continuous lymphocyte trafficking through lymph nodes is therefore fundamental to immune surveillance.

T Cells: Coordinating and Attacking



Different populations of T cells perform different functions.

CD4⁺ helper T cells help coordinate immune responses through cellular communication and cytokine signaling. They can assist B cells in producing effective antibody responses and help regulate the activity of other immune cells.

CD8⁺ cytotoxic T cells can develop the ability to recognize and destroy infected or abnormal cells displaying their specific antigen. Regulatory T cells help control immune activity and reduce inappropriate or excessive immune responses.

Once an appropriate T cell has been activated, it undergoes clonal expansion.

One antigen-specific cell can become many.

Those cells can then leave the lymph node, enter circulation, and participate in immune activity elsewhere in the body.

B Cells and Antibody Production

B cells occupy primarily follicular regions within the lymph node.

Unlike T cells, B cells can recognize intact antigen through their B-cell receptors. Following antigen recognition and appropriate signals—often including assistance from helper T cells—activated B cells can proliferate and develop into specialized populations.

Some become plasma cells, which manufacture antibodies.

Others become memory B cells, helping the immune system respond more efficiently if the same antigen is encountered again.

During an active response, structures called germinal centers can develop within lymph node follicles. Here, B cells undergo processes that help refine the antibody response and generate cells capable of producing antibodies with improved affinity for their target.

This is one reason lymph nodes can enlarge during an immune response.

The node is not simply filling with debris. Immune cells are arriving, communicating, proliferating, differentiating and organizing a response.

Antigen Presentation: Showing the Immune System What Was Found

T cells do not simply react to everything floating through lymph.


They require a much more controlled process. Antigen-presenting cells such as dendritic cells process proteins into smaller fragments and display those fragments on molecules called the major histocompatibility complex, or MHC. This is antigen presentation.

A useful way to picture it is:

Tissue → antigen → dendritic cell → lymphatic vessel → lymph node → T cell
Inside the lymph node, T cells move through the T-cell zone interacting with antigen-presenting cells.

Most encounters do not produce an immune response.

But if a T cell possesses a receptor capable of recognizing the presented antigen—and the appropriate additional signals are present—that T cell can become activated. This is one of the remarkable features of lymph nodes.



Macrophages: The First Line Inside the Node

Macrophages are positioned strategically along lymphatic sinuses.

Their location allows them to encounter lymph-borne material almost immediately after it enters the node.

They can capture and engulf microorganisms, particles and cellular debris through phagocytosis.

Subcapsular sinus macrophages are particularly important because they sit near the first region exposed to incoming lymph. Other macrophage populations are located deeper within the node, including the medullary sinuses.

These macrophages participate not only in cleanup but also in immune communication. The node is therefore performing several functions simultaneously:

sampling → capturing → processing →
presenting → communicating → responding

How Do Lymphocytes Get Into the Node?

Interestingly, lymph is not the only way immune cells enter a lymph node.


Large numbers of naïve lymphocytes arrive from the bloodstream.

Specialized blood vessels within lymph nodes called high endothelial venules, or HEVs, allow circulating lymphocytes to leave the blood and enter the lymph node.

This creates an extraordinary meeting point. From one direction comes information from the tissues through lymphatic vessels.

From another direction come circulating lymphocytes through the blood.

The lymph node brings them together. That is one of the fundamental principles of immune surveillance.



If Nothing Is Wrong

Immune surveillance does not mean that the immune system should attack everything it encounters.

Quite the opposite.

Most material encountered by the immune system does not need to trigger a destructive response.

Lymphocytes continually pass through lymph nodes looking for their specific antigens.

If they do not encounter the signals required for activation, they leave and continue circulating

When a Threat Is Identified

When the appropriate combination of antigen recognition and immune signaling occurs, the quiet surveillance environment changes.

Dendritic cells activate antigen-specific T cells. B cells recognizing relevant antigens can become activated.

T and B cells proliferate.

Plasma cells begin producing antibodies. Cytokines and chemokines coordinate communication.

Activated lymphocytes begin leaving the lymph node to participate in the immune response. The lymph node may become enlarged and tender as cellular activity increases.

What we feel as a "swollen lymph node" can therefore represent an extraordinarily complex biological response occurring at the microscopic level.

Lymph Nodes Create Immunological Memory

One of the most powerful features of adaptive immunity is memory.

After an immune response has resolved, some activated B and T lymphocytes remain as memory cells.

If the same antigen appears again, these cells can respond more rapidly and effectively than during the first encounter.

This principle is also fundamental to vaccination.

A vaccine introduces antigenic information in a controlled manner so that the adaptive immune system can develop antigen-specific responses and memory without requiring the disease itself to provide that first exposure.

Lymph nodes are major sites where those adaptive immune responses are organized.

Lymph Nodes and Abnormal Cells

Immune surveillance is not limited to microorganisms.

Cells within our bodies continually change, become damaged and die. Immune cells can recognize certain abnormal cellular signals and antigens associated with altered cells.

This is particularly relevant to cancer biology. Lymphatic vessels can transport tumor-associated antigens and antigen-presenting cells toward draining lymph nodes, potentially contributing to antitumor immune responses.

At the same time, cancer cells themselves can sometimes enter lymphatic vessels and travel to regional lymph nodes.

This helps explain why lymph nodes have such an important role in cancer evaluation and staging.

The relationship is complex: lymph nodes can participate in antitumor immunity while also serving as sites to which some cancers spread.

Why Lymphatic Flow Matters

The relationship between lymphatic flow and immunity is deeper than simply "moving waste." Lymphatic vessels transport immune cells, antigens, proteins and molecular signals from peripheral tissues toward lymph nodes.

The lymph node then provides the organized environment where that information can be examined.


The immune response depends upon movement:

from tissue → into lymph → through lymphatic vessels → into lymph nodes → between immune cells → out of the node → back toward circulation and tissues.

The lymphatic and immune systems are therefore inseparable.

A Communication Network Throughout the Body

The lymphatic system connects tissues that may be centimeters—or much farther—away from the immune cells that ultimately respond to what is happening there.



A small inflammatory event in the skin can send molecular information toward a regional lymph node.

An antigen encountered in peripheral tissue can be carried by a dendritic cell through a lymphatic vessel.

A rare antigen-specific T cell traveling through the bloodstream can enter that same node through an HEV.

Inside the lymph node, those two cells can meet.

- Recognition occurs.
- Signals are exchanged.
- Cells proliferate.

A response leaves the node and travels back into the body.

All of this can begin with microscopic events occurring within the interstitial space.

Nothing Works in Isolation

The lymphatic system is sometimes reduced to a drainage system, but drainage is only one part of its physiology.

Lymphatic vessels connect the interstitial environment to lymph nodes. Lymph nodes connect tissue information with the immune system. Immune cells move between lymph, blood, organs and tissues.

The cardiovascular system supplies circulating immune cells. The extracellular matrix influences how cells and molecules move through tissues. The nervous, endocrine and inflammatory systems contribute additional layers of communication.

Nothing works in isolation.

The lymphatic system quite literally communicates with every tissue and organ it serves.

And the lymph node demonstrates this beautifully.

It is not simply a structure that lymph passes through.

It is a place where flow becomes information, information becomes recognition, and recognition can become an immune response.

Understanding lymph nodes therefore changes the way we look at the lymphatic system itself. It is not simply moving fluid.

It is helping the body continuously ask one essential question:

What is happening in our tissues right now?

About the Author



Laine Miller, PTA, LMT, CLT, C-MLD

With more than 35 years of experience in physical therapy and bodywork, Laine is passionate about helping people understand the intricate connections within the body—especially the vital role of the lymphatic system in health, healing, and longevity.

Laine is a Certified Lymphedema Therapist, Certified Manual Lymph Drainage Therapist, Licensed Massage Therapist, and Physical Therapist Assistant.

She is the founder and developer of Lymphatic Restoration Therapy® (LRT) and the Systemic Flow™ series, and the author of *The Lymph Connection*.

References

Girard JP, Moussion C, Förster R. HEVs, lymphatics and homeostatic immune cell trafficking in lymph nodes. *Nature Reviews Immunology*. 2012;12:762–773. doi:10.1038/nri3298. [Nature](#)

von Andrian UH, Mempel TR. Homing and cellular traffic in lymph nodes. *Nature Reviews Immunology*. 2003;3:867–878. doi:10.1038/nri1222. [Nature](#)

Grant SM, Lou M, Yao L, Germain RN, Radtke AJ. The lymph node at a glance—how spatial organization optimizes the immune response. *Journal of Cell Science*. 2020;133:jcs241828. [PubMed Central \(PMC\)](#)

Förster R, Braun A, Worbs T. Lymph node homing of T cells and dendritic cells via afferent lymphatics. *Trends in Immunology*. 2012;33(6):271–280.

Russo E, Nitschké M, Halin C. Dendritic cell interactions with lymphatic endothelium. *Lymphatic Research and Biology*. 2013;11(3):172–182. [PubMed Central \(PMC\)](#)
Randolph GJ, Angeli V, Swartz MA. Dendritic-cell trafficking to lymph nodes through lymphatic vessels. *Nature Reviews Immunology*. 2005;5:617–628.

Alvarez D, Vollmann EH, von Andrian UH. Mechanisms and consequences of dendritic cell migration. *Immunity*. 2008;29(3):325–342. [PubMed Central \(PMC\)](#)

Förster R, Davalos-Misslitz AC, Rot A. CCR7 and its ligands: balancing immunity and tolerance. *Nature Reviews Immunology*. 2008;8:362–371.

Miyasaka M, Tanaka T. Lymphocyte trafficking across high endothelial venules: dogmas and enigmas. *Nature Reviews Immunology*. 2004;4:360–370. [Nature](#)

Gray EE, Cyster JG. Lymph node macrophages. *Journal of Innate Immunity*. 2012;4(5–6):424–436. [PubMed Central \(PMC\)](#)

Lämmermann T, Sixt M. The microanatomy of T-cell responses. *Immunological Reviews*. 2008;221:26–43.
Fletcher AL, Acton SE, Knoblich K. Lymph node fibroblastic reticular cells in health and disease. *Nature*

Reviews Immunology. 2015;15:350–361.
Krishnamurty AT, Turley SJ. Lymph node stromal cells: cartographers of the immune system. *Nature Immunology*. 2020;21:369–380.

Hirosue S, Dubrot J. Modes of antigen presentation by lymph node stromal cells and their immunological implications. *Frontiers in Immunology*. 2015;6:446. [PubMed Central \(PMC\)](#)

Hampton HR, Chtanova T. Lymphatic migration of immune cells. *Frontiers in Immunology*. 2019;10:1168. [PubMed Central \(PMC\)](#)

Qi H, Kastenmüller W, Germain RN. Spatiotemporal basis of innate and adaptive immunity in secondary lymphoid tissue. *Annual Review of Cell and Developmental Biology*. 2014;30:141–167.

Gerner MY, Torabi-Parizi P, Germain RN. Strategically localized dendritic cells promote rapid T cell responses to lymph-borne particulate antigens. *Immunity*. 2015;42(1):172–185.

Junt T, Scandella E, Ludewig B. Form follows function: lymphoid tissue microarchitecture in antimicrobial immune defence. *Nature Reviews Immunology*. 2008;8:764–775.

Bajénoff M, Egen JG, Koo LY, et al. Stromal cell networks regulate lymphocyte entry, migration, and territoriality in lymph nodes. *Immunity*. 2006;25(6):989–1001.

Lancaster JN. Aging of lymphoid stromal architecture impacts immune responses. *Seminars in Immunology*. 2023;70:101817. [PubMed Central \(PMC\)](#)