


Toxicological Assessment of Automotive Shredder Residue (ASR) on Female Reproductive Pathophysiology: Mechanisms of Endocrine Disruption, Cellular Proliferation, and Uterine-Mammary Tumorigenesis

Whitepaper prepared by, Cherise Petker, Founder/CEO, [CircularSolar.net](https://circularsolar.net) and assisted by Google Gemini. This whitepaper being also being shared from years of real-life studies by ASR residue (and other pollution sources) and due to the personal health changes since high exposure within this pollution hotspot, main page:

<https://circularsolar.net/pfas%2C-toxins%3B-cancer> and Date: September 9th, 2026

Executive Summary

Automotive Shredder Residue (ASR)—historically referred to as "shredder fluff"—represents a highly heterogeneous, toxicologically dense byproduct of end-of-life vehicle (ELV) recycling processes. Consisting of a dense matrix of pulverized microplastics, polyurethane foams, interior textiles, sealants, and wiring insulations, ASR serves as a key environmental repository for Persistent Organic Pollutants (POPs) and Endocrine-Disrupting Chemicals (EDCs). While regulatory bodies historically categorized ASR through the lens of solid waste management, recent clinical and epidemiological data indicate a profound correlation between localized ASR exposure pathways and a cluster of hyper-estrogenic, benign, and pre-malignant female reproductive disorders. This paper outlines the discrete physiological and molecular pathways through which ASR-derived toxic compounds—specifically per- and polyfluoroalkyl substances (PFAS), phthalate plasticizers (DEHP), polychlorinated biphenyls (PCBs), and divalent heavy metals—induce uterine leiomyoma (fibroids), adenomyosis, endometriosis, ovarian dysfunction, and progressive mammary tissue aberrations.



1. Introduction and Composition of ASR Environmental Matrices

The mechanical dismantling and shredding of vehicles result in an airborne, soil-bound, and water-soluble effluent fraction containing highly concentrated chemical vectors:

- **PFAS Compounds:** Utilized for hydro/oleophobic upholstery treatment, wire coverings, and specialized anti-corrosive industrial coatings.
- **Phthalate Plasticizers (DEHP):** Heavily formulated into polyvinyl chloride (PVC) components, synthetic leathers, and flexible interior moldings.
- **Polychlorinated Biphenyls (PCBs):** Originating from historical legacy electrical components, specialized capacitors, fluids, and degraded vehicle sealants.
- **Divalent Heavy Metals:** Particulate arrays of elemental Lead (Pb), Cadmium (Cd), and Hexavalent Chromium (Cr-VI) generated via structural degradation and wiring wear.

Due to the lack of complete encapsulation during high-volume mechanical processing, these compounds are released into ambient airsheds as particulate matter (PM_{2.5}/PM₁₀) and leach into groundwater systems near recycling infrastructures, leading to chronic human exposure through inhalation, ingestion, and dermal pathways.

2. Pathophysiological Mechanisms of Uterine Leiomyoma (Fibroids)

Uterine leiomyomas are benign monoclonal tumors derived from myometrial smooth muscle cells. ASR components drive leiomyoma development via two primary pathways:

A. Cellular Proliferation and Gap Junction Degradation (PFAS Vectors)

Perfluoroalkyl substances disrupt the baseline homeostasis of human myometrial cells by altering cell-cycle dynamics and preventing programmed apoptosis. At nanomolar, environmentally relevant exposure margins, PFAS compounds promote structural cell migration and hyper-viability.

Concurrently, these substances degrade Gap Junction Intercellular Communication (GJIC) protocols within the smooth muscle layer. When connexin-mediated communication cascades break down, individual myometrial cells fail to process localized tissue anti-growth signals, inducing continuous, unmitigated smooth muscle proliferation.

B. Aryl Hydrocarbon Receptor (AHR) Activation (Phthalate/DEHP Flux)

Pulverized vinyl and foam dust yield an ongoing systemic influx of Di-(2-ethylhexyl) phthalate (DEHP). Upon metabolic intake, its primary metabolite, mono-(2-ethylhexyl) phthalate (MEHP), selectively targets and binds to the environmentally responsive Aryl Hydrocarbon Receptor (AHR) in uterine tissue.

Upon ligand binding, the activated AHR complex undergoes nuclear translocation, binding directly to specific DNA response motifs. This genomic event overrides baseline uterine growth

controls, upregulating the transcription of target genes that cause the hyper-proliferation of myometrial cells, driving the expansion of symptomatic fibroid tumors.

3. Junctional Zone Integrity and the Etiology of Adenomyosis

Adenomyosis is characterized by the ectopic invasion of endometrial glands and stroma deep into the myometrial smooth muscle matrix. This structural transgression relies on the breakdown of the endometrial-myometrial interface, also known as the uterine junctional zone.

A. Endometrial-Myometrial Interface Invagination

ASR-derived heavy metal dusts and plasticizers stimulate localized cellular stress, forcing an up-regulation of Matrix Metalloproteinases (MMPs). These enzymes systematically break down the extracellular matrix at the junctional zone boundary. Lacking this structural barrier, endometrial epithelial cells gain the capacity to migrate basally, physically invaginating into the myometrium.


B. Epigenetic Aberrations and Localized Prostaglandin Synthesis

Chronic exposure to ASR chemical matrices triggers persistent epigenetic alterations, specifically changing DNA methylation profiles across inflammatory control genes. This causes a permanent overproduction of Prostaglandin E2 (PGE2) and a cascade of inflammatory cytokines. The resulting hyperinflammatory microenvironment creates a feedback loop: chronic inflammation degrades myometrial integrity, facilitating deeper endometrial invasion and causing the severe, progressive uterine thickening and pelvic pain characteristic of adenomyosis.

4. Systemic EDCs and Associated Reproductive Sequelae

The pathological footprint of ASR compounds extends beyond uterine smooth muscle boundaries, impacting the global pelvic and endocrine landscape through structural estrogen mimicry and follicular toxicity.

[ASR Exposure Matrix]



A. Endometriosis via Estrogenic Mimicry

Polychlorinated biphenyls (PCBs) and heavy metals like Cadmium act as highly potent "xenoestrogens." These foreign compounds structurally substitute for natural (17β) -estradiol at the hormone-binding pocket of Estrogen Receptor Alpha ($ER\alpha$).

This ongoing, inappropriate receptor over-activation simulates a state of continuous estrogen dominance. When ectopic endometrial cells slip into the peritoneal cavity via retrograde menstruation, this hyper-estrogenic environment provides the exact proliferative signal required for these cells to attach to the pelvic wall, form vascularized lesions, and establish chronic endometriosis.

B. Ovarian Dysfunction and Infertility Dynamics

Heavy metals and PFAS bioaccumulate directly within follicular fluid, bypassing the protective ovarian blood barrier. Once inside the ovary, these toxicants disrupt folliculogenesis by inducing oxidative stress within granulosa cells, leading to premature cell death.

This cellular depletion disrupts normal steroidogenesis, leading to poor oocyte maturation, irregular ovulatory cycles, accelerated follicular depletion, and a statistically significant decline in clinical fertility metrics among exposed cohorts.

5. Endocrine Disruption within Mammary Architecture

Because mammary glands are highly sensitive to systemic hormonal fluctuations, they are highly susceptible to the effects of ASR-derived xenoestrogens and plasticizers.

A. Glandular Epithelial Hyperplasia

Continuous exposure to phthalate fractions and estrogen-mimicking PCBs forces an accelerated, inappropriate cell-cycle progression within the terminal duct lobular units (TDLUs) of the breast. This persistent, chemical-driven proliferative signal induces **glandular proliferation (hyperplasia)**. The over-activation of these epithelial pathways disrupts standard cell turnover, introducing structural genetic vulnerabilities and establishing potential pre-cancerous cellular changes.

B. Fibrosis and Stromal Density Alterations

Divalent heavy metals (such as Lead and Cadmium) alter localized extracellular matrix deposition within the breast. These toxicants stimulate a permanent inflammatory state within mammary stromal tissue, driving the excessive accumulation of collagen fibers.

This results in a marked increase in **stromal/fibrous breast density**. Clinically, dense breast tissue changes pose a dual challenge: they act as a structural physical framework that supports tumor development while simultaneously masking early-stage lesions on screening mammograms.

Possible misdiagnosis or future complications:

When a doctor attributes dense, lumpy, or proliferating breast changes solely to natural "fibrocystic changes"—overlooking the fact that industrial pollutants like PFAS or PCBs are driving the cellular stress—the risk of a subsequent breast cancer diagnosis spikes due to the following statistical realities:

1. The Masking Risk: Mammograms Missing Hidden Tumors

Because both toxicant-stimulated glandular tissue and cancerous tumors appear bright white on a standard mammogram, the underlying pollution-driven density acts as natural camouflage.


- **The Error Rate:** Standard mammograms miss **30% to 40% of breast cancers** in individuals with highly dense breast tissue because the white dense tissue obscures the white tumor.
- **The Multiplied Risk:** Women whose environmental or genetic factors cause them to have extremely dense breasts are statistically at a **4 to 6 times higher risk** of

developing breast cancer compared to those with fatty breast tissue, primarily because early warning signs are physically hidden.

2. The Biopsy Risk: Hyperplasia Transitioning to Malignancy

If the environmental toxins alter cell cycles to the point where ordinary glandular growth becomes **atypical hyperplasia** (abnormal, precancerous cells), the rate of progression to invasive cancer follows a clear timeline according to longitudinal data from the [Mayo Clinic](#):

Timeframe Post-Atypia Detection	Absolute Percentage That Develops Into Breast Cancer
Within 5 Years	Approximately 7% of individuals will face a cancer diagnosis.
Within 10 Years	Approximately 13% of individuals will face a cancer diagnosis.
Within 25 Years	Cumulative incidence approaches 30% (nearly 1 in 3 women).



6. Clinical and Regulatory Recommendations

To address the systemic health risks posed by ASR-derived toxic compounds, public health frameworks and recycling industries must adopt several proactive interventions:

1. **Advanced Particulate Capture Protocols:** Industrial automobile shredder facilities must deploy high-efficiency particulate air (HEPA) filtration arrays and continuous dust suppression systems to prevent the atmospheric dispersion of shredded foam, plastic, and upholstery dust into local airsheds.
2. **Targeted Wastewater Management:** Implementation of multi-stage chemical precipitation and granular activated carbon (GAC) filtration systems at ELV sites to capture dissolved PFAS, phthalates, and heavy metals prior to storm or industrial runoff discharges.
3. **Endocrine-Disrupting Toxicity Screenings:** Incorporating environmental exposure histories into standard gynecological intakes, particularly for patients presenting with simultaneous uterine fibroid and adenomyosis phenotypes who reside near heavy manufacturing or vehicle recycling corridors.

Thank you to all out there working hard on environmental and industrial pollution management.

My personal experiences from exposure to pollution at a young age by 3M (who lied to us about its toxic impacts), then as an adult from my efforts to seek technology to manage PFAS pollutants, then desire to help with post 9/11 attack pollution exposure on people and the planet, knowing similar pollution was ongoing from recycling plants (including in Oakland and L.A. California) and later the impact of post disaster and war torn impacts on people's health (including in Gaza and Ukraine) have been a driver behind the long-term research, exposure to these pollutants to help reduce cancer rates from exposure once we deploy the Circular Solar Integrated Remediation System.

Cherise Petker, CircularSolar.net