2 of 2

(802) 540-0148 https://www.biadiagnostics.com/ Lic#TLAB0029



Sample ID: BIA250321S0003 Strain: Stink Palm

Matrix: Plant Type: Flower - Cured Sample Size: 6.44 g Lot#:

Produced: Collected: Received: 03/21/2025 Completed: 03/28/2025 Batch#: LOT 13

Client Mr Tree Lic. # CLTV0364 57 Commerce AVE South Burlington, VT 05403

Completed **Terpenes**

| | 100 | | . |
|------------------------|-------|---|---------------------|
| Analyte | LOQ | Results | Results |
| | mg/g | mg/g | % |
| Limonene | 0.010 | 6.686 | 0.669 |
| Ocimene | 0.010 | 4.016 | 0.402 |
| β-Myrcene | 0.010 | 3.525 | 0.353 |
| Linalool | 0.010 | 2.902 | 0.290 |
| β-Pinene | 0.010 | 2.389 | 0.239 |
| β-Caryophyllene | 0.010 | 1.945 | 0.195 |
| α-Pi <mark>nene</mark> | 0.010 | 1.422 | 0.142 |
| α-Humulene | 0.010 | 0.650 | 0.065 |
| Terpinolene | 0.010 | 0.442 | 0.044 |
| Camphene | 0.010 | 0.264 | 0.026 |
| y-Terpinene | 0.010 | 0.024 | 0.002 |
| α-Terpinene | 0.010 | 0.018 | 0.002 |
| 3-Carene | 0.010 | 0.013 | 0.001 |
| α-Bisabolol | 0.010 | <loq< td=""><td><loq< td=""></loq<></td></loq<> | <loq< td=""></loq<> |
| Caryophyllene Oxide | 0.010 | <loq< td=""><td><loq< td=""></loq<></td></loq<> | <loq< td=""></loq<> |
| cis-Nerolidol | 0.010 | <loq< td=""><td><loq< td=""></loq<></td></loq<> | <loq< td=""></loq<> |
| Eucalyptol | 0.010 | <loq< td=""><td><loq< td=""></loq<></td></loq<> | <loq< td=""></loq<> |
| Geraniol | 0.010 | <loq< td=""><td><loq< td=""></loq<></td></loq<> | <loq< td=""></loq<> |
| Guaiol | 0.010 | <loq< td=""><td><loq< td=""></loq<></td></loq<> | <loq< td=""></loq<> |
| Isopulegol | 0.010 | <loq< td=""><td><loq< td=""></loq<></td></loq<> | <loq< td=""></loq<> |
| p-Cymene | 0.010 | <loq< td=""><td><loq< td=""></loq<></td></loq<> | <loq< td=""></loq<> |
| trans-Nerolidol | 0.010 | <loq< td=""><td><loq< td=""></loq<></td></loq<> | <loq< td=""></loq<> |
| Total | | 24.296 | 2.430 |
| Aromas | | | |

Primary Aromas











Analyst: 045

LOQ = The lowest quantity this method can reliably detect. Any terpene that was not detected is assumed to be less than the stated LOQ

Terpene Methodology: Headspace Sampler, Gas Chromatography-Mass Spectrometry (GC-MS), using Perkin Elmer Clarus® SQ8 GC MS Reagent Blanks: < LOQs for all analytes

All results reflect dry weight of material, based on % moisture of the sample.

All moisture and water activity analysis is determined by dewpoint measurement using an AQUALAB water activity meter.



Luke Emerson-Mason Laboratory Director

Confident LIMS All Rights Reserved coa.support@confidentlims.com (866) 506-5866 www.confidentlims.com



03/28/2025