

Classified for
 "RESTRICTED USE"
 in New York State

under 6NYCRR Part 326



NET WEIGHT
.773 POUND

Active Ingredient	By Weight
*Acephate	97.4%
Inert Ingredients	2.6%
Total	100.0%

*O,S-Dimethyl acetylphosphoramidothioate
 U.S. Pat. No. 6,013,272

Manufactured for: **AMVAC Chemical Corporation**
 4695 MacArthur Court, Suite 1200, Newport Beach, CA 92660 U.S.A.
 EPA Reg. No. 5481-8978 EPA Est. No. 39578-TX-01

ACCEPTED
 VIA NOTIFICATION
 LABEL NOT REVIEWED

July 11, 2019

New York State Department
 of Environmental Conservation
 Division of Materials Management
 Pesticide Product Registration

KEEP OUT OF REACH OF CHILDREN
CAUTION
 SEE SIDE PANEL AND PAMPHLET FOR
 ADDITIONAL PRECAUTIONARY STATEMENTS AND DIRECTIONS FOR USE

ORTHENE® 97

SOLUBLE INSECTICIDE

(continued)
RESULTS FROM THE USE OF THIS PRODUCT IN ANY MANNER WHICH IS INCONSISTENT WITH THE LABEL DIRECTIONS, WARNINGS OR CAUTIONS.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW BUYER'S EXCLUSIVE REMEDY AND MANUFACTURER'S OR SELLER'S EXCLUSIVE LIABILITY FOR ANY AND ALL CLAIMS, LOSSES, DAMAGES, OR INJURIES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER OR NOT BASED IN TORT, CONTRACT, NEGLIGENCE, STRICT LIABILITY IN TORT OR OTHERWISE, SHALL BE LIMITED, AT THE MANUFACTURER'S OPTION, TO REPLACEMENT OF, OR THE REPAYMENT OF THE PURCHASE PRICE FOR, THE QUANTITY OF PRODUCT WITH RESPECT TO WHICH DAMAGES ARE CLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW MANUFACTURER OR SELLER SHALL NOT BE LIABLE FOR SPECIAL, INDIRECT OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

AMVAC offers this product, and Buyer accepts it, subject to the foregoing Limited Warranty which may be varied only by agreement in writing signed by an authorized representative of AMVAC.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage, disposal or cleaning of equipment.
PESTICIDE STORAGE: Keep pesticide in original container. Do not put concentrate or dilute into food or drink containers. Store in a cool, dry place. Protect from excessive heat. Do not contaminate food or foodstuffs. Do not store or transport near feed or food. For help with any spill, leak, fire or exposure involving this material, call day or night 1-888-462-6822.
PESTICIDE DISPOSAL: To avoid wastes, use all materials in this container by application according to label directions. If wastes cannot be avoided, offer remaining product to a waste disposal facility or pesticide disposal program (often such programs are run by state or local governments or by industry).
CONTAINER DISPOSAL: Nonrefillable container. Do not reuse or refill this container. Completely empty canister into application equipment. Do not reuse container. Dispose of empty canister in a sanitary landfill or by incineration, or offer for recycling, if available, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

©2019 AMVAC Chemical Corporation is a wholly owned subsidiary of American Vanguard Corporation. All Rights Reserved. AMERICAN VANGUARD, AMVAC and their respective logos are trademarks owned by AMVAC Chemical Corporation. Orthene is a trademark of OMS Investments, Inc. CHEMTREC is a service mark of the American Chemistry Council, Inc.

(continued)
 also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.
 Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 24 hours. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls, chemical-resistant gloves made of any waterproof material and shoes plus socks.

NON-AGRICULTURAL USE REQUIREMENTS
 The requirements in this box apply to uses of this product that are **NOT** within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses.
 Do not enter or allow others to enter until sprays have dried.

USE RESTRICTIONS
 • Not for indoor residential use.

CHEMIGATION
Apply to canneries only by sprinkler irrigation systems. Do not apply by chemigation to any other crop or this crop using any other type of irrigation system.

LIMITED WARRANTY AND DISCLAIMER
 The manufacturer warrants (a) that this product conforms to the chemical description on the label; and (b) that the directions, warnings, and other statements on this label are based upon responsible experts' evaluations of reasonable tests of effectiveness, of toxicity to laboratory animals and to plants and residues on food crops, and upon reports of field experience. Tests have not been made on all varieties of food crops and plants, or in all states or under all conditions. THIS WARRANTY DOES NOT EXTEND TO THE USE OF THIS PRODUCT CONTRARY TO LABEL INSTRUCTIONS, OR UNDER CONDITIONS NOT REASONABLY FORESEEABLE. **THESE ARE NO EXPRESS WARRANTIES OTHER THAN THOSE SET FORTH HEREIN. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW THE MANUFACTURER NEITHER MAKES NOR INTENDS, NOR DOES IT AUTHORIZE ANY AGENT OR REPRESENTATIVE, TO MAKE ANY OTHER WARRANTIES, EXPRESS OR IMPLIED, AND IT EXPRESSLY EXCLUDES AND DISCLAIMS ALL IMPLIED WARRANTIES OF MERCHANTABILITY OF FITNESS FOR A PARTICULAR PURPOSE, OR ANY WARRANTY OF QUALITY OR PERFORMANCE. THIS WARRANTY DOES NOT EXTEND TO, AND THE BUYER SHALL BE SOLELY RESPONSIBLE FOR, ANY AND ALL LOSS OR DAMAGE WHICH**

(continued)

Engineering Controls: Pilots must use an enclosed cockpit in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(6)].
 The use of human flaggers is prohibited.
 Follow the manufacturer's instructions for cleaning/maintaining PPE. If no instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS
Users should:
 • Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
 • Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
 • Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS: This pesticide is toxic to birds. For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwaters. Cover with soil or soil-incorporate spills.
 Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance, contact your State Water Board or Regional Office of the EPA.

This product and its degradate are highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds while bees are foraging the treatment area.
DIRECTIONS FOR USE
 It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.
 Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS
 Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. (continued)

FIRST AID	
Acephate is an organophosphate, cholinesterase inhibitor.	
If swallowed:	<ul style="list-style-type: none"> Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything by mouth to an unconscious person.
If in eyes:	<ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
If on skin or clothing:	<ul style="list-style-type: none"> Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
If inhaled:	<ul style="list-style-type: none"> Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

EMERGENCY INFORMATION
 Have the product container or label with you when calling a poison control center or doctor, or going for treatment. FOR THE FOLLOWING EMERGENCIES, PHONE 24 HOURS A DAY:
 For Medical Emergencies phone: 1-888-681-4261
 For Transportation Emergencies, including spill, leak or fire, phone: CHEMTREC® 1-800-424-9300
 For Product Use Information phone: AMVAC® 1-888-462-6822

NOTE TO PHYSICIAN
 This material contains a cholinesterase inhibitor. Measurement of blood cholinesterase activity may be useful in monitoring exposure but decisions regarding treatment will usually need to be made before test results are available. If signs of cholinesterase inhibition appear, atropine sulfate is antidotal. 2-PAM (PRÓTOPAM) is also antidotal and may be used in conjunction with atropine but should not be used alone.

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS & DOMESTIC ANIMALS
CAUTION: Harmful if swallowed.
PERSONAL PROTECTIVE EQUIPMENT (PPE):
 Some materials that are chemical-resistant to this product are butyl rubber, nitrile rubber, or neoprene. If you want more options, follow the instructions for category A on an EPA chemical-resistance category selection chart.
Mixers, Loaders, Applicators and Other Handlers must wear: Long-sleeved shirt and long pants, chemical-resistant gloves and shoes plus socks. **In addition, all Mixers and Loaders and all Applicators using low pressure hand wand application equipment must wear:** A NIOSH-approved dust mist filtering respirator with MSHA/NIOSH approval number prefix TC-21C or a NIOSH-approved respirator with any N, R, P or HE filter. See **Engineering Controls for additional requirements.**

TABLE 6. COTTON TANK MIXES WITH PYRETHROIDS—continued

Table with 5 columns: APPLICATION METHOD, PESTS CONTROLLED, ORTHENE 97 AND TANK MIX PARTNER, AMOUNT OF ORTHENE 97 + TANK MIX PARTNER PER ACRE, REMARKS, DAYS TO HARVEST. Includes rows for FOLIAR and For Aerial Application with various tank mix partners like ASANA XL, BAYTHROID 2 EC, etc.

For repeat applications, use a minimum spray interval of 7 days.

RESISTANCE MANAGEMENT

Cotton pest control programs, especially those for control of Silverleaf Whitefly populations, should employ a properly designed resistance management strategy. Such resistance management strategies include mixture or rotation of alternative classes of chemistry including organophosphates, carbamates, pyrethroids or insect growth regulators.

CRANBERRY

- Limit to one application per growing season.
Do not apply more than 1.0 lb. ORTHENE 97 (1.0 lb. ai) per season.
Do not apply from start of bloom until all berries set.

TABLE 7. CRANBERRY

Table with 5 columns: APPLICATION METHOD, PESTS CONTROLLED, RATE OF ORTHENE 97 PER ACRE, REMARKS, DAYS TO HARVEST. Includes rows for FOLIAR and FOLIAR By Sprinkler with various pest controls like Cranberry Blossom, Worm, etc.

Chemigation: Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the label-prescribed safety devices for public water systems are in place.

A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.

The pesticide injection pipeline must contain a functional, automatic, quick closing check valve to prevent the flow of fluid back toward the injection pump.

The pesticide injection pipeline must also contain a functional, normally closed solenoid operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.

The irrigation line or water pump must include a functional pressure switch, which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Do not apply when wind speed favors drift beyond the area intended for treatment.

Solid Set System: Apply specified dosage for the entire length of the irrigation period, or for a 30 to 60 minute period at the end of a regular irrigation set, or as a 30 to 60 minute injection as a separate application not associated with a regular irrigation. Allow time for all lines to flush the pesticide through all nozzles before turning off irrigation water.

Center Pivot Systems: Inject the specified dosage per acre continuously for one complete revolution of the system. See NOTE.

NOTE: Constant agitation must be maintained in the chemical supply tank during the entire period of insecticide application. Inject the product with a positive displacement pump into the main line ahead of a right turn to ensure adequate mixing.

Application of more than label recommended quantities of irrigation water per acre may result in decreased product performance by removing the chemical from the zone of effectiveness.

HEAD LETTUCE - Crisphead Type Only

- Do not apply more than 2 1/8 lbs./A (2 lbs. ai) ORTHENE 97 per crop cycle.
Do not feed trimmings to livestock or allow animals to graze on treated areas.
AZ & CA, (desert areas): Do not apply after first head begins to form in crops which germinate from mid-September through November.
For repeat applications, use a minimum spray interval of 3 days for application rates up to and including 1/2 lb. product/A and a minimum of 7 days for rates exceeding 1/2 lb./A.

TABLE 8. HEAD LETTUCE - Crisphead Type Only

Table with 5 columns: APPLICATION METHOD, PESTS CONTROLLED, RATES OF ORTHENE 97 PER ACRE, REMARKS, DAYS TO HARVEST. Includes rows for FOLIAR with pest controls like Astar Leafhopper, Green Peach Aphid, etc.

MINT - Peppermint and Spearmint

- Do not apply more than 2 1/8 lbs./A (2 lbs. ai) ORTHENE 97 per crop cycle.
Do not graze treated areas.
Do not feed spent mint hay to animals.

TABLE 9. MINT - Peppermint and Spearmint

Table with 5 columns: APPLICATION METHOD, PESTS CONTROLLED, RATES OF ORTHENE 97 PER ACRE, REMARKS, DAYS TO HARVEST. Includes rows for FOLIAR with pest controls like Alfalfa Looper, Aphid, Cutworm, etc.

NON-BEARING CITRUS - In Agricultural Settings (Newly Planted Groves)

- Do not graze treated areas.
Do NOT HARVEST citrus for one year (365 days) after treatment.

TABLE 10. NON-BEARING CITRUS

Table with 5 columns: APPLICATION METHOD, PESTS CONTROLLED, RATE OF ORTHENE 97 PER ACRE, REMARKS. Includes rows for FOLIAR and SOIL MOUND TREATMENT DRENCH METHOD with pest controls like Aphid, Grasshopper, etc.

* Not registered in CA for control of Citrus Blackfly and Harvester Ants.

NON-CROP AREAS

Field Borders, Fencerows, Roadsides, Ditchbanks, Borrow Pits

- Do not graze or feed vegetation cut from treated areas.

TABLE 11. NON-CROP AREAS - Field Borders, Fencerows, Roadsides, Ditchbanks, Borrow Pits

Table with 5 columns: APPLICATION METHOD, PESTS CONTROLLED, RATES OF ORTHENE 97 PER ACRE, REMARKS. Includes row for FOLIAR with pest control Grasshopper.

NON-CROP AREAS - Wasteland

- Do not make more than one application ORTHENE 97 per season.
Do not graze or feed vegetation cut from treated areas.

TABLE 12. NON-CROP AREAS - Wasteland

Table with 5 columns: APPLICATION METHOD, PESTS CONTROLLED, RATES OF ORTHENE 97 PER ACRE, REMARKS. Includes row for FOLIAR with pest controls like Black Grass, Bugs, etc.

PEANUTS

- Do not feed treated forage or hay to livestock.
Do not allow animals to graze on treated areas.
Do not apply more than 4 1/8 lbs./A (4 lbs. ai) ORTHENE 97 per crop cycle. This includes the use of ORTHENE 97 on peanuts as a foliar spray in addition to the peanut planterbox seed treatment use.
For repeat applications, use a minimum spray interval of 3 days for application rates up to and including 1/2 lb. product/A and a minimum of 7 days for rates exceeding 1/2 lb. product/A.

TABLE 13. PEANUTS

Table with 5 columns: APPLICATION METHOD, PESTS CONTROLLED, RATES OF ORTHENE 97 PER ACRE, REMARKS, DAYS TO HARVEST. Includes rows for FOLIAR with pest controls like Grasshopper, Thrips, etc.

PEPPERS - Bell

- Do not apply more than 2 1/8 lbs./A (2 lbs. ai) ORTHENE 97 per crop cycle.
For repeat applications, use a minimum spray interval of 3 days for application rates up to and including 1/2 lb. product/A and a minimum of 7 days for rates exceeding 1/2 lb. product/A.

TABLE 14. PEPPERS - Bell

Table with 5 columns: APPLICATION METHOD, PESTS CONTROLLED, RATES OF ORTHENE 97 PER ACRE, REMARKS, DAYS TO HARVEST. Includes row for FOLIAR with pest controls like Grasshopper, Cabbage Looper, etc.

PEPPERS - Non-Bell

FOR USE IN MIDWESTERN, EASTERN STATES AND PUERTO RICO ONLY.

- Do not apply more than 1 lb./A (1 lb. ai) ORTHENE 97 per crop cycle.

TABLE 15. PEPPERS - Non-Bell

Table with 5 columns: APPLICATION METHOD, PESTS CONTROLLED, RATE OF ORTHENE 97 PER ACRE, REMARKS, DAYS TO HARVEST. Includes row for FOLIAR with pest control Aphid.

SOYBEANS

- Do not graze or cut vines for hay or forage.
Do not apply more than 1.5 lbs./A (1.5 lbs. ai/A) of ORTHENE 97 per season.
Do not apply ORTHENE 97 within 14 days of harvest.
For rates up to 0.5 lb./A, at least 3 days must pass between applications of ORTHENE 97.
For rates between 0.5 lb. product/A and 1.0 lb./A, at least 7 days must pass between applications of ORTHENE 97.
Always read and follow all label directions, restrictions and precautions when using any pesticide alone or in tank mix combinations. The most restrictive labeling applies when using a tank mix.

TABLE 16. SOYBEANS

Table with 5 columns: APPLICATION METHOD, PESTS CONTROLLED, RATES OF ORTHENE 97 PER ACRE, REMARKS, DAYS TO HARVEST. Includes rows for FOLIAR with pest controls like Grasshopper, Thrips, etc.

TOBACCO (Except CA) - Transplant Water Application

- Do not apply more than 4 1/8 lbs./A (4 lbs. ai) ORTHENE 97 per crop cycle. This includes the use of ORTHENE 97 in transplant water, plant bed, soil, float bed (greenhouse) and foliar applications.

TABLE 17. TOBACCO (Except CA) - Transplant Water Application

Table with 5 columns: CROP, PESTS CONTROLLED, AMOUNT ORTHENE 97 PER ACRE, FURTHER USE INSTRUCTIONS, DAYS TO HARVEST. Includes row for Tobacco with pest controls like Cutworm, Flea Beetle, etc.

TOBACCO (Except CA) - Foliar Application

- For repeat applications use a minimum spray interval of 3 days for application rates up to and including 1/2 lb. product/A and a minimum of 7 days for rates exceeding 1/2 lb. product/A.
Do not apply more than 4 1/8 lbs./A (4 lbs. ai) ORTHENE 97 per crop cycle. This includes the use of ORTHENE 97 in transplant water, plant bed, soil, float bed (greenhouse) and foliar applications.

TABLE 18. TOBACCO (Except CA) - Foliar Application

Table with 5 columns: CROP, PESTS CONTROLLED, AMOUNT ORTHENE 97 PER ACRE, FURTHER USE INSTRUCTIONS, DAYS TO HARVEST. Includes rows for Tobacco with pest controls like Grasshopper, Flea Beetle, etc.

TOBACCO (Except CA) - Plant Bed Application

- Do not apply more than 4 1/8 lbs./A (4 lbs. ai) ORTHENE 97 per crop cycle. This includes the use of ORTHENE 97 in transplant water, plant bed, soil, float bed (greenhouse) and foliar applications.

TABLE 19. TOBACCO (Except CA) - Plant Bed Application

Table with 5 columns: CROP, PESTS CONTROLLED, AMOUNT ORTHENE 97 PER ACRE, FURTHER USE INSTRUCTIONS, DAYS TO HARVEST. Includes row for Tobacco Bed Treatment with pest controls like Cutworm, Flea Beetle, etc.

TOBACCO (Except CA) - Greenhouse Application

- Do not apply more than 4 1/8 lbs./A (4 lbs. ai) ORTHENE 97 per crop cycle. This includes the use of ORTHENE 97 in transplant water, plant bed, soil, float bed (greenhouse) and foliar applications.

TABLE 20. TOBACCO (Except CA) Greenhouse Application

Table with 5 columns: CROP, PESTS CONTROLLED, AMOUNT ORTHENE 97 PER ACRE, FURTHER USE INSTRUCTIONS, DAYS TO HARVEST. Includes row for Tobacco with pest controls like Cutworm, Flea Beetle, etc.

TOBACCO (Except CA) - Soil Application

- Do not apply more than 4 1/8 lbs./A (4 lbs. ai) ORTHENE 97 per crop cycle. This includes the use of ORTHENE 97 in transplant water, plant bed, soil, float bed (greenhouse) and foliar applications.
Treat maximum of 13 pounds per acre.
Do not treat more than once per season.

TABLE 21. TOBACCO (Except CA) - Soil Application

Table with 5 columns: CROP, PESTS CONTROLLED, AMOUNT ORTHENE 97 PER ACRE, FURTHER USE INSTRUCTIONS, DAYS TO HARVEST. Includes row for Tobacco with pest controls like Ants, Drench Method, etc.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage, disposal or cleaning of equipment.
PESTICIDE STORAGE: Keep pesticide in original container. Do not put concentrate or dilute into food or drink containers. Store in cool, dry place. Protect from excessive heat. Do not contaminate food or foodstuffs. Do not store or transport near food or feed.
PESTICIDE DISPOSAL: To avoid wastes, use all materials in this container by application according to label directions. If wastes cannot be avoided, offer remaining product to a waste disposal facility or pesticide disposal program (often such programs are run by state or local governments or by industry).

LIMITED WARRANTY AND DISCLAIMER

The manufacturer warrants (a) that this product conforms to the chemical description on the label; and (b) that the directions, warnings, and other statements on this label are based upon responsible experts' evaluations of reasonable tests of effectiveness, of toxicity to laboratory animals and to plants and residues on food crops, and upon reports of field experience. Tests have not been made on all varieties of food crops and plants, or in all states or under all conditions. THIS WARRANTY DOES NOT EXTEND TO THE USE OF THIS PRODUCT CONTRARY TO LABEL INSTRUCTIONS, OR UNDER CONDITIONS NOT REASONABLY FORESEEABLE.

THERE ARE NO EXPRESS WARRANTIES OTHER THAN THOSE SET FORTH HEREIN. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW THE MANUFACTURER NEITHER MAKES NOR INTENDS, NOR DOES IT AUTHORIZE ANY AGENT OR REPRESENTATIVE, TO MAKE ANY OTHER WARRANTIES, EXPRESS OR IMPLIED, AND IT EXPRESSLY EXCLUDES AND DISCLAIMS ALL IMPLIED WARRANTIES OF MERCHANTABILITY OF FITNESS FOR A PARTICULAR PURPOSE, OR ANY WARRANTY OF QUALITY OR PERFORMANCE. THIS WARRANTY DOES NOT EXTEND TO, AND THE BUYER SHALL BE SOLELY RESPONSIBLE FOR, ANY AND ALL LOSS OR DAMAGE WHICH RESULTS FROM THE USE OF THIS PRODUCT IN ANY MANNER WHICH IS INCONSISTENT WITH THE LABEL DIRECTIONS, WARNINGS OR CAUTIONS.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW BUYER'S EXCLUSIVE REMEDY AND MANUFACTURER'S OR SELLER'S EXCLUSIVE LIABILITY FOR ANY AND ALL CLAIMS, LOSSES, DAMAGES, OR INJURIES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER OR NOT BASED IN TORT, CONTRACT, NEGLIGENCE, STRICT LIABILITY IN TORT OR OTHERWISE, SHALL BE LIMITED, AT THE MANUFACTURER'S OPTION, TO REPLACEMENT OF, OR THE REPAYMENT OF THE PURCHASE PRICE FOR, THE QUANTITY OF PRODUCT WITH RESPECT TO WHICH DAMAGES ARE CLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW MANUFACTURER OR SELLER SHALL NOT BE LIABLE FOR SPECIAL, INDIRECT OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

AMVAC offers this product, and Buyer accepts it, subject to the foregoing Limited Warranty which may be varied only by agreement in writing signed by an authorized representative of AMVAC.

© 2019 AMVAC Chemical Corporation is a wholly owned subsidiary of American Vanguard Corporation. All Rights Reserved. AMERICAN VANGUARD, AMVAC and their respective logos are trademarks owned by AMVAC Chemical Corporation; ORTHENE is a trademark of OMS Investments, Inc. ASANA and DANITOL are trademarks of Sumitomo Chemical Co., Ltd. AMMO and CAPTURE are trademarks of Corporation. BAYTHROID and Scout X-Tra are trademarks of a Bayer Company; CHEMTREC is a service mark of the American Chemistry Council, Inc. KARATE is a trademark of Syngenta Ltd. LORSPAN is a trademark of Dow AgroSciences, LLC. EGGS/DEADHATCH is a trademark of Valent U.S.A. Corp.

