

Quick Start Guide

QUICK START GUIDE

Opioid Overdose Response Instructions

1 Identify Opioid Overdose and Check for Response



Ask person if he or she is okay and shout name.

Check for signs of opioid overdose:

- Will not wake up or respond to your voice or touch
- Breathing is very slow, irregular, or has stopped
- Center part of their eye is very small, sometimes called "pin-point pupils"


Lay the person on their back to receive a dose of Naloxone HCl Nasal Spray

2 Give Naloxone HCl Nasal Spray

Remove Naloxone HCl Nasal Spray from the box.



Peel back the tab with the circle to open the Naloxone HCl Nasal Spray



Hold the Naloxone HCl Nasal Spray with your thumb on the bottom of the plunger and your first and middle fingers on either side of the nozzle.



Gently insert the tip of the nozzle into either nostril.

- Tilt the person's head back and provide support under the neck with your hand. Gently insert the tip of the nozzle into one nostril, until your fingers on either side of the nozzle are against the bottom of the person's nose.



Press the plunger firmly to give the dose of Naloxone HCl Nasal Spray.

- Remove the Naloxone HCl Nasal Spray from the nostril after giving the dose.

3 Call for emergency medical help, Evaluate, and Support



Get emergency medical help right away.

Move the person on their side (recovery position) after giving Naloxone HCl Nasal Spray.

Watch the person closely.

If the person does not respond by waking up, to voice or touch, or breathing normally another dose may be given. Naloxone HCl Nasal Spray may be dosed every 2 to 3 minutes, if available.

Repeat Step 2 using a new Naloxone HCl Nasal Spray to give another dose in the other nostril.

If additional Naloxone HCl Nasal Sprays are available, repeat step 2 every 2 to 3 minutes until the person responds or emergency medical help is received.