

Date		Microcystin	
Sampled	Label	(ug/L)	Site
6/29/2023	Α	<0.05*	North Slalom Course
	В	<0.05*	Mid Lake
	C	<0.05*	Dam
	D	<0.05*	Shoreline Hawthorn Docks
		* Concentration	n below limit of quantification
		Method: ELISA Microcystin SAES	
		Method detection limit 0.05 ug/L	

Date		Anatoxin		
Sampled	Label	(ug/L)	Site	
6/29/2023	В	<0.15*	Mid Lake	
* Concentration below limit of quantificatio				
Method: ELISA Anatoxin				
	Method detection limit 0.15 ug/L			

Date		Saxitoxin		
Sampled	Label	(ug/L)	Site	
6/29/2023	В	<0.02*	Mid Lake	
		* Concentration below limit of quantification		
	Method: ELISA Saxitoxin			
		Method detec	tion limit 0.02 ug/L	

Date	Cylindrospermopsin			
Sampled	Label	(ug/L)	Site	
6/29/2023	В	<0.05*	Mid Lake	
	* Concentration below limit of quantification			
Method: ELISA Cylindrospermopsin				
	Method detection limit 0.05 ug/L			

## 2.3 Cyanotoxin Thresholds for Recreational Waters

Numerous risk assessment frameworks, exposure assumptions, and toxicity values from state, national, and primary literature sources were considered prior to developing the cyanotoxin thresholds. The following thresholds were established based on the best scientific information, guidance, and public policy available at the time, and are based on incidental ingestion only (Table 2).

While protective of human exposures based on current information, the thresholds given here may or may not be protective of animals such as dogs or livestock. The United States Environmental Protection Agency (U.S. EPA) issued final recommended recreational swimming advisories for two cyanotoxins, microcystins and cylindrospermopsin in June 2019 which were subsequently adopted by the State of Ohio in this document.

For a toxicity review of various cyanotoxins, exposure assumptions and threshold calculations, see Appendix D.

Table 2 — Numeric Thresholds for Cyanotoxins in Recreational Water.

Threshold (µg/L)		Anatoxin-a	Cylindrospermopsin	
Recreational Public Health Advisory	8	8	15	0.8

<sup>\*</sup>Microcystins and saxitoxin thresholds are intended to be applied to total concentrations of all reported congeners, variants, or analogs of those cyanotoxins.