

are not systematically losing (or gaining) galaxies at high redshift because of photometric errors or uncertainties, then Loh and Spillar have succeeded in measuring the volume element as a function of redshift. It does not matter whether the mass in the Universe is luminous or dark. This measurement of the geometry of the Universe yields a determination of $\Omega = 0.9^{+0.2}_{-0.5}$ (95 per cent confidence limits) and is consistent with the inflationary prediction of an Einstein-de Sitter cosmology ($\Omega = 1$).

This result is especially noteworthy because it is the first, seemingly unambiguous astronomical determination of Ω that yields a value of unity on large ($\sim 1,000$ megaparsecs) scales. Before accepting this as the new cosmological gospel, however, notes of caution have been urged. Redshift errors may be systematically underestimated because the 4,000-Å spectral indicator has only been calibrated against cluster galaxies, whereas this fea-

ture is known to be weaker for the field galaxies that dominate the Loh-Spillar sample (Richard Ellis, University of Durham, personal communication). There may be systematic errors in photometry that make the smaller, more distant galaxies appear brighter relative to the nearby galaxies, thereby mimicking the effect of high Ω (S.D. Jorgovski, personal communication). Dust in distant galaxies may also affect the photometric redshift determinations. These effects need to be critically evaluated, presumably by time-consuming spectroscopic studies, before the $\Omega = 1$ result can be considered to be definitive. For now, however, it remains a tantalizing possibility: perhaps observational cosmologists (and other astronomers) should pay more heed to theorists. □

Joseph Silk is in the Astronomy Department at the University of California, Berkeley, California 94720, U.S.A.

Mathematics

The Waring experience

from Ian Stewart

IN 1770 Joseph-Louis Lagrange proved a long-sought result in number theory, the four-squares theorem: every whole number is a sum of four perfect squares. Earlier that year Edward Waring, Lucasian Professor of Mathematics at Cambridge, had published his *Meditationes Algebraicae*. It contained a far-reaching generalization, stated without proof: every whole number is a sum of 9 cubes, 19 fourth powers "and so on". The implied conjecture, that for all n the number of perfect n th powers needed to represent any whole number is finite, became known as Waring's problem. The finiteness was proved by David Hilbert in 1909 using an identity in 25-fold multiple integrals, but his method did not yield the actual number of n th powers required. Other methods have been developed to estimate their number, and various workers have improved the upper bounds until they have become quite respectable. R.C. Vaughan has now obtained substantial improvements to these bounds when n lies between 5 and 9 (*Proc. Lond. Math. Soc.* 52, 445; 1986 and *J. Lond. Math. Soc.* 33, 227; 1986).

More precisely, let $g(n)$ be the smallest number such that every whole number is a sum of $g(n)$ n th powers. Then Lagrange's result shows that $g(2) \leq 4$. Because the number 7 actually requires all four squares ($7 = 1^2 + 1^2 + 1^2 + 2^2$ is the best we can do), $g(2)$ is exactly 4. Waring's conjecture is that $g(3) = 9$, $g(4) = 19$ and that $g(n)$ exists and is finite for all values of n .

He almost certainly hit on these numbers by trial and error. A search of all

numbers up to, say, a few thousand will show that at most 9 cubes appear to be needed; and only 2 numbers, 23 and 239, require all 9. In fact only 15 numbers seem to need 8, the largest of these being 454. In *An Introduction to the Theory of Numbers* (4th edn; Oxford University Press, 1960), G.H. Hardy and E.M. Wright comment as follows: "It is plain, if this be so, that 9 is

n	4	5	6	7	8	9	
$G(n) \leq$	19	41	87	193	425		Hardy & Littlewood (1928)
				137	163		Heilbronn (1936)
	16*	23	36				Davenport (1939, 1942)
				53	73		Narasimhamurti (1941)
		22	34	50	68	87	Thanigasalam (1982)
		21	31	45	62	82	Vaughan (1986)

* $G(4) = 16$ exactly.

not the number which is really most significant in the problem. The facts, if they be facts, that just 2 numbers require 9 cubes, and just 15 require 8, are, so to say, arithmetical flukes... The most fundamental and most difficult problem is that of deciding, not how many cubes are required for the representation of all numbers, but how many are required for the representation of all large numbers, that is, of all numbers with some finite number of exceptions".

Thus there is a second, deeper quantity, called $G(n)$, defined to be the smallest number of n th powers needed to represent any whole number, with finitely many exceptions. The "facts, if they be facts," imply that $G(3)$ is 7 (or less) rather than 9.

Because no number of the form $8k+7$ can be a sum of three squares, $G(2)$ is the same as $g(2)$, namely 4. But in general $G(n)$ will be smaller than $g(n)$.

Many authors have studied the functions $g(n)$ and $G(n)$, either for special values of n or for general ones. Thus, in 1942 Linnik showed that $G(3) \leq 7$ (so that the "facts, if they be facts" be facts). Between 1920 and 1928 Hardy and Littlewood developed an approach by way of complex function theory, and proved that $G(n)$ is never larger than $(n-2)2^{n-1}+5$. They improved this to yield $G(4) \leq 19$, $G(5) \leq 41$, $G(6) \leq 87$, $G(7) \leq 193$, $G(8) \leq 425$ and so on. By comparison the best results known by 1930 for $g(n)$ were 37, 58, 478, 3,806 and 31,353, respectively.

Vinogradov introduced powerful methods based on estimating trigonometric sums (a standard approach in analytic number theory but not an easy one to carry through), and in 1936 Heilbronn simplified the method to yield

$$G(n) \leq 6n \log n + [4 + 3 \log(3 + 2/n)]n + 3$$

whence $G(7) \leq 137$, $G(8) \leq 163$. A list of various subsequent improvements is presented in the table (for references see the works cited above). Only for $G(4)$ is the current bound known to be best possible.

Vaughan's methods are related to those of Hardy and Littlewood and Davenport, and involve very careful and delicate analytical estimates. They have applications to other questions in number theory, and may also shed light on Waring's problem for tenth and higher powers.

Although number theory may not be

the most directly useful part of mathematics, it exerts an endless fascination because of the gulf between the simplicity of its raw materials and the complexity of its methods. There is some practical payoff too (see, for example, M. Schröder's *Number Theory in Science and Communications* 2nd edn; Springer, New York, 1986). It is remarkable that a problem posed more than two centuries ago, based on simple empirical observations, continues to exercise the ingenuity of number theorists and to generate new mathematics. The Waring experience is far from over. □

Ian Stewart is in the Mathematics Institute, University of Warwick, Coventry CV4 7AL, UK.

nature

1986

NATURE VOL. 323 23 OCTOBER 1986

nature
SCIENTIFIC JOURNAL

CELL FUSION BY AIDS VIRUS GENE

Cells labelled with rhodamine isocyanate (red), infected with a recombinant vaccinia virus containing the envelope gene of HTLV-III/LAV and co-cultured with uninfected cells labelled with fluorescein isocyanate (green). The envelope protein causes the cells to fuse, producing large multinucleate giant cells that fluoresce orange-green. See page 725.

OPINION

- British science over the hill 655
 UNESCO'S new future? 656

NEWS

- US budget 657
 Unapproved drugs for export 658
 British science politics 658
 Soviet Union 659
 Mars spacecraft 659
 US innovation 660
 British research 660
 Genome sequencing 661
 Doubts about Department of Energy 661
 Contraceptive vaccines 662
 Soviet energy 662
 European microchips 663
 Nobel chemistry 663
 Soviet Antarctic research 663
 Nobel physics 663
 European research 663

CORRESPONDENCE

- Paranormal/Systematics/etc. 664

NEWS AND VIEWS

- Turbulence assails fifth force 665
 Theoretical chemistry: Back to the valence bond 666
 Roy McWeeney 666

- Nitrogen fixation: Finding a time and a place 667
 A E Walsby 667
 Melting and the surface 668
 Robert W Cahn 668
 Mantle geophysics: Variable viscosity makes waves 669
 David A Yuen 669
 Use of deuterated proteins in nuclear magnetic resonance 670
 Paul Röscher 670
 Nervous excitement over insect muscle patterns 671
 Vernon French 671
 Obituary: Fritz Lipmann 672
 William P Jencks 672
 Cosmology: Is omega equal to unity? 673
 Joseph Silk 673
 Mathematics: The Waring experience 674
 Ian Stewart 674

SCIENTIFIC CORRESPONDENCE

- Mystery solved 675
 B J Ford; A Ball 675
 Turing's fly 675
 D Gubb 675
 Imanishi's impact in Japan 675
 P J Asquith 675
 The northerly extent of Chernobyl contamination 676
 M Pourchet, J F Pinglot & J C Gascard 676

BOOK REVIEWS

- The Nazi Doctors: Medical Killing and the Psychology of Genocide 677
 by R J Lifton 677
 Benno Müller-Hill 677
 A Hotbed of Genius: The Scottish Enlightenment, 1730-90 678
 D Daiches, P Jones and J Jones, eds 678
 J V Golinski 678
 Hemoglobin: Molecular, Genetic and Clinical Aspects 679
 by H F Bunn and B G Forget 679
 D R Higgs 679
 The Evolution of the Soul 679
 by R Swinburne 679
 D M MacKay 679
 Transport and Diffusion Across Cell Membranes 680
 by W D Stein 680
 Clive Ellory 680

COMMENTARY

- National performance in basic research 681
 D C Smith, P M D Collins, D M Hicks & S Wyatt 681

REVIEW ARTICLE

- Stable and metastable metal surfaces in heterogeneous catalysis 685
 M S Spencer 685

ARTICLES

- A gene required for the specification of dorsal-ventral pattern in *Drosophila* appears to encode a serine protease 688
 R DeLotto & P Spierer 688

LETTERS TO NATURE

- Is there a common origin for the cosmic γ -ray lines at 0.51 and 1.81 MeV near the galactic centre? 692
 W R Webber, V Schönfelder & R Diehl 692

- Spluttering of sodium on the planet Mercury 694
 M A McGrath, R E Johnson & L J Lanzerotti 694

- Nature and variability of Mercury's sodium atmosphere 696
 W H Smyth 696

- The electronic structure of the benzene molecule 699
 D L Cooper, J Garratt & M Raimondi 699

- CO dissociation on potassium-promoted aluminium 701
 J Paul 701

- Icosahedral symmetry carbon cage molecules 703
 D J Klein, W A Seitz & T G Schmalz 703

- A new type of self-organization in combustion 706
 A I Lesnikov, V V Sviridov, G V Printsev, O A Ivashkevich & P N Gaponik 706

- Isotopic composition of osmium in terrestrial samples determined by accelerator mass spectrometry 707
 U Fehn, R Teng, D Elmore & P W Kubik 707

- K-Ar isochron dating of Zaire cubic diamonds 710
 S Zashu, M Ozima & O Nitoh 710

- Disoriented pathfinding by pioneer neurone growth cones deprived of filopodia by cytochalasin treatment 712
 D Bentley & A Toranion-Raymond 712
 Contents continued overleaf

Nature (ISSN 0028-0836) is published weekly on Thursday, except the last week in December, by Macmillan Journals Ltd and includes the Annual Index (mailed in February). Annual subscription for USA and Canada US \$250. USA and Canadian orders to: Nature, Subscription Dept, PO Box 1501, Neptune, NJ 07753, USA. Other orders to Nature, Brunel Road, Basingstoke, Hants RG21 2XS, UK. Second class postage paid at New York, NY 10012 and additional mailing offices. Authorization to photocopy material for internal or personal use, or internal or personal use of specific clients, is granted by Nature to libraries and others registered with the Copyright Clearance Center (CCC) Transactional Reporting Service, provided the base fee of \$1.00 a copy plus \$0.10 a page is paid directly to CCC, 21 Congress Street, Salem, Massachusetts 01970, USA. Identification code for Nature: 0028-0836/86 \$1.00 + \$0.10. US Postmaster: send address changes to Nature, 350 West 57th Street, New York, NY 10019.