


2022 Spring Snowpack Index

Paul Werth
Fire Weather Meteorologist
Weather Research and Consulting Services, LLC

The Spring Snowpack Index (SSPI) is a unique product developed by Weather Research and Consulting Services, LLC that has proven to be an effective predictor of wildland fire season severity. The SSPI combines Snow Water Equivalent (SWE) values and snowmelt rates from National Resources Conservation Service (NRCS) SNOTEL data resulting in a number that integrates both winter and spring weather (i.e. temperature, precipitation and evaporation). Our research indicates that it is a much better predictor of fire season severity than SWE alone because it is a measure of snowmelt date, fire season length, soil moisture and both live fuel and large dead fuel moisture. SSPI values have a wide range extending from -100 to 300 or higher. In general, the lower the SSPI value, the higher the probability of an active fire season.

The graphic to the right displays this year's SSPI calculated for river basins in the Western United States using NRCS SNOTEL data. The lowest SSPI values, less than minus 50, are located in Arizona, New Mexico, southern Utah and southern Nevada. Warm, dry and windy April weather produced rapid snowmelt across the Southwestern States. Unseasonably cool and wet weather significantly delayed snowmelt throughout the Pacific Northwest and Northern Rockies. The highest SSPI values, 150 or greater, are located in Washington, Oregon, Idaho and Montana.



Using updated algorithms that predict acres burned for each of the western states, the southern half of the region can expect another very active fire season. This includes the states of Arizona, New Mexico, California, Utah and Colorado. Meanwhile, the northern states of Washington, Oregon, Idaho, Montana and Wyoming will likely experience an average fire season due to the cool, wet spring and delayed snowmelt.

The table below lists the statewide SSPI averages for the past nine years. Figures in red indicate negative values with Arizona and New Mexico reporting the most. All states are showing higher SSPI values this year compared to last year indicating the winter snowpack was better and/or a below normal spring snowmelt rate. However, the improvement was negligible in Colorado, Utah, Arizona and New Mexico.

<u>State</u>	<u>2022</u>	<u>2021</u>	<u>2020</u>	<u>2019</u>	<u>2018</u>	<u>2017</u>	<u>2016</u>	<u>2015</u>	<u>2014</u>
MONTANA	125	91	101	109	154	151	67	56	166
WASHINGTON	149	104	86	73	140	146	48	24	125
WYOMING	103	92	91	83	116	155	125	49	145
IDAHO	123	28	61	112	78	200	55	9	113
CALIFORNIA	73	-6	23	165	36	317	73	4	9
COLORADO	53	43	68	122	34	94	117	49	94
OREGON	144	4	7	87	34	200	8	4	62
NEVADA	63	8	-23	132	4	192	92	-9	57
UTAH	20	12	47	152	-6	115	89	1	53
ARIZONA	-27	-77	-14	49	-7	4	-6	0	-9
NEW MEXICO	-18	-28	-12	97	-13	40	48	15	15
AVG	73	25	40	106	52	147	65	18	75

Data developed by Weather Research and Consulting Services, LLC

Paul Werth
 Fire Weather Meteorologist
 Weather Research and Consulting Services, LLC
www.fireweather.com or www.firewx.com