

## DISEASES &amp; CONDITIONS

# Osteoporosis

This article provides answers to some common questions about osteoporosis.

## What is osteoporosis?

Osteoporosis is a disease of progressive bone loss associated with an increased risk of fractures. The term osteoporosis literally means porous bone. The disease often develops unnoticed over many years, with no symptoms or discomfort until a fracture occurs. Osteoporosis often causes a loss of height and dowager's hump (a severely rounded upper back).

## Why should I be concerned about osteoporosis?

Osteoporosis is a major health problem, affecting more than 44 million Americans and contributing to an estimated 2 million bone fractures per year. According to the National Osteoporosis Foundation, the number of fractures due to osteoporosis may rise to over 3 million by the year 2025.



One in two women and one in four men older than 50 years will sustain bone fractures caused by osteoporosis. Many of these are painful fractures of the hip, spine, wrist, arm, and leg, which often occur as a result of a fall. However, performing even simple household tasks can result in a fracture of the spine if the bones have been weakened by osteoporosis.

The most serious and debilitating osteoporotic fracture is a hip fracture. Most patients who experience a hip fracture and previously lived independently will require help from their family or home care. All patients who experience a hip fracture will require walking aids for several months, and nearly half will permanently need canes or walkers to move around their house or outdoors. Hip fractures are expensive. Health care costs from hip fractures total more than \$11 billion annually, or about \$37,000 per patient.

## **What causes osteoporosis?**

Doctors do not know the exact medical causes of osteoporosis, but they have identified many of the major factors that can lead to the disease.

### ***Aging***

Everyone loses bone with age. After 35 years of age, the body builds less new bone to replace the loss of old bone. In general, the older you are, the lower your total bone mass and the greater your risk for osteoporosis.

### ***Heredity***

A family history of fractures; a small, slender body build; fair skin; and Caucasian or Asian ethnicity can increase the risk for osteoporosis. Heredity also may help explain why some people develop osteoporosis early in life.

### ***Nutrition and Lifestyle***

Poor nutrition, including a low calcium diet, low body weight, and a sedentary lifestyle have been linked to osteoporosis, as have smoking and excessive alcohol use.

### ***Medications and Other Illnesses***

Osteoporosis has been linked to the use of some medications, including steroids, and to other illnesses, including some thyroid problems.

## **What can I do to prevent osteoporosis or keep it from getting worse?**

To prevent osteoporosis, slow its progression, and protect yourself from fractures you should include adequate amounts of calcium and Vitamin D in your diet and exercise regularly.

### ***Calcium***

During the growing years, your body needs calcium to build strong bones and to create a supply of calcium reserves. Building bone mass when you are young is a good investment for your future. Inadequate calcium during growth can contribute to the development of osteoporosis later in life.

Whatever your age or health status, you need calcium to keep your bones healthy. Calcium continues to be an essential nutrient after growth because the body loses calcium every day. Although calcium cannot prevent gradual bone loss after menopause, it continues to play an essential role in maintaining bone quality. Even if women have gone through menopause or already have osteoporosis, increasing intake of calcium and Vitamin D can decrease the risk of fracture.

How much calcium you need will vary depending on your age and other factors. The National Academy of Sciences makes the following recommendations regarding daily intake of calcium:

- Males and females 9 to 18 years: 1,300 mg per day
- Women and men 19 to 50 years: 1,000 mg per day
- Pregnant or nursing women up to age 18: 1,300 mg per day
- Pregnant or nursing women 19 to 50 years: 1,000 mg per day
- Women and men over 50: 1,200 mg per day


Dairy products, including yogurt and cheese, are excellent sources of calcium. An eight-ounce glass of milk contains about 300 mg of calcium. Other calcium-rich foods include sardines with bones and green leafy vegetables, including broccoli and collard greens.

If your diet does not contain enough calcium, dietary supplements can help. Talk to your doctor before taking a calcium supplement.

## ***Vitamin D***

Vitamin D helps your body absorb calcium. The recommendation for Vitamin D is 200-600 IU (international units) daily. Supplemented dairy products are an excellent source of Vitamin D. (A cup of milk contains 100 IU of Vitamin D. A multivitamin contains 400 IU of Vitamin D.) Vitamin supplements can be taken if your diet does not contain enough of this nutrient. Again, consult with your doctor before taking a vitamin supplement. Too much Vitamin D can be toxic.

## ***Exercise Regularly***



Like muscles, bones need exercise to stay strong. No matter what your age, exercise can help minimize bone loss while providing many additional health benefits. Doctors believe that a program of moderate, regular exercise (3 to 4 times a week) is effective for the prevention and management of osteoporosis. Weight-bearing exercises such as walking, jogging, hiking, climbing stairs, dancing, treadmill exercises, and weight lifting are probably best. Falls account for 50% of fractures; therefore, even if you have low bone density, you can prevent fractures if you avoid falls. Programs that emphasize balance training, especially tai chi, should be emphasized. Consult with your doctor before beginning any exercise program.

## How is osteoporosis diagnosed?

The diagnosis of osteoporosis is usually made by your doctor using a combination of a complete medical history and physical examination, skeletal x-rays, bone densitometry, and specialized laboratory tests. If your doctor diagnoses low bone mass, he or she may want to perform additional tests to rule out the possibility of other diseases that can cause bone loss, including osteomalacia (a metabolic bone disease characterized by abnormal mineralization of bone) or hyperparathyroidism (overactivity of the parathyroid glands).

Loss of height and a stooped appearance of a person with osteoporosis results from partial collapse of weakened vertebrae.



Bone densitometry is a safe, painless x-ray technique that compares your bone density to the peak bone density that someone of your same sex and ethnicity should have reached at 20 to 25 years of age.

Bone densitometry is often performed in women at the time of menopause. Several types of bone densitometry are used today to detect bone loss in different areas of the body. Dual-energy x-ray absorptiometry (DEXA) is one of the most accurate methods, but other techniques can also identify osteoporosis, including single photon absorptiometry (SPA), quantitative computed tomography (QCT), radiographic absorptiometry, and ultrasound. Your doctor can determine which method is best suited for you.

## How is osteoporosis treated?

Because lost bone cannot be replaced, treatment for osteoporosis focuses on the prevention of further bone loss. Treatment is often a team effort involving a physician or internist, an orthopaedist, a gynecologist, and an endocrinologist.

Although exercise and nutrition therapy are often key components of a treatment plan for osteoporosis, there are other treatments as well.

## ***Estrogen Replacement Therapy***

Estrogen replacement therapy (ERT) is often recommended for women at high risk for osteoporosis to prevent bone loss and reduce fracture risk. A measurement of bone density when menopause begins may help you decide whether ERT is right for you. Hormones also prevent heart disease, improve cognitive functioning, and improve urinary function. ERT is not without some risk, including enhanced risk of breast cancer; the risks and benefits of ERT should be discussed with your doctor.

## ***Selective Estrogen Receptor Modulators***

New anti-estrogens known as SERMs (selective estrogen receptor modulators) can increase bone mass, decrease the risk of spine fractures, and lower the risk of breast cancer.

## ***Calcitonin***


Calcitonin is another medication used to decrease bone loss. A nasal spray form of this medication increases bone mass, limits spine fractures, and may offer some pain relief.

## ***Bisphosphonates***

Bisphosphonates, including alendronate, markedly increase bone mass and prevent both spine and hip fractures.

ERT, SERMs, calcitonin, and bisphosphonates all offer patient with osteoporosis an opportunity to not only increase bone mass, but also to significantly reduce fracture risk. Prevention is preferable to waiting until treatment is necessary.

Your orthopaedist is a medical doctor with extensive training in the diagnosis and nonsurgical and surgical treatment of the musculoskeletal system, including bones, joints, ligaments, tendons, muscles and nerves. This has been prepared by the American Academy of Orthopaedic Surgeons and is intended to contain current information on the subject from recognized authorities. However, it does not represent official policy of the Academy and its text should not be construed as excluding other acceptable viewpoints.



Information on this topic is also available as an OrthoInfo Basics PDF Handout.

For more information:

**Basics Handouts**

[\(/en/about-orthoinfo/OrthoInfo-Basics/basics-handouts/\)](/en/about-orthoinfo/OrthoInfo-Basics/basics-handouts/)

**Last Reviewed**

August 2009

AAOS does not endorse any treatments, procedures, products, or physicians referenced herein. This information is provided as an educational service and is not intended to serve as medical advice. Anyone seeking specific orthopaedic advice or assistance should consult his or her orthopaedic surgeon, or locate one in your area through the AAOS [Find an Orthopaedist](#) program on this website.