

Range of recommended ages for all children	Range of recommended ages for catch-up vaccination	Range of recommended ages for certain high-risk groups or populations	Recommended vaccination can begin in this age group	Recommended vaccination based on shared clinical decision-making	No Guidance/Not Applicable
--	--	---	---	--	----------------------------

Vaccine and other immunizing agents	Birth	1 mo	2 mos	4 mos	6 mos	9 mos	12 mos	15 mos
Respiratory syncytial virus ⓘ (RSV-mAb [Nirsevimab])	1 dose depending on maternal RSV vaccination status, See Notes					1 dose (8 through 19 months), See Notes		
Hepatitis B ⓘ (HepB)	1 st dose	← 2 nd dose →			← 3 rd dose →			
Rotavirus (RV) ⓘ RV1 (2-dose series); RV5 (3-dose series)			1 st dose	2 nd dose	See Notes			
Diphtheria, tetanus, & acellular pertussis ⓘ (DTaP: <7 yrs)			1 st dose	2 nd dose	3 rd dose			← 4 th dose →
Haemophilus influenzae type b ⓘ (Hib)			1 st dose	2 nd dose	See Notes		← 3 rd or 4 th dose, See Notes →	
Pneumococcal conjugate ⓘ (PCV15, PCV20)			1 st dose	2 nd dose	3 rd dose		← 4 th dose →	
Inactivated poliovirus (IPV) ⓘ			1 st dose	2 nd dose	← 3 rd dose →			
COVID-19 ⓘ (1vCOV-mRNA, 1vCOV-aPS)						1 or more doses of 2024–2025 vaccine (See Notes)		
Influenza (IIIV3, cclIV3) ⓘ						1 or 2 doses annually		
Influenza (LAIV3) ⓘ								
Measles, mumps, rubella ⓘ (MMR)						See Notes		← 1 st dose →
Varicella ⓘ (VAR)							← 1 st dose →	
Hepatitis A ⓘ (HepA)						(See Notes)		← 2-dose series, See Notes →
Tetanus, diphtheria, & acellular pertussis ⓘ (Tdap: ≥7 yrs)								
Human papillomavirus ⓘ (HPV)								
Meningococcal ⓘ (MenACWY-CRM ≥2 mos, MenACWY-TT ≥2years)			See Notes					
Meningococcal B ⓘ (MenB-4C, MenB-FHbp)								
Respiratory syncytial virus vaccine ⓘ (RSV [Abrysvo])								
Dengue ⓘ (DEN4CYD: 9-16 yrs)								
Mpox ⓘ								