



TRANSFORM FAT BURNER IV DRIP

BENEFITS

- **Enhanced Metabolism:** Contains a powerful blend of vitamins, amino acids, and antioxidants designed to boost your metabolism and support weight loss efforts.
- **Increased Energy Levels:** The combination of B vitamins and amino acids helps to increase energy levels, reduce fatigue, and improve overall stamina, aiding you in maintaining an active lifestyle.
- **Muscle Recovery and Growth:** Amino acids such as Ornithine, Arginine, Lysine, and Citrulline support muscle recovery and growth, making your workouts more effective and aiding in muscle preservation during weight loss.
- **Detoxification:** Glutathione, a potent antioxidant, helps to detoxify your body by neutralizing harmful free radicals and supporting liver function, promoting overall health and wellness.

SCIENCE

- **Lactated Ringer's/Sodium Chloride 0.9% (500-1000ml or Push 60ml):** This solution hydrates your body, replenishing essential electrolytes, and aiding in nutrient absorption and detoxification.
- **Glutathione (200mg/ml - 5ml):** Known as the "master antioxidant," glutathione protects cells from damage, supports immune function, and enhances metabolic processes.
- **Amino Blend (Ornithine 50mg, Arginine 100mg, Lysine 50mg, Citrulline 50mg - 2ml):** These amino acids are vital for protein synthesis, muscle repair, and enhancing blood flow, which can improve exercise performance and recovery.



CONTACT US

📞 239-318-0043

✉️ kyleh@thrive-ivinfusions.com

🌐 thrive-ivinfusions.com



Leave Us a
Google Review!



Follow us on
Social Media

TRANSFORM FAT BURNER IV DRIP

- **BPlex (B1, B2, B3, B5, B6 100-2-100-2-2mg/ml - 1ml):** This complex of B vitamins plays a critical role in energy production, red blood cell formation, and maintaining proper nervous system function.
- **1ml IM MIC+B12 (25mg-50mg-50mg-330mcg/ml):** This additional injection includes Methionine, Inositol, Choline, and Vitamin B12, which collectively support fat metabolism, liver function, and energy production. Note: Maximum of 1ml weekly for up to 6 weeks followed by a 3-4 week break unless otherwise advised by a healthcare provider.