

## Comments on the ACPA Discussion Paper, with special reference to goats

Dr Sandra Baxendell, PSM, BVSc (Hons), PhD, MACVSc, GCertAppSC(RurExt), GCertPSectMgt, PGDAppSc, MRurSysMan - Director, Goat Veterinary Consultancies- goatvetoz, [goatvetoz@gmail.com](mailto:goatvetoz@gmail.com), Mobile 0477 813278 [www.goatvetoz.com.au](http://www.goatvetoz.com.au)

Proposals		Comment
<p><b>3 The purpose</b> of the ACPA is to provide a reasonable balance between the welfare needs of animals and the interests of people whose livelihood is dependent on the animals.</p>	Agree	It is recognised that acts of husbandry are necessary for overall health of both animals and people handling animals. However, with advances in technology and pharmacology, pain relief and research into better options should be strongly encouraged.
<p><b>4 Prohibited events</b></p> <ul style="list-style-type: none"> <li>• Cock fighting</li> <li>• Bull fighting</li> <li>• Dog fighting</li> <li>• Coursing</li> <li>• Certain types of hunting</li> <li>• Certain events causing pain (e.g. when people attempt to catch, fight or throw animals)</li> </ul>	Agreed But do not add in goat racing	Goat racing is sometimes criticized but as long as the goats are trained first, this is a good activity for children and young people. This billy goat cart racing is also part of Australia’s culture. Goat racing should be allowed.
<p><b>5 Reporting of animal welfare concerns by veterinary professionals</b></p>	<p>Not agreed and this is also not the policy of the Australian Veterinary Association.</p> <p>Vets would continue to play an educative role with regard to the animal’s health.</p> <p>If mandatory reporting does come in, it should be</p>	<p>Issues include:</p> <ul style="list-style-type: none"> <li>• Clients would be less likely to bring forward animals for treatment if they had concerns about being reported.</li> <li>• Clients may lie about the history of their animals e.g. how much feed was available, if they thought they may be reported and vets may then be directed to other differential diagnoses.</li> <li>• This would be a breach of trust between the vet and the animal owner.</li> <li>• Vets could potentially be put at risk from Domestic Violence partners who could threaten vets or their staff</li> </ul>

	restricted to cruelty rather than neglect.	<ul style="list-style-type: none"> <li>Animal welfare concerns need to be scoped out. There would be an avalanche of reporting if every obese dog's or goat owner was referred to authorities. There are also a lot of grey areas – goats who need surgery e.g. to remove a large udder that was too close to the ground or to remove a tooth just because it is beyond their means.</li> </ul>
<b>6 Regulated surgical procedures:</b> <ul style="list-style-type: none"> <li><b>Cropping dog's ear</b></li> <li><b>Debarking operations</b></li> <li><b>Removal of cats claws</b></li> <li><b>Docking tail of horse or cow or dog</b></li> </ul>	<p>Agree</p> <p>But also add in Applying an elastrator ring to buck kids over 2 months of age. This is an Act of Veterinary Science under the Veterinary Surgeons Act</p>	<p>The castration age limit for goats should stay at &gt;2 months being an act of Vet Surgery. Male dairy goats are fertile at 3 months of age and their testes are large. Actually one of my best clients with stud Saanens confirmed a buck kid impregnated his half-sister at 2 months 3 weeks of age which was when he was moved out to join the bucks in another area despite still receiving milk via a bottle.</p>
<b>7 Possession or use of certain traps and spurs</b>	<p>Agree but more should be added i.e. Opera house traps Steel jawed traps</p>	
<b>8 Dogs</b> <ul style="list-style-type: none"> <li><b>Use of a dog to injure or kill another animal</b></li> <li><b>Closely confining dogs</b></li> <li><b>Restraint of dogs in open vehicle trays and trailers and open windows</b></li> </ul>	<p>Agree re not allowing a dog to kill other animals except for rat seeking terriers e.g. those used by local government.</p> <p>Agree re confinement but need an exception for medical conditions either temporary for recovery after surgery or if a contact or pollen allergy until treated.</p> <p>Also restraint should just be for public roads as dogs on farms will need to be able to jump on and off vehicles while on properties.</p>	<p>The change needs to allow dogs to jump on and off vehicles on farms. .</p>
<b>9 Using animals for scientific purpose</b> <ul style="list-style-type: none"> <li><b>Aligning with Scientific User</b></li> </ul>	<p>Agree but extend to the creation of breeds with defects.</p>	<p>Fainting goat genes have been imported in Australia but only currently used in meat crosses. We should not follow the US examples where the fainting gene is actively selected for and animals</p>



## ITEMS NOT COVERED IN THE DISCUSSION PAPER

- **Disbudding**

Unlike the situation with sheep and cattle, it is not possible to establish a poll goat breed as there is a linked gene to the poll gene for intersex and all pure poll female kids are sterile and can show external sexual characteristics that range from a normal female to a normal male or anything in-between. .Most often they look like a female with an enlarged clitoris. Poll goats should not be kept with horned goats due to bullying and the damage that can be given to the poll goats.

Horned goats often get caught in fences as shown in the photo below and can die of thirst.



Also some goats have such large horns they are a WH&S issue for goat handlers. I lost a tooth when an Angora buck threw back its head while being examined. A child could lose an eye. Dairy goats have never been selected for horn type so horns can be massive as shown in the following photo.





Saanen buck who got his horns caught in a fence and bled profusely.

In the UK, disbudding is only done under anaesthesia by a vet. UK vets anecdotally report that in dairy goat herds where horns are left on, damage to goats requiring veterinary treatment, surgery or destruction happen approximately every 6 months. The British Goat

Veterinary Society produced a video on how to disbud for vets and proposes to run training courses for vets using dead kids.

In the USA anyone can disbud kids where vets feel that nerves to the horn bud area take longer to develop and as long as disbudding happens in the first week then this is acceptable. In the USA a recent survey found that 64% of dairy goat operations, disbudded their kids in the first 3 weeks of life<sup>i</sup> which accounted for 67% of dairy goat breed kids. Meat goats and pet goats were also disbudded but at lower levels<sup>ii</sup>.

In Switzerland, goat farmers were given access to xylazine and ketamine (Schedule 8) drugs to anaesthetize kids prior to disbudding but a review found that farmers were not using these drugs properly and the kids disbudding was either not pain free or there were other problems<sup>iii</sup>. I do not support giving anaesthetic drugs to farmers.

There has been a lot of research looking at alternatives to hot iron disbudding in goats and all have been unsuccessful including:

- Local blocks – as goat kids need 2 blocks per side or a ring block and this risks local anaesthetic toxicity (toxic dose is 1ml for a 2kg kid, the size of some miniature breed kids in the first week after birth). Also research has shown the blocks are more painful than the procedure.<sup>ivvivi</sup>
- Prilocaine cream or EMLA cream as used in children for inserting catheters<sup>viii</sup> and this cream may have even intensified the pain.<sup>ix</sup>
- Clove oil injections.<sup>x</sup>
- Liquid nitrogen<sup>xi</sup>
- Caustic paste was only effective in just over half the kids.<sup>xii</sup>

The latest review in 2021 summarized current knowledge and found “Collectively, we did not find that any of the alternative methods of disbudding provided a feasible option over heat cautery to improve welfare.”<sup>xiii</sup>

While meloxicam has been shown to reduce the throbbing pain that occurs post disbudding, although it has no effect on the pain when the hot irons are applied<sup>xiv</sup>. A survey of British Goat Veterinary Society members found that all vets used an NSAID, generally meloxicam, as well as an anaesthesia for disbudding. NSAIDs such as meloxicam was shown to help with post disbudding pain by several researchers<sup>xxvixvii</sup>.

However there is no meloxicam product registered for use in goats. I use Buccalgesic gel every time I disbud a goat kid. My disbudding procedure is shown on this YouTube video <https://www.youtube.com/watch?v=6mT5tWD9qPE>. I recommend NSAID pain relief for disbudding provided it can be prescribed via telemedicine methods.

- **“Off Label” Prescriptions for Pain Relief products**

Neither TRI-SOLFEN nor any Meloxicam product is registered for goats and many goat owners report they have difficulty finding a veterinarian willing to treat or prescribe for their goats. Also many goats are kept in rangeland areas where veterinary services are a long distance away. For this reason I recommend that pain relief products be able to be prescribed via telemedicine to goat owners for routine husbandry procedures such as disbudding and castration. Covid 19 has shown that human prescriptions can be renewed via telemedicine, so there should be no barriers for veterinary pain relief prescriptions.

- **Access to veterinary medicines for goats**

Access to veterinary medicines is becoming more difficult for veterinarians willing to treat goats or other minor species such as alpacas. This is now a welfare issue. Currently there is only one active that is spray that can be used on dairy goats for lice but no powders nor backline treatments. Goats hate water so a spray is not ideal. Similarly there are only 2 worm drenches (Panacur and Caprimec/Virbamec) that can be used on dairy goats and one is a benzimidazole which has wide-spread resistance. I have clients with milking goats that have worms resistance to both drench families, yet I cannot legally prescribe a sheep drench from a different drench family due to the DO NOT USE statements on the sheep drench labels. Also antibiotics often have a DO NOT USE statement that prohibits use in goats. I recently had milk samples taken from a goat with mastitis and had great difficulty finding antibiotic to use that was needed based on the sensitivity results.

DO NOT USE statements should only be allowed to be placed on labels if there is a genuine danger e.g. monensin in horses or aspirin in cats, rather than just to save the manufacturers money so they don't have to do milk withdrawal period trials. Alternatively a veterinary prescription should be allowed to over-ride a DO NOT USE statement on a label. A good example is levamisole which used to be widely used in milking goats decades ago and for which the USA allows a 3 day milk with-holding period. However now there is no sheep levamisole drench that can be used in milking goats in Australia so one whole drench family cannot be used in combinations or rotations to prevent drench resistance in goats.

## References

- Alvarez, L., R. A. Nava, A. Ramírez, E. Ramírez and J. Gutiérrez (2009). "Physiological and behavioural alterations in disbudded goat kids with and without local anaesthesia." Applied Animal Behaviour Science **117**(3–4): 190-196.
- Hempstead, M. N., J. L. Fowler, J. R. Waas, M. Stewart, M. A. Sutherland, J. L. Fowler and P. J. Plummer (2019). Best-practice recommendations for disbudding dairy goat kids. Fifty-Second Annual Conference. American Association of Bovine Practitioners. St Louis, Missouri. **52**: 192-195
- Hempstead, M. N., T. M. Lindquist, J. K. Shearer, L. C. Shearer, M. A. Sutherland and P. J. Plummer (2019). Acute behavior of dairy goat kids administered lidocaine injections, topical EMLA cream or meloxicam prior to cautery disbudding American Association of Bovine Practitioners. R. A. Smith. St Louis, Missouri: 409.
- Hempstead, M. N., T. M. Lindquist, J. K. Shearer, L. C. Shearer, M. A. Sutherland and P. J. Plummer (2020). "Acute cortisol and behavior of dairy goat kids administered local anesthesia, topical anesthesia or systemic analgesia prior to cautery disbudding." Physiology & Behavior **222**: 112942.
- Hempstead, M. N., J. R. Waas, M. Stewart, S. K. Dowling, V. M. Cave, G. L. Lowe and M. A. Sutherland (2018). "Effect of isoflurane alone or in combination with meloxicam on the behavior and physiology of goat kids following cautery disbudding." Journal of Dairy Science **101**(4): 3193-3204.
- Saini, A. L., D. S. Malik and J. Mohindroo (2013). "EFFECT OF LOCAL ANAESTHETICS AND NON-STEROIDAL ANALGESICS ON MAINTENANCE BEHAVIOUR OF HOT- IRON DISBUDED BEETAL KIDS UNDER STALL- FED CONDITIONS." Indian Journal of Animal Research **47**(5): 407-411.
- Still Brooks, K. M., M. N. Hempstead, J. L. Anderson, R. L. Parsons, M. A. Sutherland, P. J. Plummer and S. T. Millman (2021). "Characterization of Efficacy and Animal Safety across Four Caprine Disbudding Methodologies." Animals (Basel) **11**(2).
- Sutherland, M., F. Huddart and M. Hempstead (2019). Pain relief and novel methods for disbudding calves and goat kids. Science week W. Goodwin. Gold Coast, Queensland, Australia NAZCVS. Animal Welfare and Pharmacology Chapters 7-8.

USDA (2020). Disbudding on U.S. Goat Operations Management Practice Briefs United States Department of Agriculture: 2.

Wagmann, N., C. Spadavecchia, U. Morath-Huss, G. Schüpbach-Regula and P. Zanolari (2018). "Evaluation of anaesthesia and analgesia quality during disbudding of goat kids by certified Swiss farmers." BMC Veterinary Research **14**(1): 220-220.

•

---

<sup>i</sup> USDA (2020). Disbudding on U.S. Goat Operations Management Practice Briefs United States Department of Agriculture: 2.

<sup>ii</sup> Ibid.

<sup>iii</sup> Wagmann, N., C. Spadavecchia, U. Morath-Huss, G. Schüpbach-Regula and P. Zanolari (2018). "Evaluation of anaesthesia and analgesia quality during disbudding of goat kids by certified Swiss farmers." BMC Veterinary Research **14**(1): 220-220.

<sup>iv</sup> Hempstead, M. N., T. M. Lindquist, J. K. Shearer, L. C. Shearer, M. A. Sutherland and P. J. Plummer (2019). Acute behavior of dairy goat kids administered lidocaine injections, topical EMLA cream or meloxicam prior to cautery disbudding American Association of Bovine Practitioners. R. A. Smith. St Louis, Missouri: 409.

<sup>v</sup>

<sup>vi</sup> Saini, A. L., D. S. Malik and J. Mohindroo (2013). "EFFECT OF LOCAL ANAESTHETICS AND NON-STEROIDAL ANALGESICS ON MAINTENANCE BEHAVIOUR OF HOT- IRON DISBUDDED BEETAL KIDS UNDER STALL- FED CONDITIONS." Indian Journal of Animal Research **47**(5): 407-411.

<sup>vii</sup> Alvarez, L., R. A. Nava, A. Ramírez, E. Ramírez and J. Gutiérrez (2009). "Physiological and behavioural alterations in disbudded goat kids with and without local anaesthesia." Applied Animal Behaviour Science **117**(3-4): 190-196.

<sup>viii</sup> Hempstead, M. N., T. M. Lindquist, J. K. Shearer, L. C. Shearer, M. A. Sutherland and P. J. Plummer (2019). Acute behavior of dairy goat kids administered lidocaine injections, topical EMLA cream or meloxicam prior to cautery disbudding American Association of Bovine Practitioners. R. A. Smith. St Louis, Missouri: 409.

<sup>ix</sup> Hempstead, M. N., T. M. Lindquist, J. K. Shearer, L. C. Shearer, M. A. Sutherland and P. J. Plummer (2020). "Acute cortisol and behavior of dairy goat kids administered local anesthesia, topical anesthesia or systemic analgesia prior to cautery disbudding." Physiology & Behavior **222**: 112942.

<sup>x</sup>

<sup>xi</sup> Sutherland, M., F. Huddart and M. Hempstead (2019). Pain relief and novel methods for disbudding calves and goat kids. Science week W. Goodwin. Gold Coast, Queensland, Australia NAZCVS. **Animal Welfare and Pharmacology Chapters** 7-8.

<sup>xii</sup> Still Brooks, K. M., M. N. Hempstead, J. L. Anderson, R. L. Parsons, M. A. Sutherland, P. J. Plummer and S. T. Millman (2021). "Characterization of Efficacy and Animal Safety across Four Caprine Disbudding Methodologies." Animals (Basel) **11**(2).

<sup>xiii</sup> Ibid.

<sup>xiv</sup> Hempstead, M. N., T. M. Lindquist, J. K. Shearer, L. C. Shearer, M. A. Sutherland and P. J. Plummer (2019). Acute behavior of dairy goat kids administered lidocaine injections, topical EMLA cream or meloxicam prior to cautery disbudding American Association of Bovine Practitioners. R. A. Smith. St Louis, Missouri: 409.

<sup>xv</sup> Hempstead, M. N., J. R. Waas, M. Stewart, S. K. Dowling, V. M. Cave, G. L. Lowe and M. A. Sutherland (2018). "Effect of isoflurane alone or in combination with meloxicam on the behavior and physiology of goat kids following cautery disbudding." Journal of Dairy Science **101**(4): 3193-3204.

<sup>xvi</sup> Saini, A. L., D. S. Malik and J. Mohindroo (2013). "EFFECT OF LOCAL ANAESTHETICS AND NON-STEROIDAL ANALGESICS ON MAINTENANCE BEHAVIOUR OF HOT- IRON DISBUDDED BEETAL KIDS UNDER STALL- FED CONDITIONS." Indian Journal of Animal Research **47**(5): 407-411.

<sup>xvii</sup> Hempstead, M. N., J. L. Fowler, J. R. Waas, M. Stewart, M. A. Sutherland, J. L. Fowler and P. J. Plummer (2019). Best-practice recommendations for disbudding dairy goat kids. Fifty-Second Annual Conference. American Association of Bovine Practitioners. St Louis, Missouri. **52**: 192-195