

TOP ICD-10CM TO HCC CODES FOR PRIMARY CARE

| RAF | HCC | DX | MORBID OBESITY |
|-------|-----|-------|---|
| 0.273 | 22 | E6601 | Morbid (severe) obesity due to excess calories |
| 0.273 | 22 | E662 | Morbid (severe) obesity with alveolar hypoventilation |
| 0.273 | 22 | Z6841 | Body mass index (BMI) 40.0-44.9, adult |
| 0.273 | 22 | Z6842 | Body mass index (BMI) 45.0-49.9, adult |
| 0.273 | 22 | Z6843 | Body mass index (BMI) 50-59.9, adult |
| 0.273 | 22 | Z6844 | Body mass index (BMI) 60.0-69.9, adult |
| 0.273 | 22 | Z6845 | Body mass index (BMI) 70 or greater, adult |

| RAF | HCC | DX | HYPERTENSION |
|--------------------------------------|--------|-------|--|
| 0 | NA | I10 | Essential (primary) hypertension, stable |
| HTN AND HEART DISEASE | | | |
| 0.323 | 85 | I110 | Hypertensive heart disease with heart failure <i>Use additional code for heart failure</i> |
| 0 | NA | I119 | Hypertensive heart disease without heart failure |
| HTN AND CKD | | | |
| 0.237 | 136 | I120 | Hypertensive chronic kidney disease with stage 5 chronic kidney disease or end stage renal disease <i>Use additional code for CKD</i> |
| 0 | NA | I129 | Hypertensive chronic kidney disease with stage 1 through stage 4 chronic kidney disease, or unspecified chronic kidney disease <i>Use additional code for CKD</i> |
| HTN AND HEART DISEASE AND CKD | | | |
| 0 | NA | I130 | Hypertensive heart and chronic kidney disease with heart failure and stage 1 through stage 4 chronic kidney disease, or unspecified chronic kidney disease <i>Use additional code for heart failure</i> <i>Use additional code for CKD</i> |
| 0 | NA | I1310 | Hypertensive heart and chronic kidney disease without heart failure, with stage 1 through stage 4 chronic kidney disease, or unspecified chronic kidney disease <i>Use additional code for CKD</i> |
| 0.237 | 136 | I1311 | Hypertensive heart and chronic kidney disease without heart failure, with stage 5 chronic kidney disease, or end stage renal disease <i>Use additional code for CKD</i> |
| 0.237 | 136/85 | I132 | Hypertensive heart and chronic kidney disease with heart failure, with stage 5 chronic kidney disease, or end stage renal disease <i>Use additional code for heart failure</i> <i>Use additional code for CKD</i> |

| | HCC | DX | HEART FAILURE |
|-------|-----|-------|--|
| 0.323 | 85 | I501 | Left ventricular failure |
| 0.323 | 85 | I5020 | Unspecified systolic (congestive) heart failure |
| 0.323 | 85 | I5021 | Acute systolic (congestive) heart failure |
| 0.323 | 85 | I5022 | Chronic systolic (congestive) heart failure |
| 0.323 | 85 | I5023 | Acute on chronic systolic (congestive) heart failure |
| 0.323 | 85 | I5030 | Unspecified diastolic (congestive) heart failure |
| 0.323 | 85 | I5031 | Acute diastolic (congestive) heart failure |
| 0.323 | 85 | I5032 | Chronic diastolic (congestive) heart failure |
| 0.323 | 85 | I5033 | Acute on chronic diastolic (congestive) heart failure |
| 0.323 | 85 | I5041 | Acute combined systolic (congestive) and diastolic (congestive) heart failure |
| 0.323 | 85 | I5042 | Chronic combined systolic (congestive) and diastolic (congestive) heart failure |
| 0.323 | 85 | I5043 | Acute on chronic combined systolic (congestive) and diastolic (congestive) heart failure |
| 0.323 | 85 | I509 | Heart failure, unspecified |

| | HCC | DX | CHRONIC KIDNEY DISEASE |
|-------|-----|------|------------------------|
| 0 | NA | N181 | Stage 1, CKD |
| 0 | NA | N182 | Stage 2, CKD |
| 0 | NA | N183 | Stage 3, CKD |
| 0.237 | 137 | N184 | Stage 4, CKD |
| 0.237 | 136 | N185 | Stage 5, CKD |
| 0.237 | 136 | N186 | Stage 6, CKD |

| RAF | HCC | DX | DIALYSIS |
|-------|-----|-------|---|
| 0.442 | 134 | Z992 | Dependence on renal dialysis |
| 0.442 | 134 | Z9115 | Patient's noncompliance with renal dialysis |

| RAF | HCC | DX | TYPE 2 DIABETES MELLITUS |
|---|-----|--------|---|
| 0.104 | 19 | E119 | Type 2 diabetes mellitus without complications |
| TYPE 2 DM WITH KIDNEY COMPLICATIONS | | | |
| 0.318 | 18 | E1121 | with diabetic nephropathy |
| 0.318 | 18 | E1122 | with diabetic chronic kidney disease <i>Use additional code for CKD</i> |
| 0.318 | 18 | E1129 | with other diabetic kidney complication |
| TYPE 2 DM WITH NERUOLGOIC COMPLICATIONS | | | |
| 0.318 | 18 | E1140 | with diabetic neuropathy, unspecified |
| 0.318 | 18 | E1141 | with diabetic mononeuropathy |
| 0.318 | 18 | E1142 | with diabetic polyneuropathy |
| 0.318 | 18 | E1143 | with diabetic autonomic (poly)neuropathy |
| 0.318 | 18 | E1144 | with diabetic amyotrophy |
| 0.318 | 18 | E1149 | with other diabetic neurological complication |
| TYPE 2 DM WITH CIRCULATORY COMPLICATIONS | | | |
| 0.318 | 18 | E1151 | with diabetic peripheral angiopathy without gangrene |
| 0.318 | 18 | E1152 | with diabetic peripheral angiopathy with gangrene |
| 0.318 | 18 | E1159 | with other circulatory complications |
| TYPE 2 DM WITH OTHER COMPLICATIONS | | | |
| 0.318 | 18 | E11610 | with diabetic neuropathic arthropathy |
| 0.318 | 18 | E11620 | with diabetic dermatitis |
| 0.318 | 18 | E11621 | with foot ulcer <i>Use additional code for foot ulcer - L97 series -</i> |
| 0.318 | 18 | E11622 | with other skin ulcer |
| 0.318 | 18 | E11628 | with other skin complications |
| 0.318 | 18 | E11630 | with periodontal disease |
| 0.318 | 18 | E11638 | with other oral complications |
| 0.318 | 18 | E11649 | with hypoglycemia without coma |
| 0.318 | 18 | E1165 | with hyperglycemia |
| 0.318 | 18 | E1169 | with other specified complication |
| 0.104 | 19 | Z794 | Long term (current) use of insulin |

| RAF | HCC | DX | COPD |
|-------|-----|------|---|
| 0.328 | 111 | j440 | COPD with acute lower respiratory infection |
| 0.328 | 111 | j441 | COPD with acute exacerbation |
| 0.328 | 111 | j449 | COPD, unspecified |

| RAF | HCC | DX | CARDIOVASCULAR |
|-------|-----|-------|-------------------------------------|
| 0.268 | 96 | I470 | Re-entry ventricular arrhythmia |
| 0.268 | 96 | I471 | Supraventricular tachycardia |
| 0.268 | 96 | I472 | Ventricular tachycardia |
| 0.268 | 96 | I479 | Paroxysmal tachycardia, unspecified |
| 0.268 | 96 | I480 | Paroxysmal atrial fibrillation |
| 0.268 | 96 | I481 | Persistent atrial fibrillation |
| 0.268 | 96 | I482 | Chronic atrial fibrillation |
| 0.268 | 96 | I483 | Typical atrial flutter |
| 0.268 | 96 | I484 | Atypical atrial flutter |
| 0.268 | 96 | I4891 | Unspecified atrial fibrillation |
| 0.268 | 96 | I4892 | Unspecified atrial flutter |
| 0.268 | 96 | I492 | Junctional premature depolarization |
| 0.268 | 96 | I495 | Sick sinus syndrome |

| RAF | HCC | DX | PSYCHIATRIC |
|-------|-----|-------|--|
| 0.395 | 58 | F22 | Delusional disorders |
| 0.395 | 58 | F24 | Shared psychotic disorder |
| 0.395 | 58 | F3010 | Manic episode without psychotic symptoms, unspecified |
| 0.395 | 58 | F3011 | Manic episode without psychotic symptoms, mild |
| 0.395 | 58 | F3012 | Manic episode without psychotic symptoms, moderate |
| 0.395 | 58 | F3013 | Manic episode, severe, without psychotic symptoms |
| 0.395 | 58 | F302 | Manic episode, severe with psychotic symptoms |
| 0.395 | 58 | F303 | Manic episode in partial remission |
| 0.395 | 58 | F304 | Manic episode in full remission |
| 0.395 | 58 | F308 | Other manic episodes |
| 0.395 | 58 | F309 | Manic episode, unspecified |
| 0.395 | 58 | F319 | Bipolar disorder, unspecified |
| 0.395 | 58 | F320 | Major depressive disorder, single episode, mild |
| 0.395 | 58 | F321 | Major depressive disorder, single episode, moderate |
| 0.395 | 58 | F322 | Major depressive disorder, single episode, severe without psychotic features |
| 0.395 | 58 | F323 | Major depressive disorder, single episode, severe with psychotic features |
| 0.395 | 58 | F324 | Major depressive disorder, single episode, in partial remission |
| 0.395 | 58 | F325 | Major depressive disorder, single episode, in full remission |
| 0.395 | 58 | F330 | Major depressive disorder, recurrent, mild |
| 0.395 | 58 | F331 | Major depressive disorder, recurrent, moderate |
| 0.395 | 58 | F332 | Major depressive disorder, recurrent severe without psychotic features |
| 0.395 | 58 | F333 | Major depressive disorder, recurrent, severe with psychotic symptoms |
| 0.395 | 58 | F3340 | Major depressive disorder, recurrent, in remission, unspecified |
| 0.395 | 58 | F3341 | Major depressive disorder, recurrent, in partial remission |
| 0.395 | 58 | F3342 | Major depressive disorder, recurrent, in full remission |
| 0.395 | 58 | F339 | Major depressive disorder, recurrent, unspecified |
| 0.395 | 58 | F348 | Other persistent mood [affective] disorders |
| 0.395 | 58 | F349 | Persistent mood [affective] disorder, unspecified |
| 0.395 | 58 | F39 | Unspecified mood [affective] disorder |

| RAF | HCC | DX | DRUG DEPENDENCE |
|-------|-----|-------|------------------------------------|
| 0.383 | 55 | F1020 | Alcohol dependence, uncomplicated |
| 0.383 | 55 | F1021 | Alcohol dependence, in remission |
| 0.383 | 55 | F1220 | Cannabis dependence, uncomplicated |
| 0.383 | 55 | F1120 | Opioid dependence, uncomplicated |

| RAF | HCC | DX | AQUIRED ABSENSE |
|-------|-----|--------|--|
| 0.588 | 189 | Z89411 | Acquired absence of right great toe |
| 0.588 | 189 | Z89412 | Acquired absence of left great toe |
| 0.588 | 189 | Z89421 | Acquired absence of other right toe(s) |
| 0.588 | 189 | Z89422 | Acquired absence of other left toe(s) |
| 0.588 | 189 | Z89431 | Acquired absence of right foot |
| 0.588 | 189 | Z89432 | Acquired absence of left foot |
| 0.588 | 189 | Z89441 | Acquired absence of right ankle |
| 0.588 | 189 | Z89442 | Acquired absence of left ankle |
| 0.588 | 189 | Z89511 | Acquired absence of right leg below knee |
| 0.588 | 189 | Z89512 | Acquired absence of left leg below knee |
| 0.588 | 189 | Z89611 | Acquired absence of right leg above knee |
| 0.588 | 189 | Z89612 | Acquired absence of left leg above knee |