

Current Biology

Expectations and Outcomes of Reserve Network Performance following Re-zoning of the Great Barrier Reef Marine Park

Highlights

- Expansion of a GBR reserve network led to substantial increases in coral trout stocks
- A large tropical cyclone impacted the reserve network, causing widespread damage
- Impacts on coral trout biomass were smaller inside NTMRs compared to fished reefs
- NTMRs may provide greater potential for replenishment after disturbances

Authors

Michael J. Emslie, Murray Logan, ..., Garry R. Russ, Hugh P.A. Sweatman

Correspondence

m.emslie@aims.gov.au

In Brief

No-take marine reserves (NTMRs) are widely advocated for conserving exploited fish stocks and biodiversity. Emslie et al. show that expanding NTMR networks had clear benefits for fishery target, but not non-target, species. A cyclone caused widespread degradation, but target species biomass was retained within NTMRs, with greater recovery potential.



Expectations and Outcomes of Reserve Network Performance following Re-zoning of the Great Barrier Reef Marine Park

Michael J. Emslie,^{1,*} Murray Logan,¹ David H. Williamson,^{2,3} Anthony M. Ayling,⁴ M. Aaron MacNeil,¹ Daniela Ceccarelli,² Alistair J. Cheal,¹ Richard D. Evans,^{5,6} Kerryn A. Johns,¹ Michelle J. Jonker,¹ Ian R. Miller,¹ Kate Osborne,¹ Garry R. Russ,^{2,3} and Hugh P.A. Sweatman¹

¹Australian Institute of Marine Science, PMB No. 3, Townsville MC, Townsville, QLD 4810, Australia

²ARC Centre of Excellence for Coral Reef Studies, James Cook University, Townsville, QLD 4811, Australia

³College of Marine and Environmental Sciences, James Cook University, Townsville, QLD 4811, Australia

⁴Sea Research, 20 Rattray Avenue, Hideaway Bay, QLD 4800, Australia

⁵Department of Parks and Wildlife, 17 Dick Perry Avenue, Kensington, Perth, WA 6151, Australia

⁶Oceans Institute, University of Western Australia, Crawley, WA 6009, Australia

*Correspondence: m.emslie@aims.gov.au

<http://dx.doi.org/10.1016/j.cub.2015.01.073>

SUMMARY

Networks of no-take marine reserves (NTMRs) are widely advocated for preserving exploited fish stocks and for conserving biodiversity. We used underwater visual surveys of coral reef fish and benthic communities to quantify the short- to medium-term (5 to 30 years) ecological effects of the establishment of NTMRs within the Great Barrier Reef Marine Park (GBRMP). The density, mean length, and biomass of principal fishery species, coral trout (*Plectropomus* spp., *Variola* spp.), were consistently greater in NTMRs than on fished reefs over both the short and medium term. However, there were no clear or consistent differences in the structure of fish or benthic assemblages, non-target fish density, fish species richness, or coral cover between NTMR and fished reefs. There was no indication that the displacement and concentration of fishing effort reduced coral trout populations on fished reefs. A severe tropical cyclone impacted many survey reefs during the study, causing similar declines in coral cover and fish density on both NTMR and fished reefs. However, coral trout biomass declined only on fished reefs after the cyclone. The GBRMP is performing as expected in terms of the protection of fished stocks and biodiversity for a developed country in which fishing is not excessive and targets a narrow range of species. NTMRs cannot protect coral reefs directly from acute regional-scale disturbance but, after a strong tropical cyclone, impacted NTMR reefs supported higher biomass of key fishery-targeted species and so should provide valuable sources of larvae to enhance population recovery and long-term persistence.

INTRODUCTION

Coral reefs are under increasing pressure, leading to debate about strategies to conserve biodiversity, enhance resilience, and maintain ecosystem processes in these habitats [1–4]. Fully protected no-take marine reserves (hereafter, NTMRs), defined as “areas of the ocean completely protected from all extractive and destructive activities” [5], are a widely advocated tool for conservation and management of marine systems [6–9]. Historically, NTMRs were conceived as a fisheries management tool to protect exploited stocks, prevent overfishing, and mitigate habitat destruction, allowing the recovery of exploited populations once fishing pressure and associated habitat destruction cease. However, in recent decades, their use has expanded to include protection of biodiversity and ecosystem processes. Whether or not NTMRs can perform these roles depends on the nature of the threats to biodiversity and the efficacy of NTMRs in countering these threats. Since NTMRs generally only eliminate extractive fishing activities, their effectiveness can vary according to size, location, and enforcement, as well as the selectivity, catch, and effort of the fishery. Hence, NTMRs would only be expected to have substantial effects on fished stocks and biodiversity under certain conditions.

There is now abundant evidence that adequately protected NTMRs are effective as fishery reserves, increasing the abundance, size, and biomass of species targeted by fisheries in both tropical and temperate systems [8, 10–18]. Importantly, NTMRs may also contribute to maintaining populations in adjacent fished areas through larval recruitment subsidies and spillover of adult fish [19–21]. While evidence suggests that NTMRs are performing successfully as fishery reserves, key questions still remain: how much area needs to be preserved to sustain fisheries at different levels of fishing pressure, and how does the spatial redistribution of fishing effort after the establishment of NTMRs affect exploited fish populations in areas that remain open to fishing?

Beyond effects on fisheries, can NTMRs effectively conserve or restore natural states of biodiversity and enhance resilience,

particularly in coral reef ecosystems? The answer to this question depends largely on the socio-economic setting of the region: specifically, the distribution and intensity of fishing pressure and the diversity of species exploited by the fishery [5, 22]. Where fisheries exploit a broad range of species that perform many ecological functions (e.g., the Caribbean, the Pacific, and Southeast Asia), and in locations where destructive fishing methods are employed (e.g., dynamite, cyanide, or muro-ami), NTMRs may be expected to significantly enhance biodiversity and maintain habitat condition. In contrast, in many developed countries, like Australia and the US, where only a limited range of high-trophic-level predatory species are targeted by fisheries and destructive fishing techniques are prohibited, enhancement of biodiversity within NTMRs may be limited and difficult to detect. For NTMRs to influence the abundance of non-targeted fish species, hard coral cover, the structure of reef fish and benthic assemblages, and biodiversity, indirect ecological processes must occur (see [4, 14, 23, 24]). Such indirect processes include trophic cascades, where targeted species exert top-down control of species at lower trophic levels, but there is little evidence of strong top-down control on species-rich coral reefs [14, 23, 25, 26]. NTMRs may protect habitat characteristics such as coral cover and benthic community composition where destructive fishing practices are used (e.g., dynamite fishing) [27], but there is little evidence that they can contribute to maintaining or enhancing coral cover in areas where less damaging fishing methods are used (e.g., spearfishing or hook and line) [14, 23, 24, 28].

While the primary goals of NTMRs are to act as fishery reserves and protect biodiversity, many of the stressors degrading coral reefs—pollution, sedimentation, coastal development, and the cumulative, escalating effects of climate change—are not related to fishing. Climatic disturbance events such as cyclones, flood plumes, and coral bleaching can severely degrade coral reefs and erode the accrued benefits of reserves at relatively local scales [29–31]. The frequency of extreme climatic disturbance events is predicted to increase in the coming decades, and it is important to consider the role that reserve networks could play in enhancing resilience and population persistence at regional and ecosystem scales [32]. While it seems obvious that reserves can do little to mitigate the acute impacts of severe climatic disturbances at local scales [33], this assumption has rarely been directly tested in a large, well-connected NTMR network [32].

The Great Barrier Reef Marine Park (GBRMP) includes a large-scale network of NTMRs that extends over 2,000 km along the northeast coast of Australia. The GBRMP has a zoning history spanning 30 years (Supplemental Experimental Procedures), and in 2004 a new zoning plan increased the total no-take reserve area from approximately 5% to 33% of the marine park. The main fishery operating within the GBRMP is a hook-and-line fishery primarily targeting coral trout (*Plectropomus* spp., *Variola* spp., family Serranidae) [34]. A limited range of other reef fishes (principally from the families Lethrinidae and Lutjanidae) are not directly targeted, but individuals that are above the minimum legal length are often retained when captured [34]; here these are termed “secondary targets” (Table S1). The GBRMP has a small and localized coastal population with moderate coastal development and has recently been exposed to a succession of severe acute disturbance events,

after which the cover of habitat-forming hard corals has declined significantly on many reefs [35–37]. The majority of the recent coral loss has occurred since 2006, after multiple storms damaged large areas of the central and southern GBRMP. Most notable was severe Tropical Cyclone Hamish in 2009 (Figure 1), which caused extensive physical damage to offshore reefs, widespread freshwater inundation of inshore reefs, and localized bleaching events [38, 39].

The GBRMP is a benchmark for the implementation of networks of reserves, particularly for coral reefs, and has inspired comparable large-scale action around the world (e.g., the US west coast, Hawaii, Mediterranean, and Coral Triangle Initiative). Because of its global importance as an example of the type of action that many believe is required to sustain coastal ecosystem services, there is general interest in how the GBRMP performs. However, any assessment of the performance of the GBRMP, or any other reserve network, must consider the disturbance history and socio-geographical settings of the region. NTMR networks in more degraded and heavily fished systems, such as the Caribbean or Southeast Asia, would be expected to perform quite differently from those in less degraded systems with lower fishing pressure.

Here we use long-term datasets (2004–2012 and 1983–2012; Table 1) from reefs spread over ~150,000 km² of the GBRMP (Figure 1) first to assess several key ecological measures of NTMR performance after a major re-zoning of the GBRMP in 2004 and second to determine the degree to which accrued NTMR benefits were affected by a tropical cyclone. Specifically, we asked three key questions:

1. Fishery effects—were the density, length, and biomass of key targeted reef fish species higher on reefs within NTMRs than on reefs that were open to fishing?
2. Biodiversity effects—did the density of non-target reef fish species, species richness of reef fishes, hard coral cover, and assemblage structure of fish and benthic communities differ between reefs in NTMRs and reefs that were open to fishing?
3. Disturbance effects—did a severe tropical cyclone affect any accrued benefits of NTMRs?

RESULTS

Fishery Effects

GBR-wide Effects of Reserve Status

Despite variability at finer temporal (among years) and spatial (among offshore sectors and inshore island groups) scales (Table S2 and Figure S1), the re-zoning of the GBRMP in 2004 resulted in clear GBRMP-wide increases in the density, length, and biomass of the primary target of the hook-and-line fishery, coral trout, on NTMR reefs relative to fished reefs (Figure 2). In inshore and offshore NTMRs, 53% and 67% of coral trout, respectively, were larger than the minimum legal size (38 cm total length [TL]), compared with 26% inshore and 56% offshore on adjacent fished reefs (Figure S2). On average, coral trout were 12% and 7% larger on inshore and offshore NTMR reefs, respectively, compared with reefs that were open to fishing (Figure 2). The differences in coral trout density and mean size translated into an




Figure 1. Map of the Study Locations

Inshore sites are located on fringing reefs surrounding high continental islands within 30 km of the coast, and offshore sites are on platform reefs >30 km from the coast. The track of Tropical Cyclone Hamish is the red line, with destructive (orange) and very destructive (red) wind fields (from the Australian Bureau of Meteorology). The gray horizontal dotted line delineates control from impact reefs used in the BACI analysis of effects of the cyclone. Reefs of the Capricorn-Bunker sector lay on the edge of the destructive wind zone but are considered to be impact reefs because they were fully exposed to the storm swell generated by the cyclone at its peak.

89% higher biomass in inshore NTMRs and an 82% higher biomass in offshore NTMRs (Figure 2). Benefits to secondary target fishes were less clear. Although secondary target fishes on offshore NTMR reefs were 1% larger and biomass was 30% greater compared with fished reefs, no such differences were evident on inshore reefs (Figure 2).

Historical Trends

A full before-after-control-impact (BACI) analysis of the effects of the 2004 re-zoning of the GBRMP was not possible due to the lack of data from offshore reefs before 2004. However, by combining three datasets spanning 1983–2012 (Supplemental Experimental Procedures), we were able to put the post-2004 changes in coral trout populations into an historical context. Before the widespread establishment of the GBRMP in the 1980s, GBR-wide coral trout biomass was $\sim 5 \text{ kg } 1,000 \text{ m}^{-2}$, subsequently declining to 1–2 $\text{kg } 1,000 \text{ m}^{-2}$ by 1996. GBR-wide

biomass on NTMR reefs had increased to $\sim 5 \text{ kg } 1000 \text{ m}^{-2}$ by the time the GBRMP was re-zoned in 2004, before again increasing rapidly to the highest levels recorded since the 1980s in 2008. GBR-wide biomass subsequently declined to $\sim 5 \text{ kg } 1,000 \text{ m}^{-2}$ coincident with the occurrence of Tropical Cyclone Hamish in 2009, but there was evidence for some recovery after the cyclone (Figure 3). Significantly, on reefs that were open to fishing, GBR-wide biomass remained stable or increased after the 2004 re-zoning, except after Tropical Cyclone Hamish, when changes in numbers were similar to those on NTMR reefs (Figure 3). Note that the catch and effort of the GBR hook-and-line fishery increased from the early 1990s until 2002, declined from 2002 until 2005, then remained stable until 2012 (Figure S3).

The overall GBR-wide time-averaged coral trout biomass was ~ 2.5 times higher on NTMR reefs than on those open to fishing (Figure 3). Offshore, this pattern was true for all sectors, and

Table 1. Methodological Details for Each Monitoring Program

Source	Spatial Coverage	Number of Reefs	Sites per Reef	Transects per Site	Data	Methods	Frequency and Use in This Study
Ayling (1983–1986)	six sectors; 11° latitude	50	1–2 (140 in total)	10 (1,400 in total)	coral trout abundance and length	UVC on 50 × 10 m belt transects	one off; historical modeling
James Cook University monitoring the effects of rezoning on inshore reefs (2004–present)	three island groups; 4° latitude	6	1–3 (34 per year in total)	5 (170 per year in total)	fish: abundance and lengths of 190 species; benthos: percent cover of hard and soft coral and algae	UVC on 50 × 6 m belt transects, 50 m line intercept transects, 50 points per transect	annually; post-re-zoning analysis and historical modeling
AIMS LTMP (1995–present)	six sectors; 11° latitude	47	3 (141 per year in total)	5 (705 per year in total)	fish: abundance of 215 species and lengths for target species; benthos: percent cover of hard and soft coral and algae	UVC belt transects; 1995–present: 50 × 5 m, 50 × 1 m damselfish; 1995–2005: video transect 40 frames, photo transects 40 frames	annually from 1995–2005, then biennially; historical modeling
AIMS monitoring the effects of rezoning (2006–present)	five regions; 9° latitude	56 (28 reserve and non-reserve pairs)	3 (168 per year in total)	5 (840 per year in total)	as for LTMP	as for LTMP	biennially since 2006; post-re-zoning analysis and historical modeling

LTMP, Long-term Monitoring Program; UVC, underwater visual census.


although the magnitude of the difference varied, the ratio of biomass in offshore NTMRs to that on fished reefs was always greater than 1.5 (Figure 3). On inshore reefs in the 1980s, coral trout biomass was generally lower than recorded offshore at that time. After 15 to 20 years of protection, biomass was greater than 1980s levels on NTMR reefs but remained similar to 1980s levels on reefs that were open to fishing (Figure 3).

Biodiversity Effects

There were few differences in the density of most non-target fish species, the percent cover of benthic organisms, and the structure of assemblages of fishes and benthic organisms between NTMR and fished reefs. On inshore reefs, benthic foragers were 21% more abundant on reefs that were open to fishing than on NTMR reefs (Figure 2). On offshore reefs, detritivores, omnivorous damselfishes, and benthic foragers were all between 13% and 35% more abundant in NTMRs compared with fished reefs (Figure 2). Species richness of reef fishes was 8% greater in offshore NTMRs than on fished reefs (Figure 2), but the species that contributed most to this difference were rare (e.g., *Chaetodon bennetti*, *Chaetodon meyeri*, *Lethrinus ornatus*, and *Lethrinus rubriopercularis*) and occurred in very low densities. There were no differences in cover of hard coral, soft coral, or algae (Figure 2) between NTMR and fished reefs. There was very little evidence that NTMR status affected the overall structure of the assemblages of fishes or benthic organisms on either inshore or offshore reefs (Figure 4). NTMR zoning status accounted for <1% of the total variation in reef fish assemblage structure, whereas differences among sectors or island groups accounted for 33%–50% of the variation.

Disturbance Effects on Offshore Reefs

In March 2009, Tropical Cyclone Hamish tracked along much of the southern GBR (Figure 1). In its wake, there were significant declines in hard coral cover and in the density of numerous fish groups, with increases in total algal cover (turf, coralline, and macro-algae) on offshore NTMR and fished reefs in the impacted region (Figure 5). There were no substantial changes in any of these variables over the same period on more northern “control” survey reefs that were not affected by the cyclone (Figure 5). While the density of coral trout declined on both NTMR and fished reefs in the impacted region, coral trout biomass only declined on fished reefs, with no concomitant change on NTMR reefs (Figure 5). At the same time, there was little or no change in the density and biomass of coral trout on reefs in the control region that were not affected by the cyclone (Figure 5). There were no significant temporal changes in density or biomass of secondary target species or in total species richness of reef fish on either NTMR or fished reefs in either the impact or control regions (Figure 5). The density of benthic foragers and obligate corallivores declined on both NTMRs and fished reefs in the impact region, but not in the control region. The density of omnivorous damselfishes and territorial farming damselfishes also declined after the cyclone, but only on reefs in the impact region that were open to fishing (Figure 5). The density of herbivorous scrapers increased in NTMRs only, whereas planktivore density increased on both NTMR and fished reefs in the impacted region with no equivalent changes on reefs in the control region (Figure 5).



DISCUSSION

This study clearly demonstrates that the GBRMP is performing as expected, given its northeastern Australian setting with relatively low fishing pressure and a fishery that targets a limited number of top-level predators. NTMRs established during the 2004 re-zoning of the GBRMP have yielded significant benefits for populations of targeted coral reef fishes on both inshore and offshore reefs within the first decade of protection. Substantial increases in the mean density, body size, and biomass of exploited species were consistently recorded on NTMR reefs, whereas there were few discernible changes on reefs that remained open to fishing. Importantly, there was no indication that the density, size, or biomass of targeted fish species was reduced on fished reefs as might occur from the displacement and concentration of fishing effort after the establishment of the NTMR network. Additionally, there were no differences in crude measures of biodiversity and, despite the major impacts of a tropical cyclone, the biomass of coral trout remained relatively stable on NTMR reefs but declined on fished reefs.

The absence of data on offshore reefs from before the new zoning plan came into effect made it difficult to attribute post-2004 increases in coral trout biomass unequivocally to NTMR protection. To address this, and to place the monitoring data

Figure 2. GBR-wide Effects of NTMRs on Fishery Target Species, Non-target Fish Groups, and Benthic Organisms

Effect sizes were averaged over all surveys since the re-zoning in 2004. Data are modeled median differences between NTMRs and fished reefs with associated 95% uncertainty intervals (UIs) for inshore (open symbols) and offshore (closed symbols) reefs. Data were modeled using a Bayesian hierarchical linear mixed model, and differences are expressed as a percentage of the value for fished reefs. A positive effect indicates higher values in NTMRs, and statistical significance is inferred where UIs do not intersect zero. *, no results are presented for excavators and detritivores on inshore reefs as the models did not converge. **, no results are presented for inshore algae as only macro algal cover was recorded. See also Figures S1 and S2.

from 2006 to 2012 into historical context, we modeled coral trout biomass from data sets spanning thirty years collected on fished and NTMR reefs. Biomass of coral trout increased over both short (2–3 years after the 2004 re-zoning) and medium (since 1996) time scales. Although such results are not without precedent [12, 14, 40], increases in coral trout biomass on NTMR reefs occurred more rapidly than in the majority of previous studies. Such short-term increases may reflect redistribution of biomass to the reserves after re-zoning. It is also possible that the study reefs were supporting high densities of sub-legal size (<38 cm TL) coral trout prior to the establishment of reserves in 2004. Given that fish body weight generally increases exponentially with increasing length [41], the rapid biomass increases on NTMR reefs may have also been at least partly due to higher numbers of fish surviving beyond 38 cm. Alternatively, the increases in coral trout biomass may simply have been a function of increasing reserve area. It is clearly not possible to apportion the contribution of these potential mechanisms to the rapid gains in coral trout biomass observed on NTMR reefs with certainty. Intuitively, however, the increase in NTMR reef area from pre-2004 to post-2004 coupled with improved surveillance and enforcement of GBRMP zoning regulations and the implementation of a range of direct fishery management actions in 2004 [42] are all likely to have contributed.

Although the benefits of NTMRs for exploited species were expected, an unanticipated result was that the reduction in the reef area available to fishers after the 2004 re-zoning did not reduce densities of coral trout on reefs that remained open to fishing. After an initial decline from 1980s levels, populations of coral trout on fished reefs remained stable or increased slightly from 1996 until 2012, suggesting that the catch rates of the GBR Line Fishery have been sustainable since the creation of the GBRMP. The increased area of NTMRs inside the GBRMP after the 2004




Figure 3. Long-Term Modelling of Coral Trout Biomass

Historical estimates of coral trout biomass in inshore and offshore NTMRs (filled symbols) and on reefs that were open to fishing (open symbols). Triangles indicate data from the dedicated post-re-zoning surveys (2006–2012), and circular symbols indicate AIMS Long-term Monitoring Program data (1995–2011). Sector labels are the same as in Figure 1. Data points for 2004 in the two inshore sectors (PA and WH) show coral trout biomass immediately prior to the re-zoning in 2004 but are coded according to the zones in place after 2004. Trends were modeled using a Bayesian hierarchical linear mixed model with a zero-inflated negative binomial distribution. The dark line and shaded band are the modeled medians \pm 95% UIs for coral trout biomass in NTMRs, and the light line and shaded band give the same information for reefs open to fishing. Black square symbols indicate median coral trout biomass in the 1980s (\pm 95% UIs) before the implementation of zoning on the GBR. The effect size plot (bottom right-hand panel) shows the modeled median ratio and associated 95% UIs of coral trout biomass in NTMRs compared with fished reefs (1980s–2012) on inshore (IN) island groups and offshore sectors. The dashed vertical line indicates equal biomass of coral trout on NTMR and fished reefs. A positive effect indicates higher values in NTMRs and statistical significance is inferred where UIs do not intersect zero. See also Figure S3.

greater and fishing effort is more broadly distributed, so the less accessible offshore populations remained relatively lightly exploited through the 1980s [45] and supported coral trout biomass similar to levels in NTMRs today. Fishing pressure on offshore reefs increased through the 1990s [46, 47] as both commercial and recreational fishing expanded [48]. The limited historical data suggest that

re-zoning may have theoretically resulted in a “squeeze effect” [43, 44], with a relocation and concentration of fishing effort on the remaining fished reefs and concomitant reductions in the abundance and biomass of target species. The lack of evidence for such an effect in the present study suggests that fishery management actions such as the GBRMP Structural Adjustment Package (GBRMPSAP), introduced shortly after the 2004 re-zoning, were effective in sustaining stock levels on fished reefs. The GBRMPSAP included a license buyout program, which successfully reduced the catch and effort of the coral reef line fishery from an all-time high in 2002 to lower but stable levels from 2005 onward.

Analysis of historical coral trout biomass suggests that populations on inshore reefs had been depleted by the 1980s, before establishment of the GBRMP. The limited area of inshore fringing reef habitat is readily accessible from the mainland, so fishing effort is highly concentrated, increasing the potential for population depletion. In contrast, the area of offshore reefs is much

the number of participants in the commercial line fishery declined from 279 in 1980/1981 to 176 in 1990, but there was an increase in catch from 201 tons 1980/1981 to 1,490 tons by 1990 [46]. This increase in the commercial catch seems the most likely explanation for the reduction in coral trout biomass on offshore reefs that we observed between the 1980s and 1995.

One of the key objectives of the 2004 re-zoning of the GBRMP was to preserve biodiversity, yet we found no large differences in coarse measures of biodiversity between fished and NTMR reefs in the present study. There was no difference in reef fish species richness between inshore NTMRs and fished reefs, and species richness was only marginally higher (8%) on offshore NTMRs than on fished reefs. This result is not surprising, as the main function of NTMRs is to reduce fishing pressure. The Reef Line Fishery operating within the GBRMP targets a narrow suite of predatory fishes and thus cannot be considered a major threat to biodiversity. In comparison, we would expect NTMRs to influence biodiversity directly in other regions of the world where




Figure 4. The Effect of NTMR Zoning on Fish and Benthic Assemblage Structure

Panels are a visualization of the structure reef fish or benthic assemblages on inshore and offshore reefs (2006–2012), based on a redundancy analysis (RDA) accounting for differences due to latitudinal sector (offshore reefs), island group (inshore reefs), and NTMR status (closed symbols, NTMR; open symbols, open to fishing). All data were standardized (row centered) prior to analysis; reef fish community data were fourth root transformed and benthic data were square root transformed to reduce the effect of highly abundant taxa.

coral reef habitats following the cyclone [49], which probably reduced prey abundance [30]. Loss of shelter may reduce the effectiveness of ambush predators such as coral trout [50], forcing them to relocate to leeward or deep water locations around the reef that were less damaged by the cyclone and still retained high levels of habitat complexity.

While coral trout density declined equally on both NTMR and fished reefs after Tropical Cyclone Hamish, NTMRs surprisingly retained significantly greater coral trout biomass than reefs that were open to fishing. Larger fishes may be better able to withstand turbulence during cyclones, be less dependent on remain-

ing reef structure after disturbances, or have a greater capacity to move to refuge areas (e.g., deeper reef habitats) and return to shallow reef areas when conditions have settled. In any case, this finding has important implications because the retention of coral trout biomass in NTMRs after Tropical Cyclone Hamish may speed recovery of depleted populations both inside and outside NTMRs via larval dispersal [20, 32].

Some non-target species (e. g. planktivores and scrapers) were more abundant after Tropical Cyclone Hamish, whereas others (e. g. obligate corallivores and benthic foragers) declined in abundance. Such changes can be explained by increases to algal cover and reductions to hard coral cover [31, 51, 52]; however, there was no indication that the responses of these fishes, or of algae and hard coral, differed between NTMRs and reefs open to fishing. Like other large-scale disturbances such as coral bleaching events [33] and flood plumes [30], large storms appear to swamp any differences in resistance between NTMRs and fished reefs. Marine reserves provide no direct protection from storms, flood plumes, or temperature anomalies at local scales, but the establishment of a large network of NTMRs inside the GBRMP spaced over 1,000s of kilometers ensured that there were protected reefs that were unaffected by Tropical Cyclone Hamish; these remained as potential sources for re-seeding damaged reefs, thereby maintaining biodiversity and the persistence of coral trout populations at regional and ecosystem scales [30].

fishers target a wide range of species that perform many ecological functions, often using methods that destroy coral habitat. In contrast to recent work in the Caribbean [23], our results suggest that the current levels of fishing exert little top-down control on the abundant and speciose reef fish assemblages in the GBRMP. The structure of reef fish and benthic assemblages appears to be largely driven by bottom-up processes, such as exposure, variability in larval supply, and the effects of disturbances such as large-scale storms.

Tropical Cyclone Hamish caused widespread declines in coral cover on both NTMR and fished reefs across a broad swathe of the southern GBRMP in 2009. Such broad-scale damage to habitat-forming hard corals commonly has direct effects on reef-associated species such as fishes [31], and in the months following the cyclone, commercial fishers reported that coral trout catch rates had declined [34]. Our analysis of the impacts of Tropical Cyclone Hamish indicated that NTMR and fished reefs fared equally poorly by most metrics, including a 50% reduction in hard coral cover and in coral trout density on both NTMR and fished reefs. The reductions in coral trout density may reflect mortality or movement to less damaged reef areas [30]. There was some recovery after the storm, and the average size of coral trout was similar before and after the cyclone, which implies that relocation was more likely than widespread mortality. Such movement may be a response to the dramatic reduction in benthic habitat complexity in shallow




Figure 5. The Impacts of Tropical Cyclone Hamish on Communities of Fishes and Benthos on Offshore Reefs of the GBR

Analyses were based on a BACI design applied to the Markov chain Monte Carlo samples from the Bayesian hierarchical linear mixed model for each response variable. Plots give the median differences between values from before (2006–2008) and after (2010–2012) Tropical Cyclone Hamish ($\pm 95\%$ UIs) for reefs in the impact and control zones of the GBRMP. Closed symbols indicate NTMR reefs, and open symbols refer to reefs that were open to fishing. Control reefs (Cairns and Townsville sectors) were outside the destructive wind fields of Tropical Cyclone Hamish (Figure 1), whereas impact reefs (Pompey, Swain, and Capricorn Bunker sectors) were directly affected.

occur more frequently as climate change progresses [53]. The establishment of highly connected networks of NTMRs can contribute to a secure future for coral reefs, but effective measures to reduce land-based threats and to mitigate climate change will also be essential.

EXPERIMENTAL PROCEDURES

Sampling Protocols

Two systematic monitoring programs were instigated to assess the ecological effects of the new NTMRs after the implementation of the new GBRMP zoning plan in 2004. A team from James Cook University began surveying reef fish and benthic communities at three “inshore” island groups (fringing reefs on high continental islands within 30 km of the coast—Palm Islands, Whitsunday Islands, and Keppel Islands; Figure 1) in 2004 (prior to the re-zoning), while a team from the Australian Institute of Marine Science (AIMS) began surveys in five “offshore” latitudinal sectors (platform reefs >30 km from the coast—Cairns, Townsville, Pompey, Swain and Capricorn-Bunker) of the GBRMP in 2006 (Figure 1). Both programs surveyed NTMR reefs that were paired with similar reefs open to fishing. Despite minor differences in the details of the sampling protocols, comparable methods were used to collect all data (Table 1). Further details can be found in the [Supplemental Experimental Procedures](#).

Data Analyses

Benthic data (hard coral, soft coral, and algae) were expressed as percent cover. On the GBR, fishers using hook and line retain all species of “coral trout” (*Plectropomus* spp. and *Variola* spp.; family Serranidae) that are above the minimum legal size (38 cm TL), so density, size, and biomass estimates for all of these species were pooled. In addition, several species of “secondary targets,” which are not the main targets of fishers, are retained if caught (Table S1). Fish surveys using UVC recorded the counts and total lengths of coral trout and secondary target species on belt transects, whereas other reef fishes that were not targeted by fishing were only counted. All reef fish data were standardized by converting raw counts to densities $1,000\text{ m}^{-2}$. Biomass ($\text{kg } 1,000\text{ m}^{-2}$) was calculated for coral trout and secondary target species from estimated fish lengths (TL cm) using published length-weight relationships [54, 55]. We categorized non-target fishes into functional groups (Table S1). Further details can be found in the [Supplemental Experimental Procedures](#).

The spatial and temporal variation in the effects of NTMRs on the density and species richness of fish taxa and the percent cover of hard coral, soft coral, and algae were estimated using Bayesian hierarchical linear mixed models

Conclusions

The GBRMP zoning management plan appears to be performing as expected, given its geographic and socio-economic context. The expansion of NTMRs within the GBRMP coupled with effective direct fishery management actions have ensured adequate protection for stocks of key targeted coral reef fish species of the commercial and recreational fisheries and have lowered overall fishery catch to what currently appears to be sustainable levels [47]. Time will tell whether such levels prove to be sustainable, but if global temperatures and disturbance frequency increase in the future, we will face the prospect of having to reduce fishing pressure as target populations, both inside and outside NTMRs, suffer increasingly from non-fishery impacts. Monitoring and adaptive management would appear pertinent if we are to respond appropriately to changing conditions in the future and preserve fish stocks. There was little evidence of increased biodiversity within NTMRs compared with fished reefs, but this is not surprising given the limited range of species that are targeted by the fishery and the coarse measures of biodiversity used in this study. That the devastating effects of a severe tropical cyclone affected both NTMR and fished reefs equally is a timely reminder that NTMRs are not, by themselves, the solution for the full range of threats currently afflicting coral reefs. Pollution, sedimentation, coastal development, and the escalating effects of climate change all act at regional and global scales. Should we expect NTMRs to safeguard coral reefs from these threats? An encouraging finding from this study was that NTMRs can retain some benefits for key fishery-targeted species in the face of strong tropical cyclones that are predicted to

[56] (for details of models, see the [Supplemental Experimental Procedures](#)). Inferences about specific spatial and temporal differences between NTMRs and reefs open to fishing were based on 95% Bayesian UIs for modeled higher posterior density (HPD) median effects. Differences between values for NTMR and fished reefs were then expressed as a percentage of the value on the fished reefs, such that a higher value in NTMRs compared with fished reefs would yield a positive difference, whereas a lower value would give a negative difference.

Offshore reefs were not surveyed systematically before the new zoning plan was implemented in 2004, thus precluding the use of BACI analysis. However, estimates of coral trout biomass were available for 187 offshore reefs over the period of 1983–2012. Biomass samples from NTMRs and fished reefs in each latitudinal sector in each year were used to model trends using a Bayesian hierarchical linear mixed model. All models of biomass of coral trout were estimated using a linked, zero-inflated negative binomial (ZINB) model [57] in a Bayesian framework, using the PyMC package [58] for the Python programming language (for full model details, see the [Supplemental Experimental Procedures](#)). Coral trout were also surveyed on inshore reefs in the Palm and Whitsunday Islands (but not in the Keppel Islands) in the 1980s, and these estimates were compared with post-2004 values from these inshore island groups and also modeled using a linked ZINB model.

We explored the structure of reef fish and benthic assemblages graphically using RDA, looking for differences in assemblage structure attributable to reserve protection. Data were constrained by environmental predictors, in this case latitude (sector or island group) and zoning status (NTMR or open to fishing). The resulting variation in community structure was then partitioned among the constraining variables.

Finally, using Tropical Cyclone Hamish as a case study, we examined the effect of a regional scale disturbance on any effects of offshore NTMRs. Tropical Cyclone Hamish passed over reefs in the southern Pompey, Swain, and Capricorn-Bunker sectors in March 2009 ([Figure 1](#)). We applied a BACI design to the data and used Bayesian hierarchical models described above to evaluate the effects of the cyclone on the density and biomass of coral trout, on secondary target fishes, on functional groups of non-target fishes, and on hard coral cover at NTMR and fished reefs in the affected sectors. Further details of the BACI design can be found in the [Supplemental Experimental Procedures](#).

SUPPLEMENTAL INFORMATION

Supplemental Information includes Supplemental Results, Supplemental Experimental Procedures, three figures, and two tables and can be found with this article online at <http://dx.doi.org/10.1016/j.cub.2015.01.073>.

ACKNOWLEDGMENTS

Funding for the inshore reef monitoring program was provided by the Australian Research Council (ARC), the CRC Reef Research Centre, the Australian Government's Marine and Tropical Sciences Research Facility (MTSRF), and the National Environmental Research Program (NERP). The offshore reef monitoring program was primarily funded by the Australian Institute of Marine Science (AIMS). Additional funding was provided by the CRC Reef Research Centre, the Australian Government's Marine and Tropical Sciences Research Facility (MTSRF), and the National Environmental Research Program (NERP). We thank Tim Simmonds for assistance with the figures.

Received: October 4, 2014
 Revised: December 16, 2014
 Accepted: January 30, 2015
 Published: March 26, 2015

REFERENCES

- Hughes, T.P., Baird, A.H., Bellwood, D.R., Card, M., Connolly, S.R., Folke, C., Grosberg, R., Hoegh-Guldberg, O., Jackson, J.B.C., Kleypas, J., et al. (2003). Climate change, human impacts, and the resilience of coral reefs. *Science* 301, 929–933.
- Pandolfi, J.M., Bradbury, R.H., Sala, E., Hughes, T.P., Bjorndal, K.A., Cooke, R.G., McArdle, D., McClenachan, L., Newman, M.J.H., Paredes, G., et al. (2003). Global trajectories of the long-term decline of coral reef ecosystems. *Science* 301, 955–958.
- Bellwood, D.R., Hughes, T.P., Folke, C., and Nyström, M. (2004). Confronting the coral reef crisis. *Nature* 429, 827–833.
- Jackson, J.B.C., Donovan, M.K., Cramer, K.L., and Lam, V.V. (2014). Status and trends of caribbean coral reefs: 1970–2012. Global Coral Reef Monitoring Network, International Union for the Conservation of Nature. http://cmsdata.iucn.org/downloads/caribbean_coral_reefs__status_report_1970_2012.pdf.
- Lubchenco, J., Palumbi, S.R., Gaines, S.D., and Andelman, S. (2003). Plugging a hole in the ocean: the emerging science of marine reserves. *Ecol. Appl.* 13, S3–S7.
- Bohnsack, J.A. (1993). Marine reserves: they enhance fisheries, reduce conflicts and protect resources. *Oceanus* 36, 63–71.
- Halpern, B.S., and Warner, R.R. (2002). Marine reserves have rapid and lasting effects. *Ecol. Lett.* 5, 361–366.
- Halpern, B.S. (2003). The impact of marine reserves: do reserves work and does reserve size matter? *Ecol. Appl.* 13, 117–137.
- Claudet, J., Osenberg, C.W., Benedetti-Cecchi, L., Domenici, P., Garcia-Charton, J.-A., Pérez-Ruzafa, A., Badalamenti, F., Bayle-Sempere, J., Brito, A., Bulleri, F., et al. (2008). Marine reserves: size and age do matter. *Ecol. Lett.* 11, 481–489.
- Russ, G.R., and Alcala, A.C. (1996). Marine reserves: rates and patterns of recovery and decline of large predatory fish. *Ecol. Appl.* 6, 947–961.
- Williamson, D.H., Russ, G.R., and Ayling, A.M. (2004). No-take marine reserves increase abundance and biomass of reef fish on inshore fringing reefs of the Great Barrier Reef. *Environ. Conserv.* 31, 149–159.
- McClanahan, T.R., Graham, N.A.J., Calnan, J.M., and MacNeil, M.A. (2007). Toward pristine biomass: reef fish recovery in coral reef marine protected areas in Kenya. *Ecol. Appl.* 17, 1055–1067.
- Russ, G.R., Cheal, A.J., Dolman, A.M., Emslie, M.J., Evans, R.D., Miller, I., Sweatman, H., and Williamson, D.H. (2008). Rapid increase in fish numbers follows creation of world's largest marine reserve network. *Curr. Biol.* 18, R514–R515.
- Babcock, R.C., Shears, N.T., Alcala, A.C., Barrett, N.S., Edgar, G.J., Lafferty, K.D., McClanahan, T.R., and Russ, G.R. (2010). Decadal trends in marine reserves reveal differential rates of change in direct and indirect effects. *Proc. Natl. Acad. Sci. USA* 107, 18256–18261.
- Mosquera, I., Côté, I.M., Jennings, S., and Reynolds, J.D. (2000). Conservation benefits of marine reserves for fish populations. *Anim. Conserv.* 3, 321–332.
- Francini-Filho, R.B., and Moura, R.L. (2008). Evidence for spillover of reef fishes from a no-take marine reserve: an evaluation using the before-after control-impact (BACI) approach. *Fish. Res.* 93, 346–356.
- Lester, S.E., Halpern, B.S., Grorud-Colvert, K., Lubchenco, J., Ruttenberg, B.I., Gaines, S.D., Airame, S., and Warner, R.R. (2009). Biological effects within no-take marine reserves: a global synthesis. *Mar. Ecol. Prog. Ser.* 384, 33–46.
- Edgar, G.J., Stuart-Smith, R.D., Willis, T.J., Kininmonth, S., Baker, S.C., Banks, S., Barrett, N.S., Becerro, M.A., Bernard, A.T.F., Berkhout, J., et al. (2014). Global conservation outcomes depend on marine protected areas with five key features. *Nature* 506, 216–220.
- Abesamis, R.A., and Russ, G.R. (2005). Density-dependent spillover from a marine reserve: long-term evidence. *Ecol. Appl.* 15, 1798–1812.
- Harrison, H.B., Williamson, D.H., Evans, R.D., Almany, G.R., Thorrold, S.R., Russ, G.R., Feldheim, K.A., van Herwerden, L., Planes, S., Srinivasan, M., et al. (2012). Larval export from marine reserves and the recruitment benefit for fish and fisheries. *Curr. Biol.* 22, 1023–1028.
- Kerwath, S.E., Winker, H., Götz, A., and Attwood, C.G. (2013). Marine protected area improves yield without disadvantaging fishers. *Nat. Commun.* 4, 2347.

22. Graham, N.A.J., Ainsworth, T.D., Baird, A.H., Ban, N.C., Bay, L.K., Cinner, J.E., De Freitas, D.M., Diaz-Pulido, G., Dornelas, M., Dunn, S.R., et al. (2011). From microbes to people: tractable benefits of no-take areas for coral reefs. *Oceanogr. Mar. Biol. Annu. Rev.* **49**, 105–135.
23. Mumby, P.J., and Harborne, A.R. (2010). Marine reserves enhance the recovery of corals on Caribbean reefs. *PLoS ONE* **5**, e8657.
24. Selig, E.R., Casey, K.S., and Bruno, J.F. (2012). Temperature-driven coral decline: the role of marine protected areas. *Glob. Chang. Biol.* **18**, 1561–1570.
25. Graham, N.A.J., Evans, R.D., and Russ, G.R. (2003). The effects of marine reserve protection on the trophic relationships of reef fishes on the Great Barrier Reef. *Environ. Conserv.* **30**, 200–208.
26. McClanahan, T.R., Graham, N.A.J., MacNeil, M.A., Muthiga, N.A., Cinner, J.E., Bruggemann, J.H., and Wilson, S.K. (2011). Critical thresholds and tangible targets for ecosystem-based management of coral reef fisheries. *Proc. Natl. Acad. Sci. USA* **108**, 17230–17233.
27. Alcala, A.C., and Russ, G.R. (2006). No-take marine reserves and reef fisheries management in the Philippines: a new people power revolution. *Ambio* **35**, 245–254.
28. Selig, E.R., and Bruno, J.F. (2010). A global analysis of the effectiveness of marine protected areas in preventing coral loss. *PLoS ONE* **5**, e9278.
29. Jones, G.P., McCormick, M.I., Srinivasan, M., and Eagle, J.V. (2004). Coral decline threatens fish biodiversity in marine reserves. *Proc. Natl. Acad. Sci. USA* **101**, 8251–8253.
30. Williamson, D.H., Ceccarelli, D.M., Evans, R.D., Jones, G.P., and Russ, G.R. (2014). Habitat dynamics, marine reserve status, and the decline and recovery of coral reef fish communities. *Ecol. Evol.* **4**, 337–354.
31. Graham, N.A.J., Wilson, S.K., Jennings, S., Polunin, N.V.C., Bijoux, J.P., and Robinson, J. (2006). Dynamic fragility of oceanic coral reef ecosystems. *Proc. Natl. Acad. Sci. USA* **103**, 8425–8429.
32. Almany, G.R., Connolly, S.R., Heath, D.D., Hogan, J.D., Jones, G.P., McCook, L.J., Mills, M., Pressey, R.L., and Williamson, D.H. (2009). Connectivity, biodiversity conservation and the design of marine reserve networks for coral reefs. *Coral Reefs* **28**, 339–351.
33. Graham, N.A., McClanahan, T.R., MacNeil, M.A., Wilson, S.K., Polunin, N.V., Jennings, S., Chabanet, P., Clark, S., Spalding, M.D., Letourneur, Y., et al. (2008). Climate warming, marine protected areas and the ocean-scale integrity of coral reef ecosystems. *PLoS ONE* **3**, e3039.
34. DEEDI (2012). Annual status report 2011: coral reef fin fish fishery. The State of Queensland, Department of Employment, Economic Development and Innovation. https://www.daff.qld.gov.au/_data/assets/pdf_file/0017/66212/ASR_CRFFF2011.pdf
35. Osborne, K., Dolman, A.M., Burgess, S.C., and Johns, K.A. (2011). Disturbance and the dynamics of coral cover on the Great Barrier Reef (1995–2009). *PLoS ONE* **6**, e17516.
36. Sweatman, H., Delean, S., and Syms, C. (2011). Assessing the loss of coral cover on Australia's Great Barrier Reef over two decades, with implications for longer-term trends. *Coral Reefs* **30**, 521–531.
37. De'ath, G., Fabricius, K.E., Sweatman, H., and Puotinen, M. (2012). The 27-year decline of coral cover on the Great Barrier Reef and its causes. *Proc. Natl. Acad. Sci. USA* **109**, 17995–17999.
38. Diaz-Pulido, G., McCook, L.J., Dove, S., Berkelmans, R., Roff, G., Kline, D.I., Weeks, S., Evans, R.D., Williamson, D.H., and Hoegh-Guldberg, O. (2009). Doom and boom on a resilient reef: climate change, algal overgrowth and coral recovery. *PLoS ONE* **4**, e5239.
39. Berkelmans, R., Jones, A.M., and Schaffelke, B. (2012). Salinity thresholds of *Acropora* spp. on the Great Barrier Reef. *Coral Reefs* **31**, 1103–1110.
40. Russ, G.R., and Alcala, A.C. (2010). Decadal-scale rebuilding of predator biomass in Philippine marine reserves. *Oecologia* **163**, 1103–1106.
41. Kulbicki, M., Mou Tham, G., Thollot, P., and Wantiez, L. (1993). Length-weight relationships of fish from the lagoon of New Caledonia. *Naga ICLARM Q.* **16**, 26–30.
42. Australian Government Department of Environment and Heritage (2004). Marine protected areas and displaced fishing: a policy statement. <http://webarchive.nla.gov.au/gov/20070830033442/http://www.environment.gov.au/coasts/mpa/publications/displaced-fishing.html>.
43. Halpern, B.S., Gaines, S.D., and Warner, R.R. (2004). Confounding effects of the export of production and the displacement of fishing effort from marine reserves. *Ecol. Appl.* **14**, 1248–1256.
44. Buxton, C.D., Hartmann, K., Kearney, R., and Gardner, C. (2014). When is spillover from marine reserves likely to benefit fisheries? *PLoS ONE* **9**, e107032, <http://dx.doi.org/10.1371/journal.pone.0107032>.
45. Blarney, R.K., and Hundloe, T.J. (1991). Characteristics of Recreational Boat Fishing in the Great Barrier Reef Region. (Townsville: GBRMPA).
46. Williams, D.M., and Russ, G.R. (1994). In Review of Data on Fishes of Commercial and Recreational Fishing Interest in the Great Barrier Reef, *Volume 1*. (Townsville: GBRMPA).
47. Leigh, G.M., Campbell, A.B., Ludnow, C.P., and O'Neill, M.F. (2014). Stock Assessment of the Queensland East Coast Common Coral Trout (*Plectropomus leopardus*) Fishery. (Brisbane: Department of Agriculture, Fisheries and Forestry).
48. Fembach, M. (2008). A review of recreational activities undertaken in the Great Barrier Reef Marine Park (recreation review stage 1), Research Publication No. 93 (Townsville: GBRMPA). http://www.gbrmpa.gov.au/managing-the-reef/great-barrier-reef-outlook-report/outlook-report-2009/test-outlook-online?sq_content_src=%2BdXJsPWh0dHAIM0EIMkYIMkZ3d3ctb28uZ2JybXBhLmdvdvdi5hdSUyRm91dGxvb2slMkZncmVhdF9iYXJyaWVvYX3JlZWZfb3V0bG9va19yZXBvcnQIMkZvdXRsb29rX3JlcG9ydCUyRmV2aWRlbnNIJTJGMDFfc3RhbmRhcmlfZXZpZGVuY2VfcGFmZTEzMi5odG1sJmFsbD0x.
49. Emslie, M.J., Cheal, A.J., and Johns, K.A. (2014). Retention of habitat complexity minimizes disassembly of reef fish communities following disturbance: a large-scale natural experiment. *PLoS ONE* **9**, e105384.
50. Wen, C.K.C., Pratchett, M.S., Almany, G.R., and Jones, G.P. (2013). Role of prey availability in microhabitat preferences of juvenile coral trout (*Plectropomus*: Serranidae). *J. Exp. Mar. Biol. Ecol.* **443**, 39–45.
51. Berumen, M.L., and Pratchett, M.S. (2006). Recovery without resilience: persistent disturbance and long-term shifts in the structure of fish and coral communities at Tiahura Reef, Moorea. *Coral Reefs* **25**, 647–653.
52. Emslie, M.J., Pratchett, M.S., and Cheal, A.J. (2011). Effects of different disturbance types on butterflyfish communities of Australia's Great Barrier Reef. *Coral Reefs* **30**, 461–471.
53. Knutson, T.R., McBride, J.L., Chan, J., Emanuel, K., Holland, G., Landsea, C., Held, I., Kossin, J.P., Srivastava, A.K., and Sugi, M. (2010). Tropical cyclones and climate change. *Nat. Geosci.* **3**, 157–163.
54. Kulbicki, M., Guillemot, N., and Amand, M. (2005). A general approach to length-weight relationships for New Caledonian lagoon fishes. *Cybio* **29**, 235–252.
55. Froese, R., and Pauly, D., eds. (2014). FishBase, version 06/2014. <http://www.fishbase.org>.
56. Gelman, A., and Hill, J. (2007). *Data Analysis using Regression and Multilevel/Hierarchical Models*. (Cambridge: Cambridge University Press).
57. Smith, A.N.H., Anderson, M.J., and Millar, R.B. (2012). Incorporating the intraspecific occupancy-abundance relationship into zero-inflated models. *Ecology* **93**, 2526–2532.
58. Patil, A., Huard, D., and Fonnesbeck, C.J. (2010). PyMC: Bayesian stochastic modelling in Python. *J. Stat. Softw.* **35**, 1–81.