



BREED MIX

Dachshund : 100.0%

GENETIC STATS

Predicted adult weight: **15 lbs**Life stage: **Young adult**Based on your dog's date of birth provided.

TEST DETAILS

Kit number: EM-69092754 Swab number: 31220511509341

Registration: American Kennel Club

(AKC)








DNA Test Report Test Date: May 26th, 2023

embk.me/trumansagirlsbestfriend





The Dachshund was bred originally in Germany to flush out Badgers and other den animals in the 15th century. The breed, originally known as the Teckel, was refined by German Foresters to have the elongated shape that is advantageous for fitting into tight animal burrows. Dachshunds are often viewed as a symbol for Germany. For example, a Dachshund named Waldi was the first official mascot of the 1972 Summer Olympics held in Munich. Dachshunds are one of the most popular breeds in the United States, ranking 13th in AKC's most popular breeds. The Dachshund's personality is described as energetic, clever, and persistent to the point of stubbornness.

Alternative Names

Dachshund (Miniature), Dachshund (Standard)

Fun Fact

The name Dachshund is derived from "Dachs Krieger" meaning "Badger Warrior", who knew your Dachshund has such a fearsome name!

RELATED BREEDS



Basset HoundCousin breed



BeagleCousin breed



Bloodhound Cousin breed



Otterhound Cousin breed

Registration:







MATERNAL LINE



Through Bob's mitochondrial DNA we can trace his mother's ancestry back to where dogs and people first became friends. This map helps you visualize the routes that his ancestors took to your home. Their story is described below the map.

HAPLOGROUP: B1

B1 is the second most common maternal lineage in breeds of European or American origin. It is the female line of the majority of Golden Retrievers, Basset Hounds, and Shih Tzus, and about half of Beagles, Pekingese and Toy Poodles. This lineage is also somewhat common among village dogs that carry distinct ancestry from these breeds. We know this is a result of B1 dogs being common amongst the European dogs that their conquering owners brought around the world, because nowhere on earth is it a very common lineage in village dogs. It even enables us to trace the path of (human) colonization: Because most Bichons are B1 and Bichons are popular in Spanish culture, B1 is now fairly common among village dogs in Latin America.

HAPLOTYPE: B77/B81

Part of the B1 haplogroup, the B77/B81 haplotype occurs most frequently in Shih Tzus, Small Poodles, and American Bullies.

Registration: American Kennel Club

Kembark

(AKC)





PATERNAL LINE



Through Bob's Y chromosome we can trace his father's ancestry back to where dogs and people first became friends. This map helps you visualize the routes that his ancestors took to your home. Their story is described below the map.

HAPLOGROUP: A2b

A2b appears to have split a few times in succession, which means that some of the Central Asian male ancestors of this lineage went their separate ways before their respective Y chromosomes made their rounds. There is not much diversity in this lineage, meaning that it has only begun to take off recently. Two iconic breeds, the Dachshund and Bloodhound, represent this lineage well. Over half of Rottweilers are A2b, as are the majority of Labrador Retrievers and Cavalier King Charles Spaniels. While A2a is restricted mostly to East Asia, this paternal line is also found among European breeds.

HAPLOTYPE: Hc.9

Part of the A2b haplogroup, this haplotype is found in village dogs spanning South America, Africa, and the South Pacific. Among the breeds we have spotted it in, the most frequent occurrences are in Dachshund, Bloodhound, American Eskimo Dog, and Jack Russell Terrier.

Registration: American Kennel Club

Kembark

(AKC)





TRAITS: COAT COLOR

TRAIT RESULT

E Locus (MC1R)

The E Locus determines if and where a dog can produce dark (black or brown) hair. Dogs with two copies of the recessive **e** variant do not produce dark hairs and will express a red pigment called pheomelanin over their entire body. The shade of red, which can range from a deep copper to white, depends on other genetic factors, including the Intensity loci. In addition to determining if a dog can develop dark hairs, the E Locus can give a dog a black "mask" or "widow's peak" unless the dog has overriding coat color genetic factors.

No dark mask or grizzle (EE)

Dogs with one or two copies of the E^m variant may have a melanistic mask (dark facial hair as commonly seen in the German Shepherd Dog and Pug). In the absence of E^m, dogs with the E^g variant can have a "grizzle" phenotype (darker color on the head and top with a melanistic "widow's peak" and a lighter underside, commonly seen in the Afghan Hound and Borzoi and also referred to as "domino"). In the absence of both E^m and E variants, dogs with the E^a or E^h variants can express the grizzle phenotype. Additionally, a dog with any combination of two of the E^g, E^a, or E^h variants (example: E^gE^a) is also expected to express the grizzle phenotype.

K Locus (CBD103)

The K Locus K^B allele "overrides" the A Locus, meaning that it prevents the A Locus genotype from affecting coat color. For this reason, the K^B allele is referred to as the "dominant black" allele. As a result, dogs with at least one K^B allele will usually have solid black or brown coats (or red/cream coats if they are ee at the E Locus) regardless of their genotype at the A Locus, although several other genes could impact the dog's coat and cause other patterns, such as white spotting. Dogs with the $k^y k^y$ genotype will show a coat color pattern based on the genotype they have at the A Locus. Dogs who test as $K^B k^y$ may be brindle rather than black or brown.

More likely to have a patterned haircoat $(k^{y}k^{y})$









TRAITS: COAT COLOR (CONTINUED)

TRAIT RESULT

Intensity Loci

Areas of a dog's coat where dark (black or brown) pigment is not expressed either contain red/yellow pigment, or no pigment at all. Five locations across five chromosomes explain approximately 70% of red pigmentation "intensity" variation across all dogs. Dogs with a result of Intense Red Pigmentation will likely have deep red hair like an Irish Setter or "apricot" hair like some Poodles, dogs with a result of Intermediate Red Pigmentation will likely have tan or yellow hair like a Soft-Coated Wheaten Terrier, and dogs with Dilute Red Pigmentation will likely have cream or white hair like a Samoyed. Because the mutations we test may not directly cause differences in red pigmentation intensity, we consider this to be a linkage test.

Any light hair likely apricot or red (Intense Red Pigmentation)

A Locus (ASIP)

The A Locus controls switching between black and red pigment in hair cells, but it will only be expressed in dogs that are not **ee** at the E Locus and are **k**^y**k**^y at the K Locus. Sable (also called "Fawn") dogs have a mostly or entirely red coat with some interspersed black hairs. Agouti (also called "Wolf Sable") dogs have red hairs with black tips, mostly on their head and back. Black and tan dogs are mostly black or brown with lighter patches on their cheeks, eyebrows, chest, and legs. Recessive black dogs have solid-colored black or brown coats.

Black/Brown and tan coat color pattern (a^ta^t)

D Locus (MLPH)

The D locus result that we report is determined by three different genetic variants that can work together to cause diluted pigmentation. These are the common **d** allele, also known as "**d1**", and the less common alleles known as "**d2**" and "**d3**". Dogs with two **d** alleles, regardless of which variant, will have all black pigment lightened ("diluted") to gray, or brown pigment lightened to lighter brown in their hair, skin, and sometimes eyes. There are many breed-specific names for these dilute colors, such as "blue", "charcoal", "fawn", "silver", and "Isabella". Note that in certain breeds, dilute dogs have a higher incidence of Color Dilution Alopecia. Dogs with one **d** allele will not be dilute, but can pass the **d** allele on to their puppies.

Dark areas of hair and skin are not lightened (Dd)







TRAITS: COAT COLOR (CONTINUED)

TRAIT RESULT

Cocoa (HPS3)

Dogs with the **coco** genotype will produce dark brown pigment instead of black in both their hair and skin. Dogs with the **Nco** genotype will produce black pigment, but can pass the **co** allele on to their puppies. Dogs that have the **coco** genotype as well as the **bb** genotype at the B locus are generally a lighter brown than dogs that have the **Bb** or **BB** genotypes at the B locus.

No co alleles, not expressed (NN)

B Locus (TYRP1)

Dogs with two copies of the **b** allele produce brown pigment instead of black in both their hair and skin.

Dogs with one copy of the **b** allele will produce black pigment, but can pass the **b** allele on to their puppies.

E Locus **ee** dogs that carry two **b** alleles will have red or cream coats, but have brown noses, eye rims, and footpads (sometimes referred to as "Dudley Nose" in Labrador Retrievers). "Liver" or "chocolate" is the preferred color term for brown in most breeds; in the Doberman Pinscher it is referred to as "red".

Black or gray hair and skin (Bb)

Saddle Tan (RALY)

The "Saddle Tan" pattern causes the black hairs to recede into a "saddle" shape on the back, leaving a tan face, legs, and belly, as a dog ages. The Saddle Tan pattern is characteristic of breeds like the Corgi, Beagle, and German Shepherd. Dogs that have the II genotype at this locus are more likely to be mostly black with tan points on the eyebrows, muzzle, and legs as commonly seen in the Doberman Pinscher and the Rottweiler. This gene modifies the A Locus at allele, so dogs that do not express at are not influenced by this gene.

Not saddle tan patterned (II)

S Locus (MITF)

The S Locus determines white spotting and pigment distribution. MITF controls where pigment is produced, and an insertion in the MITF gene causes a loss of pigment in the coat and skin, resulting in white hair and/or pink skin. Dogs with two copies of this variant will likely have breed-dependent white patterning, with a nearly white, parti, or piebald coat. Dogs with one copy of this variant will have more limited white spotting and may be considered flash, parti or piebald. This MITF variant does not explain all white spotting patterns in dogs and other variants are currently being researched. Some dogs may have small amounts of white on the paws, chest, face, or tail regardless of their S Locus genotype.

Likely solid colored, but may have small amounts of white (Ssp)

Registration:







TRAITS: COAT COLOR (CONTINUED)

TRAIT RESULT

M Locus (PMEL)

Merle coat patterning is common to several dog breeds including the Australian Shepherd, Catahoula Leopard Dog, and Shetland Sheepdog, among many others. Merle arises from an unstable SINE insertion (which we term the "M*" allele) that disrupts activity of the pigmentary gene PMEL, leading to mottled or patchy coat color. Dogs with an M*m result are likely to be phenotypically merle or could be "non-expressing" merle, meaning that the merle pattern is very subtle or not at all evident in their coat. Dogs with an M*M* result are likely to be phenotypically merle or double merle. Dogs with an mm result have no merle alleles and are unlikely to have a merle coat pattern.

Note that Embark does not currently distinguish between the recently described cryptic, atypical, atypical+, classic, and harlequin merle alleles. Our merle test only detects the presence, but not the length of the SINE insertion. We do not recommend making breeding decisions on this result alone. Please pursue further testing for allelic distinction prior to breeding decisions.

One merle allele; may express merle (M*m)

Note: This locus includes several alleles. At the time this dog was genotyped Embark we could not distinguish all of the possible alleles.

R Locus (USH2A)

The R Locus regulates the presence or absence of the roan coat color pattern. Partial duplication of the USH2A gene is strongly associated with this coat pattern. Dogs with at least one **R** allele will likely have roaning on otherwise uniformly unpigmented white areas. Roan appears in white areas controlled by the S Locus but not in other white or cream areas created by other loci, such as the E Locus with **ee** along with Dilute Red Pigmentation by I Locus (for example, in Samoyeds). Mechanisms for controlling the extent of roaning are currently unknown, and roaning can appear in a uniform or non-uniform pattern. Further, non-uniform roaning may appear as ticked, and not obviously roan. The roan pattern can appear with or without ticking.

Likely no impact on coat pattern (rr)

H Locus (Harlequin)

This pattern is recognized in Great Danes and causes dogs to have a white coat with patches of darker pigment. A dog with an **Hh** result will be harlequin if they are also **M*m** or **M*M*** at the M Locus and are not **ee** at the E locus. Dogs with a result of **hh** will not be harlequin. This trait is thought to be homozygous lethal; a living dog with an **HH** genotype has never been found.

No harlequin alleles (hh)

Registration:







TRAITS: OTHER COAT TRAITS

TRAIT RESULT

Furnishings (RSPO2)

Dogs with one or two copies of the **F** allele have "furnishings": the mustache, beard, and eyebrows characteristic of breeds like the Schnauzer, Scottish Terrier, and Wire Haired Dachshund. A dog with two **I** alleles will not have furnishings, which is sometimes called an "improper coat" in breeds where furnishings are part of the breed standard. The mutation is a genetic insertion which we measure indirectly using a linkage test highly correlated with the insertion.

Likely unfurnished (no mustache, beard, and/or eyebrows) (II)









TRAITS: OTHER COAT TRAITS (CONTINUED)

TRAIT RESULT

Coat Length (FGF5)

The FGF5 gene affects hair length in many species, including cats, dogs, mice, and humans. In dogs, an **Lh** allele confers a long, silky hair coat across many breeds, including Yorkshire Terriers, Cocker Spaniels, and Golden Retrievers, while the **Sh** allele causes a shorter coat, as seen in the Boxer or the American Staffordshire Terrier. In certain breeds, such as the Pembroke Welsh Corgi and French Bulldog, the long haircoat is described as "fluffy". The coat length determined by FGF5, as reported by us, is influenced by four genetic variants that work together to promote long hair.

The most common of these is the **Lh1** variant (G/T, CanFam3.1, chr32, g.4509367) and the less common ones are **Lh2** (C/T, CanFam3.1, chr32, g.4528639), **Lh3** (16bp deletion, CanFam3.1, chr32, g.4528616), and **Lh4** (GG insertion, CanFam3.1, chr32, g.4528621). The FGF5_Lh1 variant is found across many dog breeds. The less common alleles, FGF5_Lh2, have been found in the Akita, Samoyed, and Siberian Husky, FGF5_Lh3 have been found in the Eurasier, and FGF5_Lh4 have been found in the Afghan Hound, Eurasier, and French Bulldog.

Likely short or midlength coat (ShLh)

The **Lh** alleles have a recessive mode of inheritance, meaning that two copies of the **Lh** alleles are required to have long hair. The presence of two Lh alleles at any of these FGF5 loci is expected to result in long hair. One copy each of **Lh1** and **Lh2** have been found in Samoyeds, one copy each of **Lh1** and **Lh3** have been found in Eurasiers, and one copy each of **Lh1** and **Lh4** have been found in the Afghan Hounds and Eurasiers.

Interestingly, the Lh3 variant, a 16 base pair deletion, encompasses the Lh4 variant (GG insertion). The presence of one or two copies of Lh3 influences the outcome at the Lh4 locus. When two copies of Lh3 are present, there will be no reportable result for the FGF5_Lh4 locus. With one copy of Lh3, Lh4 can have either one copy of the variant allele or the normal allele. The overall FGF5 result remains unaffected by this.









TRAITS: OTHER COAT TRAITS (CONTINUED)

TRAIT RESULT

Shedding (MC5R)

Dogs with at least one copy of the ancestral **C** allele, like many Labradors and German Shepherd Dogs, are heavy or seasonal shedders, while those with two copies of the **T** allele, including many Boxers, Shih Tzus and Chihuahuas, tend to be lighter shedders. Dogs with furnished/wire-haired coats caused by RSPO2 (the furnishings gene) tend to be low shedders regardless of their genotype at this gene.

Likely light shedding (TT)

Coat Texture (KRT71)

Dogs with a long coat and at least one copy of the **T** allele have a wavy or curly coat characteristic of Poodles and Bichon Frises. Dogs with two copies of the ancestral **C** allele are likely to have a straight coat, but there are other factors that can cause a curly coat, for example if they at least one **F** allele for the Furnishings (RSPO2) gene then they are likely to have a curly coat. Dogs with short coats may carry one or two copies of the **T** allele but still have straight coats.

Likely straight coat (CC)

Hairlessness (FOXI3)

A duplication in the FOXI3 gene causes hairlessness over most of the body as well as changes in tooth shape and number. This mutation occurs in Peruvian Inca Orchid, Xoloitzcuintli (Mexican Hairless), and Chinese Crested (other hairless breeds have different mutations). Dogs with the **NDup** genotype are likely to be hairless while dogs with the **NN** genotype are likely to have a normal coat. The **DupDup** genotype has never been observed, suggesting that dogs with that genotype cannot survive to birth. Please note that this is a linkage test, so it may not be as predictive as direct tests of the mutation in some lines.

Very unlikely to be hairless (NN)

Hairlessness (SGK3)

Hairlessness in the American Hairless Terrier arises from a mutation in the SGK3 gene. Dogs with the **DD** result are likely to be hairless. Dogs with the **ND** genotype will have a normal coat, but can pass the **D** variant on to their offspring.

Very unlikely to be hairless (NN)

Registration:







TRAITS: OTHER COAT TRAITS (CONTINUED)

TRAIT RESULT

Oculocutaneous Albinism Type 2 (SLC45A2)

Dogs with two copies **DD** of this deletion in the SLC45A2 gene have oculocutaneous albinism (OCA), also known as Doberman Z Factor Albinism, a recessive condition characterized by severely reduced or absent pigment in the eyes, skin, and hair. Affected dogs sometimes suffer from vision problems due to lack of eye pigment (which helps direct and absorb ambient light) and are prone to sunburn. Dogs with a single copy of the deletion **ND** will not be affected but can pass the mutation on to their offspring. This particular mutation can be traced back to a single white Doberman Pinscher born in 1976, and it has only been observed in dogs descended from this individual. Please note that this is a linkage test, so it may not be as predictive as direct tests of the mutation in some lines.

Likely not albino (NN)









TRAITS: OTHER BODY FEATURES

TRAIT RESULT

Muzzle Length (BMP3)

Dogs in medium-length muzzle (mesocephalic) breeds like Staffordshire Terriers and Labradors, and long muzzle (dolichocephalic) breeds like Whippet and Collie have one, or more commonly two, copies of the ancestral \mathbf{C} allele. Dogs in many short-length muzzle (brachycephalic) breeds such as the English Bulldog, Pug, and Pekingese have two copies of the derived \mathbf{A} allele. At least five different genes affect muzzle length in dogs, with BMP3 being the only one with a known causal mutation. For example, the skull shape of some breeds, including the dolichocephalic Scottish Terrier or the brachycephalic Japanese Chin, appear to be caused by other genes. Thus, dogs may have short or long muzzles due to other genetic factors that are not yet known to science.

Likely medium or long muzzle (CC)

Tail Length (T)

Whereas most dogs have two **C** alleles and a long tail, dogs with one **G** allele are likely to have a bobtail, which is an unusually short or absent tail. This mutation causes natural bobtail in many breeds including the Pembroke Welsh Corgi, the Australian Shepherd, and the Brittany Spaniel. Dogs with **GG** genotypes have not been observed, suggesting that dogs with the **GG** genotype do not survive to birth. Please note that this mutation does not explain every natural bobtail! While certain lineages of Boston Terrier, English Bulldog, Rottweiler, Miniature Schnauzer, Cavalier King Charles Spaniel, and Parson Russell Terrier, and Dobermans are born with a natural bobtail, these breeds do not have this mutation. This suggests that other unknown genetic mutations can also lead to a natural bobtail.

Likely normal-length tail (CC)

Hind Dewclaws (LMBR1)

Common in certain breeds such as the Saint Bernard, hind dewclaws are extra, nonfunctional digits located midway between a dog's paw and hock. Dogs with at least one copy of the **T** allele have about a 50% chance of having hind dewclaws. Note that other (currently unknown to science) mutations can also cause hind dewclaws, so some **CC** or **TC** dogs will have hind dewclaws.

Unlikely to have hind dew claws (CC)

Registration:







TRAITS: OTHER BODY FEATURES (CONTINUED)

TRAIT RESULT

Blue Eye Color (ALX4)

Embark researchers discovered this large duplication associated with blue eyes in Arctic breeds like Siberian Husky as well as tri-colored (non-merle) Australian Shepherds. Dogs with at least one copy of the duplication (**Dup**) are more likely to have at least one blue eye. Some dogs with the duplication may have only one blue eye (complete heterochromia) or may not have blue eyes at all; nevertheless, they can still pass the duplication and the trait to their offspring. **NN** dogs do not carry this duplication, but may have blue eyes due to other factors, such as merle. Please note that this is a linkage test, so it may not be as predictive as direct tests of the mutation in some lines.

Less likely to have blue eyes (NN)

Back Muscling & Bulk, Large Breed (ACSL4)

The **T** allele is associated with heavy muscling along the back and trunk in characteristically "bulky" large-breed dogs including the Saint Bernard, Bernese Mountain Dog, Greater Swiss Mountain Dog, and Rottweiler. The "bulky" **T** allele is absent from leaner shaped large breed dogs like the Great Dane, Irish Wolfhound, and Scottish Deerhound, which are fixed for the ancestral **C** allele. Note that this mutation does not seem to affect muscling in small or even mid-sized dog breeds with notable back muscling, including the American Staffordshire Terrier, Boston Terrier, and the English Bulldog.

Likely normal muscling (CC)









TRAITS: BODY SIZE

TRAIT		RESULT
Body Size (IGF1) The I allele is associated with smaller body size.	Smaller (II)	
Body Size (IGFR1) The A allele is associated with smaller body size.	Smaller (AA)	
Body Size (STC2) The A allele is associated with smaller body size.	Larger (TT)	
Body Size (GHR - E191K) The A allele is associated with smaller body size.	Smaller (AA)	
Body Size (GHR - P177L) The T allele is associated with smaller body size.	Larger (CC)	





TRAITS: PERFORMANCE

TRAIT RESULT

Altitude Adaptation (EPAS1)

This mutation causes dogs to be especially tolerant of low oxygen environments (hypoxia), such as those found at high elevations. Dogs with at least one $\bf A$ allele are less susceptible to "altitude sickness." This mutation was originally identified in breeds from high altitude areas such as the Tibetan Mastiff.

Normal altitude tolerance (GG)

Appetite (POMC)

This mutation in the POMC gene is found primarily in Labrador and Flat Coated Retrievers. Compared to dogs with no copies of the mutation (NN), dogs with one (ND) or two (DD) copies of the mutation are more likely to have high food motivation, which can cause them to eat excessively, have higher body fat percentage, and be more prone to obesity. Read more about the genetics of POMC, and learn how you can contribute to research, in our blog post (https://embarkvet.com/resources/blog/pomc-dogs/). We measure this result using a linkage test.

Normal food motivation (NN)









HEALTH REPORT

How to interpret Bob's genetic health results:

If Bob inherited any of the variants that we tested, they will be listed at the top of the Health Report section, along with a description of how to interpret this result. We also include all of the variants that we tested Bob for that we did not detect the risk variant for.

A genetic test is not a diagnosis

This genetic test does not diagnose a disease. Please talk to your vet about your dog's genetic results, or if you think that your pet may have a health condition or disease.

Summary

Of the 256 genetic health risks we analyzed, we found 3 results that you should learn about.

O Increased risk results (1)

Intervertebral Disc Disease (Type I)

Notable results (2)

ALT Activity

Progressive Retinal Atrophy, crd4/cord1

Clear results

Breed-relevant (6)

Other (246)

Registration: American Kennel Club

Hembark





BREED-RELEVANT RESULTS

Research studies indicate that these results are more relevant to dogs like Bob, and may influence his chances of developing certain health conditions.

○ Intervertebral Disc Disease (Type I) (FGF4 retrogene - CFA12)	Increased risk
Progressive Retinal Atrophy, crd4/cord1 (RPGRIP1)	Notable
	Clear
Mucopolysaccharidosis Type IIIA, Sanfilippo Syndrome Type A, MPS IIIA (SGSH Exon 6, Dachshund Variant)	Clear
Narcolepsy (HCRTR2 Exon 1, Dachshund Variant)	Clear
Neuronal Ceroid Lipofuscinosis 1, NCL 1 (PPT1 Exon 8, Dachshund Variant 1)	Clear
Neuronal Ceroid Lipofuscinosis 2, NCL 2 (TPP1 Exon 4, Dachshund Variant 2)	Clear
Osteogenesis Imperfecta (SERPINH1, Dachshund Variant)	Clear

Registration: American Kennel Club (AKC)







OTHER RESULTS

Research has not yet linked these conditions to dogs with similar breeds to Bob. Review any increased risk or notable results to understand his potential risk and recommendations.

ALT Activity (GPT)	Notable
② 2-DHA Kidney & Bladder Stones (APRT)	Clear
Acral Mutilation Syndrome (GDNF-AS, Spaniel and Pointer Variant)	Clear
Alaskan Husky Encephalopathy (SLC19A3)	Clear
Alaskan Malamute Polyneuropathy, AMPN (NDRG1 SNP)	Clear
Alexander Disease (GFAP)	Clear
Anhidrotic Ectodermal Dysplasia (EDA Intron 8)	Clear
Autosomal Dominant Progressive Retinal Atrophy (RHO)	Clear
Bald Thigh Syndrome (IGFBP5)	Clear
Bernard-Soulier Syndrome, BSS (GP9, Cocker Spaniel Variant)	Clear
Bully Whippet Syndrome (MSTN)	Clear
Canine Elliptocytosis (SPTB Exon 30)	Clear
Canine Fucosidosis (FUCA1)	Clear
Canine Leukocyte Adhesion Deficiency Type I, CLAD I (ITGB2, Setter Variant)	Clear
Canine Leukocyte Adhesion Deficiency Type III, CLAD III (FERMT3, German Shepherd Variant)	Clear
Canine Multifocal Retinopathy, cmr1 (BEST1 Exon 2)	Clear
Canine Multifocal Retinopathy, cmr2 (BEST1 Exon 5, Coton de Tulear Variant)	Clear
 Canine Multifocal Retinopathy, cmr3 (BEST1 Exon 10 Deletion, Finnish and Swedish Lapphund, Lapponian Herder Variant) 	Clear





 ✓ Canine Multiple System Degeneration (SERAC1 Exon 15, Kerry Blue Terrier Variant) ✓ Cardiomyopathy and Juvenile Mortality (YARS2) 	lear lear
	lear
Centronuclear Myopathy, CNM (PTPLA)	lear
Cerebellar Hypoplasia (VLDLR, Eurasier Variant)	lear
Chondrodystrophy (ITGA10, Norwegian Elkhound and Karelian Bear Dog Variant)	lear
Cleft Lip and/or Cleft Palate (ADAMTS20, Nova Scotia Duck Tolling Retriever Variant)	lear
Cleft Palate, CP1 (DLX6 intron 2, Nova Scotia Duck Tolling Retriever Variant)	lear
Cobalamin Malabsorption (CUBN Exon 8, Beagle Variant)	lear
Cobalamin Malabsorption (CUBN Exon 53, Border Collie Variant)	lear
○ Collie Eye Anomaly (NHEJ1)	lear
	lear
Openity County Station Discussion (NCDIII Objection Variant)	lear
Congenital Cornification Disorder (NSDHL, Chihuahua Variant)	
	lear
Congenital Hypothyroidism (TPO, Rat, Toy, Hairless Terrier Variant)	
 Congenital Hypothyroidism (TPO, Rat, Toy, Hairless Terrier Variant) Congenital Hypothyroidism (TPO, Tenterfield Terrier Variant) 	lear
 ✓ Congenital Hypothyroidism (TPO, Rat, Toy, Hairless Terrier Variant) ✓ Congenital Hypothyroidism (TPO, Tenterfield Terrier Variant) ✓ Congenital Hypothyroidism with Goiter (TPO Intron 13, French Bulldog Variant) 	lear lear





OTHER RESULTS

Congenital Myasthenic Syndrome, CMS (COLQ, Labrador Retriever Variant)	Clear
Congenital Myasthenic Syndrome, CMS (COLQ, Golden Retriever Variant)	Clear
Congenital Myasthenic Syndrome, CMS (CHAT, Old Danish Pointing Dog Variant)	Clear
Congenital Myasthenic Syndrome, CMS (CHRNE, Jack Russell Terrier Variant)	Clear
Congenital Stationary Night Blindness (LRIT3, Beagle Variant)	Clear
Congenital Stationary Night Blindness (RPE65, Briard Variant)	Clear
	Clear
Craniomandibular Osteopathy, CMO (SLC37A2 Intron 16, Basset Hound Variant)	Clear
Cystinuria Type I-A (SLC3A1, Newfoundland Variant)	Clear
Cystinuria Type II-A (SLC3A1, Australian Cattle Dog Variant)	Clear
Cystinuria Type II-B (SLC7A9, Miniature Pinscher Variant)	Clear
Oay Blindness (CNGB3 Deletion, Alaskan Malamute Variant)	Clear
Oay Blindness (CNGA3 Exon 7, German Shepherd Variant)	Clear
Oay Blindness (CNGA3 Exon 7, Labrador Retriever Variant)	Clear
Oay Blindness (CNGB3 Exon 6, German Shorthaired Pointer Variant)	Clear
O Deafness and Vestibular Syndrome of Dobermans, DVDob, DINGS (MYO7A)	Clear
	Clear
Degenerative Myelopathy, DM (SOD1A)	Cicai

Registration: American Kennel Club (AKC)







OTHER RESULTS

O Dental-Skeletal-Retinal Anomaly (MIA3, Cane Corso Variant)	Clear
Oiffuse Cystic Renal Dysplasia and Hepatic Fibrosis (INPP5E Intron 9, Norwich Terrier Variant)	Clear
Oilated Cardiomyopathy, DCM (RBM20, Schnauzer Variant)	Clear
Oilated Cardiomyopathy, DCM1 (PDK4, Doberman Pinscher Variant 1)	Clear
Dilated Cardiomyopathy, DCM2 (TTN, Doberman Pinscher Variant 2)	Clear
Disproportionate Dwarfism (PRKG2, Dogo Argentino Variant)	Clear
Ory Eye Curly Coat Syndrome (FAM83H Exon 5)	Clear
Opstrophic Epidermolysis Bullosa (COL7A1, Central Asian Shepherd Dog Variant)	Clear
Opstrophic Epidermolysis Bullosa (COL7A1, Golden Retriever Variant)	Clear
Early Bilateral Deafness (LOXHD1 Exon 38, Rottweiler Variant)	Clear
Early Onset Adult Deafness, EOAD (EPS8L2 Deletion, Rhodesian Ridgeback Variant)	Clear
Early Onset Cerebellar Ataxia (SEL1L, Finnish Hound Variant)	Clear
Ehlers Danlos (ADAMTS2, Doberman Pinscher Variant)	Clear
Enamel Hypoplasia (ENAM Deletion, Italian Greyhound Variant)	Clear
Enamel Hypoplasia (ENAM SNP, Parson Russell Terrier Variant)	Clear
Episodic Falling Syndrome (BCAN)	Clear
Exercise-Induced Collapse, EIC (DNM1)	Clear

Registration: American Kennel Club (AKC)







OTHER RESULTS

Factor XI Deficiency (F11 Exon 7, Kerry Blue Terrier Variant)	Clear
Familial Nephropathy (COL4A4 Exon 3, Cocker Spaniel Variant)	Clear
Familial Nephropathy (COL4A4 Exon 30, English Springer Spaniel Variant)	Clear
Fanconi Syndrome (FAN1, Basenji Variant)	Clear
Fetal-Onset Neonatal Neuroaxonal Dystrophy (MFN2, Giant Schnauzer Variant)	Clear
⊘ Glanzmann's Thrombasthenia Type I (ITGA2B Exon 13, Great Pyrenees Variant)	Clear
⊘ Glanzmann's Thrombasthenia Type I (ITGA2B Exon 12, Otterhound Variant)	Clear
Globoid Cell Leukodystrophy, Krabbe disease (GALC Exon 5, Terrier Variant)	Clear
Glycogen Storage Disease Type IA, Von Gierke Disease, GSD IA (G6PC, Maltese Variant)	Clear
Glycogen Storage Disease Type IIIA, GSD IIIA (AGL, Curly Coated Retriever Variant)	Clear
Glycogen storage disease Type VII, Phosphofructokinase Deficiency, PFK Deficiency (PFKM, Whippet and English Springer Spaniel Variant)	Clear
Glycogen storage disease Type VII, Phosphofructokinase Deficiency, PFK Deficiency (PFKM, Wachtelhund Variant)	Clear
GM1 Gangliosidosis (GLB1 Exon 2, Portuguese Water Dog Variant)	Clear
	Clear

Hembark





OTHER RESULTS

☑ Golden Retriever Progressive Retinal Atrophy 2, GR-PRA2 (TTC8) Clear ☑ Goniodysgenesis and Glaucoma, Pectinate Ligament Dysplasia, PLD (OLFM3) Clear ☑ Hemophilia A (F8 Exon 11, German Shepherd Variant 1) Clear ☑ Hemophilia A (F8 Exon 1, German Shepherd Variant 2) Clear ☑ Hemophilia A (F8 Exon 10, Boxer Variant) Clear ☑ Hemophilia B (F9 Exon 7, Terrier Variant) Clear ☑ Hemophilia B (F9 Exon 7, Rhodesian Ridgeback Variant) Clear ☑ Hereditary Ataxia, Cerebellar Degeneration (RAB24, Old English Sheepdog and Gordon Setter Variant) Clear ☑ Hereditary Cataracts (HSF4 Exon 9, Australian Shepherd Variant) Clear ☑ Hereditary Footpad Hyperkeratosis (FAM83G, Terrier and Kromfohrlander Variant) Clear ☑ Hereditary Nasal Parakeratosis (SUV39H2 Intron 4, Greyhound Variant) Clear ☑ Hereditary Nasal Parakeratosis, HNPK (SUV39H2) Clear ☑ Hereditary Vitamin D-Resistant Rickets (VDR) Clear ☑ Hypocatalasia, Acatalasemia (CAT) Clear ☑ Hypomyelination and Tremors (FNIP2, Weimaraner Variant) Clear ☑ Hypophosphatasia (ALPL Exon 9, Karelian Bear Dog Variant) Clear
 → Hemophilia A (F8 Exon 11, German Shepherd Variant 1) → Hemophilia A (F8 Exon 1, German Shepherd Variant 2) → Hemophilia A (F8 Exon 10, Boxer Variant) → Hemophilia B (F9 Exon 7, Terrier Variant) → Hemophilia B (F9 Exon 7, Rhodesian Ridgeback Variant) → Hemophilia B (F9 Exon 7, Rhodesian Ridgeback Variant) → Hereditary Ataxia, Cerebellar Degeneration (RAB24, Old English Sheepdog and Gordon Setter Variant) → Hereditary Cataracts (HSF4 Exon 9, Australian Shepherd Variant) → Hereditary Footpad Hyperkeratosis (FAM83G, Terrier and Kromfohrlander Variant) → Hereditary Footpad Hyperkeratosis (DSG1, Rottweiler Variant) → Hereditary Nasal Parakeratosis (SUV39H2 Intron 4, Greyhound Variant) → Hereditary Vitamin D-Resistant Rickets (VDR) → Hereditary Vitamin D-Resistant Rickets (VDR) → Hypocatalasia, Acatalasemia (CAT) → Hypomyelination and Tremors (FNIP2, Weimaraner Variant) → Clear → Hypomyelination and Tremors (FNIP2, Weimaraner Variant)
 ○ Hemophilia A (F8 Exon 1, German Shepherd Variant 2) ○ Hemophilia A (F8 Exon 10, Boxer Variant) ○ Clear ○ Hemophilia B (F9 Exon 7, Terrier Variant) ○ Clear ○ Hemophilia B (F9 Exon 7, Rhodesian Ridgeback Variant) ○ Clear ○ Hereditary Ataxia, Cerebellar Degeneration (RAB24, Old English Sheepdog and Gordon Setter Variant) ○ Clear ○ Hereditary Cataracts (HSF4 Exon 9, Australian Shepherd Variant) ○ Clear ○ Hereditary Footpad Hyperkeratosis (FAM83G, Terrier and Kromfohrlander Variant) ○ Clear ○ Hereditary Footpad Hyperkeratosis (DSG1, Rottweiler Variant) ○ Clear ○ Hereditary Nasal Parakeratosis (SUV39H2 Intron 4, Greyhound Variant) ○ Clear ○ Hereditary Vitamin D-Resistant Rickets (VDR) ○ Clear ○ Hypocatalasia, Acatalasemia (CAT) ○ Clear ○ Hypomyelination and Tremors (FNIP2, Weimaraner Variant) ○ Clear ○ Clear ○ Hypomyelination and Tremors (FNIP2, Weimaraner Variant)
 ✓ Hemophilia A (F8 Exon 10, Boxer Variant) ✓ Hemophilia B (F9 Exon 7, Terrier Variant) ✓ Hemophilia B (F9 Exon 7, Rhodesian Ridgeback Variant) ✓ Hereditary Ataxia, Cerebellar Degeneration (RAB24, Old English Sheepdog and Gordon Setter Variant) ✓ Hereditary Cataracts (HSF4 Exon 9, Australian Shepherd Variant) ✓ Hereditary Footpad Hyperkeratosis (FAM83G, Terrier and Kromfohrlander Variant) ✓ Hereditary Footpad Hyperkeratosis (DSG1, Rottweiler Variant) ✓ Hereditary Nasal Parakeratosis (SUV39H2 Intron 4, Greyhound Variant) ✓ Hereditary Vitamin D-Resistant Rickets (VDR) ✓ Hypocatalasia, Acatalasemia (CAT) ✓ Hypomyelination and Tremors (FNIP2, Weimaraner Variant)
 → Hemophilia B (F9 Exon 7, Terrier Variant) → Hemophilia B (F9 Exon 7, Rhodesian Ridgeback Variant) → Hereditary Ataxia, Cerebellar Degeneration (RAB24, Old English Sheepdog and Gordon Setter Variant) → Hereditary Cataracts (HSF4 Exon 9, Australian Shepherd Variant) → Hereditary Footpad Hyperkeratosis (FAM83G, Terrier and Kromfohrlander Variant) → Hereditary Footpad Hyperkeratosis (DSG1, Rottweiler Variant) → Hereditary Nasal Parakeratosis (SUV39H2 Intron 4, Greyhound Variant) → Hereditary Nasal Parakeratosis, HNPK (SUV39H2) → Hereditary Vitamin D-Resistant Rickets (VDR) → Hypocatalasia, Acatalasemia (CAT) → Hypomyelination and Tremors (FNIP2, Weimaraner Variant)
 → Hemophilia B (F9 Exon 7, Rhodesian Ridgeback Variant) → Hereditary Ataxia, Cerebellar Degeneration (RAB24, Old English Sheepdog and Gordon Setter Variant) → Hereditary Cataracts (HSF4 Exon 9, Australian Shepherd Variant) → Hereditary Footpad Hyperkeratosis (FAM83G, Terrier and Kromfohrlander Variant) → Hereditary Footpad Hyperkeratosis (DSG1, Rottweiler Variant) → Hereditary Nasal Parakeratosis (SUV39H2 Intron 4, Greyhound Variant) → Hereditary Nasal Parakeratosis, HNPK (SUV39H2) → Hereditary Vitamin D-Resistant Rickets (VDR) → Hypocatalasia, Acatalasemia (CAT) → Hypomyelination and Tremors (FNIP2, Weimaraner Variant)
 ✓ Hereditary Ataxia, Cerebellar Degeneration (RAB24, Old English Sheepdog and Gordon Setter Variant) ✓ Hereditary Cataracts (HSF4 Exon 9, Australian Shepherd Variant) ✓ Hereditary Footpad Hyperkeratosis (FAM83G, Terrier and Kromfohrlander Variant) ✓ Clear ✓ Hereditary Footpad Hyperkeratosis (DSG1, Rottweiler Variant) ✓ Clear ✓ Hereditary Nasal Parakeratosis (SUV39H2 Intron 4, Greyhound Variant) ✓ Clear ✓ Hereditary Nasal Parakeratosis, HNPK (SUV39H2) ✓ Hereditary Vitamin D-Resistant Rickets (VDR) ✓ Hypocatalasia, Acatalasemia (CAT) ✓ Clear ✓ Hypomyelination and Tremors (FNIP2, Weimaraner Variant)
 ✓ Hereditary Cataracts (HSF4 Exon 9, Australian Shepherd Variant) ✓ Hereditary Footpad Hyperkeratosis (FAM83G, Terrier and Kromfohrlander Variant) ✓ Hereditary Footpad Hyperkeratosis (DSG1, Rottweiler Variant) ✓ Hereditary Nasal Parakeratosis (SUV39H2 Intron 4, Greyhound Variant) ✓ Hereditary Nasal Parakeratosis, HNPK (SUV39H2) ✓ Hereditary Vitamin D-Resistant Rickets (VDR) ✓ Hypocatalasia, Acatalasemia (CAT) ✓ Hypomyelination and Tremors (FNIP2, Weimaraner Variant)
 ✓ Hereditary Footpad Hyperkeratosis (FAM83G, Terrier and Kromfohrlander Variant) ✓ Hereditary Footpad Hyperkeratosis (DSG1, Rottweiler Variant) ✓ Hereditary Nasal Parakeratosis (SUV39H2 Intron 4, Greyhound Variant) ✓ Hereditary Nasal Parakeratosis, HNPK (SUV39H2) ✓ Hereditary Vitamin D-Resistant Rickets (VDR) ✓ Hypocatalasia, Acatalasemia (CAT) ✓ Hypomyelination and Tremors (FNIP2, Weimaraner Variant)
 ✓ Hereditary Footpad Hyperkeratosis (DSG1, Rottweiler Variant) ✓ Hereditary Nasal Parakeratosis (SUV39H2 Intron 4, Greyhound Variant) ✓ Hereditary Nasal Parakeratosis, HNPK (SUV39H2) ✓ Hereditary Vitamin D-Resistant Rickets (VDR) ✓ Hypocatalasia, Acatalasemia (CAT) ✓ Hypomyelination and Tremors (FNIP2, Weimaraner Variant) ✓ Clear
 ✓ Hereditary Nasal Parakeratosis (SUV39H2 Intron 4, Greyhound Variant) ✓ Hereditary Nasal Parakeratosis, HNPK (SUV39H2) ✓ Hereditary Vitamin D-Resistant Rickets (VDR) ✓ Hypocatalasia, Acatalasemia (CAT) ✓ Hypomyelination and Tremors (FNIP2, Weimaraner Variant)
 ✓ Hereditary Nasal Parakeratosis, HNPK (SUV39H2) ✓ Hereditary Vitamin D-Resistant Rickets (VDR) ✓ Hypocatalasia, Acatalasemia (CAT) ✓ Hypomyelination and Tremors (FNIP2, Weimaraner Variant) Clear
 ✓ Hereditary Vitamin D-Resistant Rickets (VDR) ✓ Hypocatalasia, Acatalasemia (CAT) ✓ Hypomyelination and Tremors (FNIP2, Weimaraner Variant)
 Hypocatalasia, Acatalasemia (CAT) Hypomyelination and Tremors (FNIP2, Weimaraner Variant)
Hypomyelination and Tremors (FNIP2, Weimaraner Variant) Clear
Hypophosphatasia (ALPL Exon 9, Karelian Bear Dog Variant) Clear
⊘ Ichthyosis (NIPAL4, American Bulldog Variant) Clear

Registration: American Kennel Club (AKC)







☑ Ichthyosis (ASPRV1 Exon 2, German Shepherd Variant) Clear ☑ Ichthyosis (SLC27A4, Great Dane Variant) Clear ☑ Ichthyosis, Epidermolytic Hyperkeratosis (KRT10, Terrier Variant) Clear ☑ Ichthyosis, ICH1 (PNPLA1, Golden Retriever Variant) Clear ☑ Inflammatory Myopathy (SLC25A12) Clear ☑ Inherited Myopathy of Great Danes (BIN1) Clear ☑ Inherited Selected Cobalamin Malabsorption with Proteinuria (CUBN, Komondor Variant) Clear ☑ Intestinal Lipid Malabsorption (ACSL5, Australian Kelpie) Clear ☑ Junctional Epidermolysis Bullosa (LAMA3 Exon 66, Australian Cattle Dog Variant) Clear ☑ Juvenile Epidermolysis Bullosa (LAMB3 Exon 11, Australian Shepherd Variant) Clear ☑ Juvenile Epidermolysis and Polyneuropathy (RAB3GAP1, Rottweiller Variant) Clear ☑ Juvenile Myoclonic Epilepsy (DIRAS1) Clear ☑ L-2-Hydroxyglutaricaciduria, L2HGA (L2HGDH, Staffordshire Bull Terrier Variant) Clear ☑ Lagotto Storage Disease (ATG4D) Clear ☑ Laryngeal Paralysis (RAPGEF6, Miniature Bull Terrier Variant) Clear ☑ Late Onset Spinocerebellar Ataxia (CAPN1) Clear ☑ Late-Onset Neuronal Ceroid Lipofuscinosis, NCL 12 (ATP13A2, Australian Cattle Dog Variant) Clear		
✓ Ichthyosis, Epidermolytic Hyperkeratosis (KRT10, Terrier Variant) Clear ✓ Ichthyosis, ICH1 (PNPLA1, Golden Retriever Variant) Clear ✓ Inflammatory Myopathy (SLC25A12) Clear ✓ Inherited Myopathy of Great Danes (BIN1) Clear ✓ Inherited Selected Cobalamin Malabsorption with Proteinuria (CUBN, Komondor Variant) Clear ✓ Intestinal Lipid Malabsorption (ACSL5, Australian Kelpie) Clear ✓ Junctional Epidermolysis Bullosa (LAMA3 Exon 66, Australian Cattle Dog Variant) Clear ✓ Junctional Epidermolysis Bullosa (LAMB3 Exon 11, Australian Shepherd Variant) Clear ✓ Juvenile Epilepsy (LGI2) Clear ✓ Juvenile Laryngeal Paralysis and Polyneuropathy (RAB3GAP1, Rottweiler Variant) Clear ✓ Juvenile Myoclonic Epilepsy (DIRAS1) Clear ✓ Lagotto Storage Disease (ATG4D) Clear ✓ Lagotto Storage Disease (ATG4D) Clear ✓ Laryngeal Paralysis (RAPGEF6, Miniature Bull Terrier Variant) Clear ✓ Late Onset Spinocerebellar Ataxia (CAPN1) Clear	O Ichthyosis (ASPRV1 Exon 2, German Shepherd Variant)	Clear
⊘ Ichthyosis, ICH1 (PNPLA1, Golden Retriever Variant) Clear ⊘ Inflammatory Myopathy (SLC25A12) Clear ⊘ Inherited Myopathy of Great Danes (BIN1) Clear ⊘ Inherited Selected Cobalamin Malabsorption with Proteinuria (CUBN, Komondor Variant) Clear ⊘ Intestinal Lipid Malabsorption (ACSL5, Australian Kelpie) Clear ⊘ Junctional Epidermolysis Bullosa (LAMA3 Exon 66, Australian Cattle Dog Variant) Clear ⊘ Junctional Epidermolysis Bullosa (LAMB3 Exon 11, Australian Shepherd Variant) Clear ⊘ Juvenile Epilepsy (LGI2) Clear ⊘ Juvenile Laryngeal Paralysis and Polyneuropathy (RAB3GAP1, Rottweiler Variant) Clear ⊘ Juvenile Myoclonic Epilepsy (DIRAS1) Clear ⊘ Lagotto Storage Disease (ATG4D) Clear ⊘ Lagotto Storage Disease (ATG4D) Clear ⊘ Laryngeal Paralysis (RAPGEF6, Miniature Bull Terrier Variant) Clear ⊘ Late Onset Spinocerebellar Ataxia (CAPN1) Clear	O Ichthyosis (SLC27A4, Great Dane Variant)	Clear
✓ Inflammatory Myopathy (SLC25A12) Clear ✓ Inherited Myopathy of Great Danes (BIN1) Clear ✓ Inherited Selected Cobalamin Malabsorption with Proteinuria (CUBN, Komondor Variant) Clear ✓ Intestinal Lipid Malabsorption (ACSL5, Australian Kelpie) Clear ✓ Junctional Epidermolysis Bullosa (LAMA3 Exon 66, Australian Cattle Dog Variant) Clear ✓ Junctional Epidermolysis Bullosa (LAMB3 Exon 11, Australian Shepherd Variant) Clear ✓ Juvenile Epilepsy (LGI2) Clear ✓ Juvenile Laryngeal Paralysis and Polyneuropathy (RAB3GAP1, Rottweiler Variant) Clear ✓ Juvenile Myoclonic Epilepsy (DIRAS1) Clear ✓ L-2-Hydroxyglutaricaciduria, L2HGA (L2HGDH, Staffordshire Bull Terrier Variant) Clear ✓ Lagotto Storage Disease (ATG4D) Clear ✓ Laryngeal Paralysis (RAPGEF6, Miniature Bull Terrier Variant) Clear ✓ Late Onset Spinocerebellar Ataxia (CAPN1) Clear	Olichthyosis, Epidermolytic Hyperkeratosis (KRT10, Terrier Variant)	Clear
⊘ Inherited Myopathy of Great Danes (BIN1) Clear ⊘ Inherited Selected Cobalamin Malabsorption with Proteinuria (CUBN, Komondor Variant) Clear ⊘ Intestinal Lipid Malabsorption (ACSL5, Australian Kelpie) Clear ⊘ Junctional Epidermolysis Bullosa (LAMA3 Exon 66, Australian Cattle Dog Variant) Clear ⊘ Junctional Epidermolysis Bullosa (LAMB3 Exon 11, Australian Shepherd Variant) Clear ⊘ Juvenile Epilepsy (LGI2) Clear ⊘ Juvenile Laryngeal Paralysis and Polyneuropathy (RAB3GAP1, Rottweiler Variant) Clear ⊘ Juvenile Myoclonic Epilepsy (DIRAS1) Clear ⊘ L-2-Hydroxyglutaricaciduria, L2HGA (L2HGDH, Staffordshire Bull Terrier Variant) Clear ⊘ Lagotto Storage Disease (ATG4D) Clear ⊘ Laryngeal Paralysis (RAPGEF6, Miniature Bull Terrier Variant) Clear ⊘ Late Onset Spinocerebellar Ataxia (CAPN1) Clear	Olichthyosis, ICH1 (PNPLA1, Golden Retriever Variant)	Clear
Inherited Selected Cobalamin Malabsorption with Proteinuria (CUBN, Komondor Variant) Clear Intestinal Lipid Malabsorption (ACSL5, Australian Kelpie) Clear Junctional Epidermolysis Bullosa (LAMA3 Exon 66, Australian Cattle Dog Variant) Clear Junctional Epidermolysis Bullosa (LAMB3 Exon 11, Australian Shepherd Variant) Clear Juvenile Epilepsy (LGI2) Clear Juvenile Laryngeal Paralysis and Polyneuropathy (RAB3GAP1, Rottweiler Variant) Clear Juvenile Myoclonic Epilepsy (DIRAS1) Clear L-2-Hydroxyglutaricaciduria, L2HGA (L2HGDH, Staffordshire Bull Terrier Variant) Clear Lagotto Storage Disease (ATG4D) Clear Late Onset Spinocerebellar Ataxia (CAPN1) Clear	✓ Inflammatory Myopathy (SLC25A12)	Clear
 ☑ Intestinal Lipid Malabsorption (ACSL5, Australian Kelpie) ☑ Junctional Epidermolysis Bullosa (LAMA3 Exon 66, Australian Cattle Dog Variant) ☑ Junctional Epidermolysis Bullosa (LAMB3 Exon 11, Australian Shepherd Variant) ☑ Clear ☑ Juvenile Epidepsy (LGI2) ☑ Juvenile Laryngeal Paralysis and Polyneuropathy (RAB3GAP1, Rottweiler Variant) ☑ Juvenile Myoclonic Epidepsy (DIRAS1) ☑ L-2-Hydroxyglutaricaciduria, L2HGA (L2HGDH, Staffordshire Bull Terrier Variant) ☑ Lagotto Storage Disease (ATG4D) ☑ Laryngeal Paralysis (RAPGEF6, Miniature Bull Terrier Variant) ☑ Clear ☑ Late Onset Spinocerebellar Ataxia (CAPN1) 		Clear
 ✓ Junctional Epidermolysis Bullosa (LAMA3 Exon 66, Australian Cattle Dog Variant) ✓ Junctional Epidermolysis Bullosa (LAMB3 Exon 11, Australian Shepherd Variant) ✓ Juvenile Epilepsy (LGI2) ✓ Juvenile Laryngeal Paralysis and Polyneuropathy (RAB3GAP1, Rottweiler Variant) ✓ Juvenile Myoclonic Epilepsy (DIRAS1) ✓ L-2-Hydroxyglutaricaciduria, L2HGA (L2HGDH, Staffordshire Bull Terrier Variant) ✓ Lagotto Storage Disease (ATG4D) ✓ Laryngeal Paralysis (RAPGEF6, Miniature Bull Terrier Variant) ✓ Late Onset Spinocerebellar Ataxia (CAPN1) 	Inherited Selected Cobalamin Malabsorption with Proteinuria (CUBN, Komondor Variant)	Clear
 ✓ Junctional Epidermolysis Bullosa (LAMB3 Exon 11, Australian Shepherd Variant) ✓ Juvenile Epilepsy (LGI2) ✓ Juvenile Laryngeal Paralysis and Polyneuropathy (RAB3GAP1, Rottweiler Variant) ✓ Juvenile Myoclonic Epilepsy (DIRAS1) ✓ L-2-Hydroxyglutaricaciduria, L2HGA (L2HGDH, Staffordshire Bull Terrier Variant) ✓ Lagotto Storage Disease (ATG4D) ✓ Laryngeal Paralysis (RAPGEF6, Miniature Bull Terrier Variant) ✓ Clear ✓ Late Onset Spinocerebellar Ataxia (CAPN1) 	Intestinal Lipid Malabsorption (ACSL5, Australian Kelpie)	Clear
 ✓ Juvenile Epilepsy (LGI2) ✓ Juvenile Laryngeal Paralysis and Polyneuropathy (RAB3GAP1, Rottweiler Variant) ✓ Juvenile Myoclonic Epilepsy (DIRAS1) ✓ L-2-Hydroxyglutaricaciduria, L2HGA (L2HGDH, Staffordshire Bull Terrier Variant) ✓ Lagotto Storage Disease (ATG4D) ✓ Laryngeal Paralysis (RAPGEF6, Miniature Bull Terrier Variant) ✓ Clear ✓ Late Onset Spinocerebellar Ataxia (CAPN1) 	Junctional Epidermolysis Bullosa (LAMA3 Exon 66, Australian Cattle Dog Variant)	Clear
 ✓ Juvenile Laryngeal Paralysis and Polyneuropathy (RAB3GAP1, Rottweiler Variant) ✓ Juvenile Myoclonic Epilepsy (DIRAS1) ✓ L-2-Hydroxyglutaricaciduria, L2HGA (L2HGDH, Staffordshire Bull Terrier Variant) ✓ Lagotto Storage Disease (ATG4D) ✓ Laryngeal Paralysis (RAPGEF6, Miniature Bull Terrier Variant) ✓ Clear ✓ Late Onset Spinocerebellar Ataxia (CAPN1) ✓ Clear 	Junctional Epidermolysis Bullosa (LAMB3 Exon 11, Australian Shepherd Variant)	Clear
 ✓ Juvenile Myoclonic Epilepsy (DIRAS1) ✓ L-2-Hydroxyglutaricaciduria, L2HGA (L2HGDH, Staffordshire Bull Terrier Variant) ✓ Lagotto Storage Disease (ATG4D) ✓ Laryngeal Paralysis (RAPGEF6, Miniature Bull Terrier Variant) ✓ Late Onset Spinocerebellar Ataxia (CAPN1) 	Juvenile Epilepsy (LGI2)	Clear
 ∠ L-2-Hydroxyglutaricaciduria, L2HGA (L2HGDH, Staffordshire Bull Terrier Variant) ∠ Lagotto Storage Disease (ATG4D) ∠ Laryngeal Paralysis (RAPGEF6, Miniature Bull Terrier Variant) ∠ Late Onset Spinocerebellar Ataxia (CAPN1) 	Juvenile Laryngeal Paralysis and Polyneuropathy (RAB3GAP1, Rottweiler Variant)	Clear
 ∠ Lagotto Storage Disease (ATG4D) ∠ Laryngeal Paralysis (RAPGEF6, Miniature Bull Terrier Variant) ∠ Late Onset Spinocerebellar Ataxia (CAPN1) Clear	Juvenile Myoclonic Epilepsy (DIRAS1)	Clear
 ✓ Laryngeal Paralysis (RAPGEF6, Miniature Bull Terrier Variant) ✓ Late Onset Spinocerebellar Ataxia (CAPN1) Clear	L-2-Hydroxyglutaricaciduria, L2HGA (L2HGDH, Staffordshire Bull Terrier Variant)	Clear
 ✓ Late Onset Spinocerebellar Ataxia (CAPN1) 	Lagotto Storage Disease (ATG4D)	Clear
	Laryngeal Paralysis (RAPGEF6, Miniature Bull Terrier Variant)	Clear
Late-Onset Neuronal Ceroid Lipofuscinosis, NCL 12 (ATP13A2, Australian Cattle Dog Variant)	Late Onset Spinocerebellar Ataxia (CAPN1)	Clear
	Late-Onset Neuronal Ceroid Lipofuscinosis, NCL 12 (ATP13A2, Australian Cattle Dog Variant)	Clear





	Clear
	Clear
	Clear
Leukodystrophy (TSEN54 Exon 5, Standard Schnauzer Variant)	Clear
	Clear
 Limb Girdle Muscular Dystrophy (SGCD, Boston Terrier Variant) 	Clear
O Long QT Syndrome (KCNQ1)	Clear
Lundehund Syndrome (LEPREL1)	Clear
Macular Corneal Dystrophy, MCD (CHST6)	Clear
Malignant Hyperthermia (RYR1)	Clear
May-Hegglin Anomaly (MYH9)	Clear
Methemoglobinemia (CYB5R3, Pit Bull Terrier Variant)	Clear
	Clear
Microphthalmia (RBP4 Exon 2, Soft Coated Wheaten Terrier Variant)	Clear
Mucopolysaccharidosis IIIB, Sanfilippo Syndrome Type B, MPS IIIB (NAGLU, Schipperke Variant)	Clear
Mucopolysaccharidosis Type IIIA, Sanfilippo Syndrome Type A, MPS IIIA (SGSH Exon 6, New Zealand Huntaway Variant)	Clear
Mucopolysaccharidosis Type VI, Maroteaux-Lamy Syndrome, MPS VI (ARSB Exon 5, Miniature Pinscher Variant)	Clear
Mucopolysaccharidosis Type VII, Sly Syndrome, MPS VII (GUSB Exon 3, German Shepherd Variant)	Clear





Mucopolysaccharidosis Type VII, Sly Syndrome, MPS VII (GUSB Exon 5, Terrier Brasileiro Variant)	Clear
Multiple Drug Sensitivity (ABCB1)	Clear
Muscular Dystrophy (DMD, Cavalier King Charles Spaniel Variant 1)	Clear
Muscular Dystrophy (DMD, Golden Retriever Variant)	Clear
Musladin-Lueke Syndrome, MLS (ADAMTSL2)	Clear
Myasthenia Gravis-Like Syndrome (CHRNE, Heideterrier Variant)	Clear
Myotonia Congenita (CLCN1 Exon 23, Australian Cattle Dog Variant)	Clear
Myotonia Congenita (CLCN1 Exon 7, Miniature Schnauzer Variant)	Clear
Narcolepsy (HCRTR2 Intron 4, Doberman Pinscher Variant)	Clear
Narcolepsy (HCRTR2 Intron 6, Labrador Retriever Variant)	Clear
Nemaline Myopathy (NEB, American Bulldog Variant)	Clear
Neonatal Cerebellar Cortical Degeneration (SPTBN2, Beagle Variant)	Clear
Neonatal Encephalopathy with Seizures, NEWS (ATF2)	Clear
Neonatal Interstitial Lung Disease (LAMP3)	Clear
Neuroaxonal Dystrophy, NAD (VPS11, Rottweiler Variant)	Clear
Neuroaxonal Dystrophy, NAD (TECPR2, Spanish Water Dog Variant)	Clear
Neuronal Ceroid Lipofuscinosis 10, NCL 10 (CTSD Exon 5, American Bulldog Variant)	Clear
Neuronal Ceroid Lipofuscinosis 5, NCL 5 (CLN5 Exon 4 SNP, Border Collie Variant)	Clear





Neuronal Ceroid Lipofuscinosis 5, NCL 5 (CLN5 Exon 4 Deletion, Golden Retriever Variant)	Clear
Neuronal Ceroid Lipofuscinosis 6, NCL 6 (CLN6 Exon 7, Australian Shepherd Variant)	Clear
Neuronal Ceroid Lipofuscinosis 7, NCL 7 (MFSD8, Chihuahua and Chinese Crested Variant)	Clear
Neuronal Ceroid Lipofuscinosis 8, NCL 8 (CLN8, Australian Shepherd Variant)	Clear
Neuronal Ceroid Lipofuscinosis 8, NCL 8 (CLN8 Exon 2, English Setter Variant)	Clear
Neuronal Ceroid Lipofuscinosis 8, NCL 8 (CLN8 Insertion, Saluki Variant)	Clear
Neuronal Ceroid Lipofuscinosis, Cerebellar Ataxia, NCL4A (ARSG Exon 2, American Staffordshire Terrier Variant)	Clear
Oculocutaneous Albinism, OCA (SLC45A2 Exon 6, Bullmastiff Variant)	Clear
Oculocutaneous Albinism, OCA (SLC45A2, Small Breed Variant)	Clear
Oculoskeletal Dysplasia 2 (COL9A2, Samoyed Variant)	Clear
Osteochondrodysplasia (SLC13A1, Poodle Variant)	Clear
Osteogenesis Imperfecta (COL1A2, Beagle Variant)	Clear
Osteogenesis Imperfecta (COL1A1, Golden Retriever Variant)	Clear
P2Y12 Receptor Platelet Disorder (P2Y12)	Clear
Pachyonychia Congenita (KRT16, Dogue de Bordeaux Variant)	Clear
Paroxysmal Dyskinesia, PxD (PIGN)	Clear
Persistent Mullerian Duct Syndrome, PMDS (AMHR2)	Clear
Pituitary Dwarfism (POU1F1 Intron 4, Karelian Bear Dog Variant)	Clear





Platelet Factor X Receptor Deficiency, Scott Syndrome (TMEM16F)	Clear
Polycystic Kidney Disease, PKD (PKD1)	Clear
Pompe's Disease (GAA, Finnish and Swedish Lapphund, Lapponian Herder Variant)	Clear
Prekallikrein Deficiency (KLKB1 Exon 8)	Clear
Primary Ciliary Dyskinesia, PCD (NME5, Alaskan Malamute Variant)	Clear
Primary Ciliary Dyskinesia, PCD (CCDC39 Exon 3, Old English Sheepdog Variant)	Clear
Primary Hyperoxaluria (AGXT)	Clear
Primary Lens Luxation (ADAMTS17)	Clear
Primary Open Angle Glaucoma (ADAMTS17 Exon 11, Basset Fauve de Bretagne Variant)	Clear
Primary Open Angle Glaucoma (ADAMTS10 Exon 17, Beagle Variant)	Clear
Primary Open Angle Glaucoma (ADAMTS10 Exon 9, Norwegian Elkhound Variant)	Clear
Primary Open Angle Glaucoma and Primary Lens Luxation (ADAMTS17 Exon 2, Chinese Shar-Pei Variant)	Clear
Progressive Retinal Atrophy (SAG)	Clear
Progressive Retinal Atrophy (IFT122 Exon 26, Lapponian Herder Variant)	Clear
Progressive Retinal Atrophy, Bardet-Biedl Syndrome (BBS2 Exon 11, Shetland Sheepdog Variant)	Clear
Progressive Retinal Atrophy, CNGA (CNGA1 Exon 9)	Clear
Progressive Retinal Atrophy, crd1 (PDE6B, American Staffordshire Terrier Variant)	Clear
Progressive Retinal Atrophy, PRA1 (CNGB1)	Clear





⊘ Progressive Retinal Atrophy, prcd (PRCD Exon 1) ○ Cl ○ Progressive Retinal Atrophy, prcd (PRCD Exon 1) ○ Cl ○ Progressive Retinal Atrophy, prcd (PRCD Exon 1) ○ Cl ○ Progressive Retinal Atrophy, prcd (PRCD Exon 1) ○ Cl ○ Progressive Retinal Atrophy, prcd (PRCD Exon 1) ○ P	clear Clear
	loar
Progressive Retinal Atrophy, rcd1 (PDF6B Exon 21, Irish Setter Variant)	icai
(1 2 2 2 2 × 1 × 1 × 1 × 1 × 1 × 1 × 1 ×	lear
⊘ Progressive Retinal Atrophy, rcd3 (PDE6A) ○ CI	lear
Proportionate Dwarfism (GH1 Exon 5, Chihuahua Variant)	lear
⊘ Protein Losing Nephropathy, PLN (NPHS1)	lear
Pyruvate Dehydrogenase Deficiency (PDP1, Spaniel Variant)	lear
Pyruvate Kinase Deficiency (PKLR Exon 5, Basenji Variant)	lear
Pyruvate Kinase Deficiency (PKLR Exon 7, Beagle Variant)	lear
Pyruvate Kinase Deficiency (PKLR Exon 10, Terrier Variant)	lear
Pyruvate Kinase Deficiency (PKLR Exon 7, Labrador Retriever Variant)	lear
Pyruvate Kinase Deficiency (PKLR Exon 7, Pug Variant)	lear
Raine Syndrome (FAM20C)	lear
Recurrent Inflammatory Pulmonary Disease, RIPD (AKNA, Rough Collie Variant)	lear
Renal Cystadenocarcinoma and Nodular Dermatofibrosis (FLCN Exon 7)	lear
Retina Dysplasia and/or Optic Nerve Hypoplasia (SIX6 Exon 1, Golden Retriever Variant)	lear
Sensory Neuropathy (FAM134B, Border Collie Variant)	lear
	lear





Severe Combined Immunodeficiency, SCID (RAG1, Wetterhoun Variant)	Clear
Shaking Puppy Syndrome (PLP1, English Springer Spaniel Variant)	Clear
Shar-Pei Autoinflammatory Disease, SPAID, Shar-Pei Fever (MTBP)	Clear
Skeletal Dysplasia 2, SD2 (COL11A2, Labrador Retriever Variant)	Clear
Skin Fragility Syndrome (PKP1, Chesapeake Bay Retriever Variant)	Clear
Spinocerebellar Ataxia (SCN8A, Alpine Dachsbracke Variant)	Clear
Spinocerebellar Ataxia with Myokymia and/or Seizures (KCNJ10)	Clear
Spongy Degeneration with Cerebellar Ataxia 1 (KCNJ10)	Clear
Spongy Degeneration with Cerebellar Ataxia 2 (ATP1B2)	Clear
Stargardt Disease (ABCA4 Exon 28, Labrador Retriever Variant)	Clear
Succinic Semialdehyde Dehydrogenase Deficiency (ALDH5A1 Exon 7, Saluki Variant)	Clear
Thrombopathia (RASGRP1 Exon 5, American Eskimo Dog Variant)	Clear
Thrombopathia (RASGRP1 Exon 5, Basset Hound Variant)	Clear
Thrombopathia (RASGRP1 Exon 8, Landseer Variant)	Clear
	Clear
Ullrich-like Congenital Muscular Dystrophy (COL6A3 Exon 10, Labrador Retriever Variant)	Clear
Ullrich-like Congenital Muscular Dystrophy (COL6A1 Exon 3, Landseer Variant)	Clear
Unilateral Deafness and Vestibular Syndrome (PTPRQ Exon 39, Doberman Pinscher)	Clear





OTHER RESULTS

	Clear
	Clear
✓ Von Willebrand Disease Type II, Type II vWD (VWF, Pointer Variant)	Clear
Over Willebrand Disease Type III, Type III vWD (VWF Exon 4, Terrier Variant)	Clear
On Willebrand Disease Type III, Type III vWD (VWF Intron 16, Nederlandse Kooikerhondje Variant)	Clear
On Willebrand Disease Type III, Type III vWD (VWF Exon 7, Shetland Sheepdog Variant)	Clear
X-Linked Hereditary Nephropathy, XLHN (COL4A5 Exon 35, Samoyed Variant 2)	Clear
X-Linked Myotubular Myopathy (MTM1, Labrador Retriever Variant)	Clear
X-Linked Progressive Retinal Atrophy 1, XL-PRA1 (RPGR)	Clear
X-linked Severe Combined Immunodeficiency, X-SCID (IL2RG Exon 1, Basset Hound Variant)	Clear
X-linked Severe Combined Immunodeficiency, X-SCID (IL2RG, Corgi Variant)	Clear
Xanthine Urolithiasis (XDH, Mixed Breed Variant)	Clear
β-Mannosidosis (MANBA Exon 16, Mixed-Breed Variant)	Clear
Mast Cell Tumor	No result

Registration: American Kennel Club (AKC)







HEALTH REPORT



Increased risk result

Intervertebral Disc Disease (Type I)

Truman's a Girls Best Friend inherited both copies of the variant we tested for Chondrodystrophy and Intervertebral Disc Disease, CDDY/IVDD, Type I IVDD

Bob is at increased risk for Type I IVDD

How to interpret this result

Bob has two copies of an FGF4 retrogene on chromosome 12. In some breeds such as Beagles, Cocker Spaniels, and Dachshunds (among others) this variant is found in nearly all dogs. While those breeds are known to have an elevated risk of IVDD, many dogs in those breeds never develop IVDD. For mixed breed dogs and purebreds of other breeds where this variant is not as common, risk for Type I IVDD is greater for individuals with this variant than for similar dogs.

What is Chondrodystrophy and Intervertebral Disc Disease, CDDY/IVDD, Type I IVDD?

Type I Intervertebral Disc Disease (IVDD) is a back/spine issue that refers to a health condition affecting the discs that act as cushions between vertebrae. With Type I IVDD, affected dogs can have a disc event where it ruptures or herniates towards the spinal cord. This pressure on the spinal cord causes neurologic signs which can range from a wobbly gait to impairment of movement. Chondrodystrophy (CDDY) refers to the relative proportion between a dog's legs and body, wherein the legs are shorter and the body longer. There are multiple different variants that can cause a markedly chondrodystrophic appearance as observed in Dachshunds and Corgis. However, this particular variant is the only one known to also increase the risk for IVDD.

When signs & symptoms develop in affected dogs

Signs of CDDY are recognized in puppies as it affects body shape. IVDD is usually first recognized in adult dogs, with breed specific differences in age of onset.

Signs & symptoms

Research indicates that dogs with one or two copies of this variant have a similar risk of developing IVDD. However, there are some breeds (e.g. Beagles and Cocker Spaniels, among others) where this variant has been passed down to nearly all dogs of the breed and most do not show overt clinical signs of the disorder. This suggests that there are other genetic and environmental factors (such as weight, mobility, and family history) that contribute to an individual dog's risk of developing clinical IVDD. Signs of IVDD include neck or back pain, a change in your dog's walking pattern (including dragging of the hind limbs), and paralysis. These signs can be mild to severe, and if your dog starts exhibiting these signs, you should schedule an appointment with your veterinarian for a diagnosis.

How vets diagnose this condition

For CDDY, dogs with one copy of this variant may have mild proportional differences in their leg length. Dogs with two copies of this variant will often have visually longer bodies and shorter legs. For IVDD, a neurological exam will be performed on any dog showing suspicious signs. Based on the result of this exam, radiographs to detect the presence of calcified discs or advanced imaging (MRI/CT) to detect a disc rupture may be recommended.

How this condition is treated

Registration:







HEALTH REPORT



Notable result

ALT Activity

Truman's a Girls Best Friend inherited both copies of the variant we tested for Alanine Aminotransferase Activity

Why is this important to your vet?

Bob has two copies of a variant in the GPT gene and is likely to have a lower than average baseline ALT activity. ALT is a commonly used measure of liver health on routine veterinary blood chemistry panels. As such, your veterinarian may want to watch for changes in Bob's ALT activity above their current, healthy, ALT activity. As an increase above Bob's baseline ALT activity could be evidence of liver damage, even if it is within normal limits by standard ALT reference ranges.

What is Alanine Aminotransferase Activity?

Alanine aminotransferase (ALT) is a clinical tool that can be used by veterinarians to better monitor liver health. This result is not associated with liver disease. ALT is one of several values veterinarians measure on routine blood work to evaluate the liver. It is a naturally occurring enzyme located in liver cells that helps break down protein. When the liver is damaged or inflamed, ALT is released into the bloodstream.

How vets diagnose this condition

Genetic testing is the only way to provide your veterinarian with this clinical tool.

How this condition is treated

Veterinarians may recommend blood work to establish a baseline ALT value for healthy dogs with one or two copies of this variant.









HEALTH REPORT



Notable result

Progressive Retinal Atrophy, crd4/cord1

Truman's a Girls Best Friend inherited one copy of the variant we tested for Progressive Retinal Atrophy, crd4/cord1

What does this result mean?

This variant should not impact Bob's health. This variant is inherited in an autosomal recessive manner, meaning that a dog needs two copies of the variant to show signs of this condition. Bob is unlikely to develop this condition due to this variant because he only has one copy of the variant.

Impact on Breeding

Your dog carries this variant and will pass it on to ~50% of his offspring. You can email breeders@embarkvet.com to discuss with a genetic counselor how the genotype results should be applied to a breeding program.

What is Progressive Retinal Atrophy, crd4/cord1?

PRA-CRD4/cord1 is a retinal disease that causes progressive, non-painful vision loss over a 1-2 year period. The retina contains cells, called photoreceptors, that collect information about light and send signals to the brain. There are two types of photoreceptors: rods, for night vision and movement, and cones, for day vision and color. This type of PRA leads to early loss of cone cells, causing day blindness before night blindness.

When signs & symptoms develop in affected dogs

The earliest ophthalmic signs are typically present by 6 months of age. There is a wide range in the age of when dogs become clinically affected, although the average age is approximately 5 years. Dogs as young as 6 months may be blind, while dogs as old as 10 may still have vision.

How vets diagnose this condition

Veterinarians use a focused light to examine the pupils. In affected dogs, the pupils will appear more dilated and slower to contract. Your vet may also use a lens to visualize the retina at the back of the eye to look for changes in the optic nerve or blood vessels. You may be referred to a veterinary ophthalmologist for a definitive diagnosis.

How this condition is treated

Currently, there is no definitive treatment for PRA. Supplements, including antioxidants, have been proposed for management of the disease, but have not been scientifically proven effective.

Actions to take if your dog is affected

- Careful monitoring by your veterinarian will be required for the rest of your affected dog's life as secondary complications, including cataracts, can develop.
- With blind dogs, keeping furniture in the same location, making sure they are on a leash in unfamiliar territory, and training them to understand verbal commands are some of the ways to help them at home.










INBREEDING AND DIVERSITY

CATEGORY RESULT

Coefficient Of Inbreeding

Our genetic COI measures the proportion of your dog's genome where the genes on the mother's side are identical by descent to those on the father's side.



10%

MHC Class II - DLA DRB1

A Dog Leukocyte Antigen (DLA) gene, DRB1 encodes a major histocompatibility complex (MHC) protein involved in the immune response. Some studies have shown associations between certain DRB1 haplotypes and autoimmune diseases such as Addison's disease (hypoadrenocorticism) in certain dog breeds, but these findings have yet to be scientifically validated.

High Diversity

How common is this amount of diversity in purebreds:




MHC Class II - DLA DQA1 and DQB1

DQA1 and DQB1 are two tightly linked DLA genes that code for MHC proteins involved in the immune response. A number of studies have shown correlations of DQA-DQB1 haplotypes and certain autoimmune diseases; however, these have not yet been scientifically validated.

High Diversity

How common is this amount of diversity in purebreds:



Registration: American Kennel Club

(AKC)