



ANESTHESIA CONSIDERATIONS REGARDING PATIENT MEDICATIONS

Sometimes, anesthesia, oftentimes delivered with conscious sedation or monitored anesthesia care (MAC) is utilized during certain surgeries or procedure. Rarely, general anesthesia (GA) is required. There are certain considerations that need to be evaluated to effectively deliver safe and timely care.

MEDICATIONS THAT MAY NEED TO BE STOPPED PRIOR TO ANESTHESIA

Please ask your healthcare provider directly with any questions.

Medication (trade name)	(generic name)	Recommended discontinuation time (days)
Jardiance	Empagliflozin	3 days
Farxiga	Depagliflozin	3 days
Invokana	Canagliflozin	3 days
Ozempic	Semaglutide	If taken daily: hold day of the procedure If taken weekly, hold 7 days
Wegovy	Semaglutide	If taken daily: hold day of the procedure If taken weekly, hold 7 days
Trulicity	Dulaglutide	If taken daily: hold day of the procedure If taken weekly, hold 7 days
Mounjaro	Tirzepatide	If taken daily: hold day of the procedure If taken weekly, hold 7 days
Cozaar	Losartan	Hold day of the procedure, per Anesthesia Provider
Diovan	Valsartan	Hold day of the procedure, per Anesthesia Provider
Zestril; Prinivil; Obrelis	Lisinopril	Hold day of the procedure, per Anesthesia Provider

