



# BOVANS WHITE COMMERCIAL MANAGEMENT GUIDE

# INTRODUCTION

This guide highlights feeding, management and environmental factors, which can help achieve maximum profit from this genetically superior white egg layer.

These, combined with sound stock sense, are the prime ingredients required for good performance. It is

recommended that the guide be read completely prior to setting up a management program to ensure co-ordination of all phases of the life of the flock. This is important, particularly where growing and laying phases are separate operations. Improper treatment in the growing period can limit the pullet's potential.

## **ISA North America**

650 Riverbend Drive, Suite C  
Kitchener, Ontario  
Canada N2K 3S2

T +1 519 578 2740

F +1 519 578 1870

[www.isapoultry.com](http://www.isapoultry.com)

## **Institut de Sélection Animale BV**

Villa 'de Körver'  
Sporstraat 69, 5831 CK Boxmeer  
P.O. Box 114, 5830 AC Boxmeer  
The Netherlands-EU  
T +31 485 319 111  
F +31 485 319 112

[www.isapoultry.com](http://www.isapoultry.com)



# TABLE OF CONTENTS

<b>INTRODUCTION .....</b>	<b>1</b>	<b>BODY WEIGHTS &amp; APPROXIMATE FEED CONSUMPTION FOR BOVANS WHITE PULLETS.....</b>	<b>14</b>
<b>TABLE OF CONTENTS .....</b>	<b>2</b>	<b>UNIFORMITY .....</b>	<b>14</b>
<b>COMMERCIAL PERFORMANCE OBJECTIVES BOVANS WHITE..</b>	<b>4</b>	<b>WEIGH SCALES.....</b>	<b>15</b>
<b>SUGGESTED RATIONS.....</b>	<b>5</b>	<b>FEED IN GROWING PERIOD .....</b>	<b>15</b>
<b>RECOMMENDED PULLET FEEDING PROGRAM .....</b>	<b>5</b>	<b>MOVING TO THE LAYING HOUSE.....</b>	<b>15</b>
<b>RECOMMENDED VITAMIN TRACE MINERAL LEVELS .....</b>	<b>6</b>	<b>LIGHTING MANAGEMENT .....</b>	<b>15</b>
<b>BOVANS WHITE FEEDING RECOMMENDATIONS LAYING PERIOD.....</b>	<b>7</b>	<b>INTRODUCTION .....</b>	<b>15</b>
<b>THE BROODING PERIOD.....</b>	<b>8</b>	<b>LIGHTING PROGRAM FOR CONTROLLED ENVIRONMENT HOUSING .....</b>	<b>15</b>
<b>RECOMMENDED SPACE ALLOWANCE FROM DAY OLD TO 6 WEEKS.....</b>	<b>8</b>	<b>LIGHTING PROGRAM FOR OPEN-SIDED HOUSES AT THE EQUATOR.....</b>	<b>16</b>
PLANNING & PREPARATION .....	8	<b>LIGHTING PROGRAM FOR OPEN-SIDED HOUSES - NORTHERN HEMISPHERE.....</b>	<b>16</b>
CLEANING .....	8	<b>LIGHTING PROGRAM FOR OPEN-SIDED HOUSES - SOUTHERN HEMISPHERE .....</b>	<b>17</b>
ISOLATION.....	8	<b>THE BASICS OF DAYLENGTH ADJUSTMENT.....</b>	<b>17</b>
WATER.....	9	REARING .....	17
FEED .....	9	CONTROLLED ENVIRONMENT HOUSING .....	17
DANGER SIGNS.....	9	OPEN-SIDED HOUSING .....	17
<b>CAGE BROODING</b> .....	<b>9</b>	START OF LAY.....	17
TEMPERATURE .....	9	ON-OFF TIMES & TIME CLOCKS .....	18
HUMIDITY .....	9	<b>SUGGESTED LIGHT INTENSITIES .....</b>	<b>18</b>
PAPER.....	9	<b>LIGHT INTENSITY.....</b>	<b>18</b>
<b>FLOOR BROODING</b> .....	<b>9</b>	DIMMERS .....	18
TEMPERATURE .....	9	<b>THE PRE-LAY AND LAYING PERIODS ....</b>	<b>19</b>
CHICK GUARDS .....	9	<b>LAYING HOUSE REQUIREMENTS.....</b>	<b>19</b>
LITTER .....	10	<b>ADULT SPACE REQUIREMENT.....</b>	<b>19</b>
<b>BEAK TRIMMING .....</b>	<b>10</b>	<b>FEEDING DURING PRE-LAYING PERIOD .....</b>	<b>19</b>
AT HATCH .....	10	<b>FEEDING AT ONSET OF PRODUCTION .....</b>	<b>19</b>
AT 6-9 DAYS .....	10	<b>PROTEIN.....</b>	<b>20</b>
AT 10-12 WEEKS .....	10	<b>BODY WEIGHT IN THE ADULT PERIOD .....</b>	<b>20</b>
<b>INSOLUBLE GRIT .....</b>	<b>11-12</b>	<b>ENERGY.....</b>	<b>20</b>
<b>VENTILATION.....</b>	<b>11-12</b>	<b>BODY WEIGHT, EGG SIZE AND PRODUCTION INCREASE..</b>	<b>20</b>
<b>THE GROWING PERIOD .....</b>	<b>12</b>		
<b>REARING SPACE REQUIREMENTS .....</b>	<b>12</b>		
<b>MONITORING DEVELOPMENT.....</b>	<b>12</b>		
<b>REARING SPACE REQUIREMENTS 6-18 WEEKS .....</b>	<b>13</b>		
WEIGHING.....	13		
GROWTH RATE CONTROL .....	13		



ADJUSTED DENSITY FEEDING.....	20	PERITONITIS IN LAYERS .....	28
POST PEAK PROTEIN AND ENERGY .....	21	DISEASES OF THE DIGESTIVE SYSTEM .....	28
POST PEAK BODY WEIGHTS .....	21	DISEASES AFFECTING THE NERVOUS SYSTEM .....	28
EGG SIZE .....	21	DISEASES AFFECTING THE URINARY AND REPRODUCTIVE TRACTS .....	28
TEMPERATURE AND VENTILATION .....	21	DISEASES AFFECTING THE IMMUNE SYSTEM .....	28
CALCIUM .....	21	DISEASE AGENTS OF CONCERN FOR FOOD SAFETY ....	28
PHOSPHORUS .....	21		
<b>PULLET AND HEN HEALTH .....</b>	<b>22</b>	<b>APPENDIX 1.....</b>	<b>29</b>
BIOSECURITY .....	22	SUGGESTED DAILY NUTRIENT INTAKE FOR BOVANS WHITE COMMERCIAL PULLETS.....	29
WELFARE AND POULTRY HUSBANDRY .....	22		
DISEASE PREVENTION BY IMMUNIZATION .....	22	<b>APPENDIX 2.....</b>	<b>30</b>
TYPES OF VACCINES .....	22	SUGGESTED DAILY NUTRIENT ALLOWANCES FOR BOVANS WHITE COMMERCIAL.....	30
VACCINATION METHODS.....	23		
OCULAR (EYE DROP), BEAK DIPPING AND INTRANASAL VACCINATION .....	23	<b>APPENDIX 3.....</b>	<b>32</b>
SUBCUTANEOUS AND INTRAMUSCULAR INJECTION ....	23	BOVANS WHITE EGG PRODUCTION.....	32
TRANSCUTANEOUS INJECTION (WING WEB) .....	23		
VENT BRUSH VACCINATION .....	24	<b>APPENDIX 4.....</b>	<b>34</b>
IN OVO INJECTION .....	24	BOVANS WHITE EGG WEIGHT DISTRIBUTION (%) – CANADIAN SYSTEM .....	34
DRINKING WATER (ORAL) VACCINATION .....	24		
VACCINATION THROUGH A MEDICATOR .....	24	<b>APPENDIX 5.....</b>	<b>36</b>
WATER VACCINATION .....	24	BOVANS WHITE EGG WEIGHT DISTRIBUTION (%) – U.S.A. SYSTEM .....	36
SPRAY VACCINATION .....	25		
PARASITE CONTROL .....	26	<b>REARING GRAPH.....</b>	<b>38</b>
VACCINATION AGAINST COCCIDIOSIS .....	26	<b>PRODUCTION RECORDING GRAPH.....</b>	<b>39</b>
HISTOMONIASIS AND ROUND WORMS .....	27		
MITES .....	27	<b>EGG MASS &amp; CONSUMPTION GRAPH ..</b>	<b>40</b>
CONTROLLING GROUPS OF DISEASES BY VACCINATION .	27		
RESPIRATORY DISEASES .....	27		

The performance data contained in this document was obtained from results and experience from our own research flocks and flocks of our customers. In no way does the data contained in this document constitute a warranty or guarantee of the same performance under different conditions of nutrition, density or physical or biological environment. In particular (but without limitation of the foregoing) we do not grant any warranties regarding the fitness for purpose, performance, use, nature or quality of the flocks. ISA makes no representation as to the accuracy or completeness of the information contained in this document.



TABLE 1

## COMMERCIAL PERFORMANCE OBJECTIVES

**Bovans White****18-90 weeks**

<b>Body weight:</b>		
at 18 weeks	1260 g	2.78 lb.
at 30 weeks	1628 g	3.59 lb.
at 90 weeks	1720 g	3.79 lb
<b>Hen housed production:</b>		
at 60 weeks	254 eggs	
at 72 weeks	321 eggs	
at 90 weeks	411 eggs	
Age at 50% rate of lay	143 days	
Age at peak production	27-28 weeks	
Peak production	96 %	
<b>Production rate:</b>		
at 60 weeks	86.4 %	
at 72 weeks	80.4 %	
at 90 weeks	70.0 %	
<b>Total egg mass:</b>		
at 60 weeks	15.4 kg	
at 72 weeks	19.7 kg	
at 90 weeks	25.5 kg	
Average egg weight	62.2 g	49.4 lbs./100
Feed conversion* (ratio by weight)	2.08 kg/kg	
Average feed consumption during lay per day	106 g	23.4 lbs./100
Shell strength	4100 g	
Haug units	83.0	
Liveability (18-90 weeks)	94.1 %	

\* Data concerning feed conversion is based on controlled environmental temperature and phase feed under this guide's recommendation (Tables 2-4)



## SUGGESTED RATIONS

The choice of diet will be determined by factors such as price, type of ingredients and climate. The Bovans White is capable of good performance on a wide range of different feeds, as long as they are balanced properly. Suggested ration specifications are shown in Tables 2, 3 and 4.

TABLE 2

### RECOMMENDED PULLET FEEDING PROGRAM

Nutrients	Starter 1-8 weeks	Grower 1 9-12 weeks	Grower 2 13-16 or 17 weeks
Crude Protein (%)	20.5-21.0	18.0-18.5	17.0-17.5
ME (kcal/kg)	2850-2950	2750-2850	2700-2800
(kcal/lb)	1295-1340	1250-1295	1225-1270
Linoleic Acid (%)	1.30	1.00	1.00
Amino Acids (%)			
Methionine	0.48	0.40	0.38
Methionine + Cystine	0.83	0.70	0.65
Lysine	1.10	1.00	0.90
Tryptophan	0.23	0.21	0.18
Threonine	0.75	0.70	0.60
Arginine	1.20	1.10	1.00
Minerals (%)			
Calcium	1.05	1.05	1.10
Available Phosphorus	0.45-0.50	0.45-0.50	0.42-0.47
Sodium	0.18	0.18	0.18
Chloride (max.)	0.25	0.25	0.25

#### NOTE:

If pullets achieved 1200-1225 grams or 2.64-2.70 lbs. of body weight and good uniformity at 16 weeks of age it is possible to stimulate them with light and feed to achieve earlier egg production.

Feed Chick starter and grower rations containing a coccidiostat from day-old to 8-12 weeks of age when rearing flocks on litter floors without the use of a coccidiosis vaccine.



TABLE 3

**RECOMMENDED VITAMIN TRACE MINERAL LEVELS****added per tonne of complete feed**

Nutrients		Starter	Grower	Layer
Vitamin A	IU	12,000,000	9,000,000	11,000,000
Vitamin D3	ICU	2,500,000	2,500,000	3,000,000
Vitamin E	IU	30,000	20,000	20,000
Vitamin K3	g	3.0	3.0	3.0
Thiamine	g	2.5	2.5	2.2
Riboflavin	g	7.0	5.0	6.5
Pantothenic Acid	g	12.0	9.0	10.0
Niacin	g	40.0	30.0	40.0
Pyridoxine	g	5.0	3.5	4.5
Biotin	g	0.2	0.2	0.2
Folic Acid	g	1.0	1.0	1.0
Vitamin B12	g	0.03	0.02	0.02
Choline	g	1,000	800	1,000
Iron	g	80	80	80
Copper	g	10	10	10
Manganese	g	85	85	85
Zinc	g	80	80	80
Iodine	g	1.0	1.0	1.0
Selenium	g	0.3	0.3	0.3

**NOTE:**

Antioxidants should be added at levels recommended by the manufacturer. Antioxidants are especially important in hot climates and where fats are added to the ration.



TABLE 4

**BOVANS WHITE FEEDING RECOMMENDATIONS LAYING PERIOD**

Nutrients Age Range	Layer I 16/17-38 weeks	Layer II * 39-51 weeks	Layer III ** 52-64 weeks	Layer IV *** 65-77 weeks	Layer V **** 78-end
Feed Cons. Range	21-22.5 lbs/100 95-102 g/bird	21.4-22.9 lbs/100 97-104 g/bird	22-23 lbs/100 100-105 g/bird	22-23 lbs/100 100-105 g/bird	22-23 lbs/100 100-105 g/bird
Crude Protein (%)	18.0-18.5	17.5-18.0	16.5-17.0	15.5-16.0	15.0-15.5
ME (kcal/kg)	2860-2900	2850-2880	2840-2860	2820-2840	2820-2840
(kcal/lb)	1300-1320	1295-1310	1290-1300	1280-1290	1280-1290
Linoleic Acid (%)	1.90	1.60	1.40	1.30	1.30
Amino Acids (%)					
Methionine	0.48	0.45	0.42	0.39	0.36
Methionine + Cystine	0.80	0.76	0.72	0.68	0.62
Lysine	0.98	0.94	0.90	0.86	0.82
Tryptophan	0.22	0.19	0.18	0.17	0.16
Threonine	0.70	0.66	0.63	0.61	0.57
Arginine	0.96	0.92	0.88	0.84	0.80
Minerals (%)					
Calcium	4.1-4.2	4.2-4.3	4.3-4.4	4.4-4.5	4.6-4.7
Available Phosphorus	0.45-0.48	0.43-0.46	0.40-0.43	0.37-0.40	0.34-0.37
Sodium	0.18	0.18	0.18	0.18	0.18
Chloride (max.)	0.25	0.25	0.25	0.25	0.25

\* Changes from Layer I to Layer II should be made based on daily egg mass. After peak mass has been achieved (about 39 weeks of age), change from Layer I to Layer II.

\*\* Changes from Layer II to Layer III should be made around 52 weeks, when egg mass is about 57 grams.

\*\*\* Changes from Layer III to Layer IV should be made around 65 weeks, when egg mass is about 54 grams.

\*\*\*\* Changes from Layer IV to Layer V should be made around 78 weeks, when egg mass is about 50 grams. Changes should be subtle, if possible mixing both formulas for a week between each phase.

**NOTE:**

Daily egg mass output can be calculated by multiplying the actual hen-day rate of egg production by the average egg weight in grams (e.g. a flock laying 93% with an average egg weight of 60 grams has a daily egg mass output of 55.8 grams per bird).





# THE BROODING PERIOD

## FIRST DAY TO 6TH WEEK

The objective is for uninterrupted growth in order to achieve the correct body weight and frame development from day old through sexual maturity and during the egg production period.

The bird's ability to resist disease, to respond to vaccines, to reach the correct mature body size and, eventually, to

perform to its genetic potential, depends greatly on what happens during these first 6 weeks.

Lighting and vaccination programs, as well as body weight, all start during the brooding period. These, and other management aspects, are dealt with in the appropriate chapters.

TABLE 5

## RECOMMENDED SPACE ALLOWANCE FROM DAY OLD TO 6 WEEKS for optimum performance

	Cages	Litter	Litter & Slats
Floor	160 cm <sup>2</sup> (24.8 sq.in.) / bird	18 birds/m <sup>2</sup> (0.6 sq.ft./bird)	
Feeder    Trough Pans	5.0 cm (2") / bird -	5.0 cm (2") / bird 4/100 birds	
Drinker    Trough Birds/Round Bell Drinker	2.5 cm (1") / bird -	2.5 cm (1") / bird 90	
Cups or nipples	maximum 10 birds/cup or nipple a minimum of 2 cups or nipples per cage	maximum 10 birds/cup or nipple	

### NOTES:

1. With all types of equipment, do not exceed the manufacturer's recommendations for minimum floor, feeder and drinker space.
2. These recommendations reflect general practice. In some countries, legislation dictates greater minimum space allowances. Comply with the law.
3. In some countries, floor, feeder and drinker allowances are less than required for optimum performance. Under these conditions, lower performance is expected and accepted. Bovans birds perform competitively under all circumstances.
4. In hot weather conditions, allow 25% more space than figures above.

### PLANNING & PREPARATION

"All in, all out" is recommended.

### CLEANING

Remove all traces of manure, litter, dust, feathers, feed and any other residues from previous flocks. If stored or spread, keep manure at least 300 m (1,000 ft.) downwind from poultry houses.

Take steps to eliminate rodents, wild birds, insects and other pests. Carry out all repairs.

Flush and sanitize waterlines and tanks. Empty and clean bulk feed bins and storerooms.

The house and equipment, after cleaning and disinfecting, must be left to air dry for at least 10 days before the new flock arrives.

### ISOLATION

If buildings are less than 45 m (150 ft) apart, place young stock upwind from older birds. The attendant should not go near older flocks. If this is unavoidable, visit the younger flock first and change footwear, cap and overalls before tending the older birds. Hands should be washed too, using plenty of soap and water.

Place a 10 cm (4") deep bath of disinfectant solution at each house door. All who enter must first dip their footwear.

Clean the bath and replenish the disinfectant at least once daily. Keep the footbath sheltered from sun and rain.

Unless their presence is essential, allow no visitors in, or close to, the poultry house.



## WATER

Water intake restriction, whether accidental or deliberate, will reduce feed intake and growth rate. Water is also essential for body temperature regulation.

Before chicks arrive, have clean, fresh water ready. In a cool climate, it should be at room temperature, but even in hot regions, never warmer than 35°C (95°F).

Provide plenty of easily accessible, well-illuminated drinkers. For a fast start, especially when new arrivals have been heat stressed or dehydrated, give water, but no feed for the first two hours.

To limit wastage, begin raising drinkers as soon as possible. Nipples should be within “stretching” reach overhead. Cups and troughs should have their rims at mid neck height. Water depth must be sufficient to allow immersion of the whole beak, but for baby chicks, never so deep as to risk drowning.

## FEED

Feed conversion efficiency will never be better than during these first few weeks of life. Take full advantage of this. Go for maximum intake and thus maximum growth in the “Brooding Period”.

Use a good quality diet (see Tables 2-4). Crumbles are better than mash at this stage. Provide plenty of feeder space. Place the first feed on clean paper or new fibre egg trays, but as soon as possible, have the chicks eat from regular feed troughs.

Keep feeders full at the beginning but, within a week, reduce feed depth to minimize wastage. Chicks must still be able to eat without restriction.

By the end of the second week, the rims of both manual and automatic troughs should be at the level of the bird’s back.

## DANGER SIGNS

Watch for, listen, and react to the chick’s complaints. Their welfare is your profit. They can help you avoid mistakes.

Some indications are:

Loud chirping – hunger, cold, fear

Huddling together – cold, draughts

Prostrate, listless – too hot

## CAGE BROODING

### TEMPERATURE

Preheat the house for 24 hours before chicks arrive, maintaining 30-32°C (86-90°F) at cage level. Check temperatures at several different locations in the house.

To maintain health, growth and comfort of the flock, reduce temperature by about 3°C (5°F) per week. Aim for 22°C (72°F) at 21 days, but temper all adjustments to the behaviour and the real needs of the birds. When stressed, sick, or reacting to a vaccine, a flock’s need for warmth may increase temporarily, even beyond the age of 6 weeks.

Excessively high temperatures reduce feed intake. Any apparent advantage of lower consumption is more than wiped out by slower development and the lost opportunity to maximize growth while feed efficiency is at its peak.

Caged birds cannot escape to more comfortable areas. They rely on the attendant to get their environmental conditions right the first time.

## HUMIDITY

For optimum feather growth, health and feed conversion, maintain a relative humidity in the house of not less than 50%. If necessary, the cage house walkways and walls can be hosed down with a fine spray 3 or 4 times daily. Avoid getting water on the feed.

## PAPER

Several layers of newsprint, or similar absorbent, non slip paper over the wire floors will allow chicks immediate full use of the available space. It may also conserve heat and stop draughts.

Spread the paper under all drinkers to give quick, easy access to water. Peel back one layer at a time, as it becomes soiled. Remove paper altogether by 10 days. Not all cage designs need paper on the floor. Besides the high labour requirement it entails, paper may interfere with ventilation. Follow the recommendations of the cage manufacturer.

## FLOOR BROODING

### TEMPERATURE

Preheat for 24 hours before chick arrival and maintain a room temperature of 24°C (75°F).

Brooder temperatures, before chick arrival, should start at 32°C (90°F), 5 cm (2”) above the litter at the edge of the canopy. Aim to reduce this by 3°C (5°F) each week to a room temperature of 22°C (72°F) at 21 days.

Do not rely only on automatic controls or thermostats. Be guided by chick behaviour and see that their real needs are met.

## CHICK GUARDS

Use circular screens 38-46 cm (15-18”) high to confine chicks to the heated area. In cold seasons, the circle can be 1 m (about 3’ 3”) from the edge of the heater canopy. In hot seasons, leave 2 m (6’ 6”) or more space.

Cardboard or similar flexible, draught proof material can be used, but when night temperatures stay above 30°C (86°F), small mesh wire netting makes a better substitute.

Space all feeders and drinkers within the circle, but never under the direct heat of the brooder.

Gradually enlarge the circle, at least every second day, adding equipment as necessary and spacing it evenly. By 6 -10 days, remove the chick guards altogether.



## LITTER

Litter should be able to absorb and release moisture quickly. It should be non abrasive and non toxic.

Use 8 cm (3") of litter spread evenly over the floor. Concrete floors are recommended. Level the litter carefully in the brooder area. Do not cover it. Mould grows quickly even in new litter when paper or other materials exclude air.

Chick box lids, or similar containers, make good first time feeders and help to exclude litter from feeder troughs. Place drinkers on small boards for the same reason, but always ensure that the chicks have easy access to water.

Most manure will accumulate where the chicks rest. From the first week, turn this litter over to prevent caking. Let it become evenly "humid" rather than wet in spots. Promptly remove badly caked or soaking wet litter.

## **BEAK TRIMMING**

Where windowed or open sided houses are used, or where intensive systems of management are employed and when high-energy diets are fed, beak trimming is recommended for the control of cannibalism and to improve feed conversion. For the least interruption in growth, beaks should be precision trimmed by infrared beak treatment at hatch or at 6-9 days of age, not later. Trimmed carefully, they will not need touching again, but for assurance, inspect all at 10-12 weeks of age, when touch up is still feasible. Trimming after 10 days of age creates an unnecessary stress at a critical age, when a reduction in growth rate is least desired. As an alternative to trimming at this time, to burn the tip of the beak and subsequently do a precision trimming at 10-12 weeks has proven to be satisfactory, with least interference in growth and precise shape and size of beak.

### AT HATCH

- Use standard beak treatment protocol for the infrared beak treatment at hatch (i.e. 27/23c interface plate, glass radius mirror, treatment level 47 – chicks from PS flock younger than 30 weeks of age; 25/23 with treatment level 50 – chicks from PS flock 30 weeks and older).
- Infrared beak treated chicks should have very easy access to the water upon arrival to the brooding facility. Watering system could impact livability, weight and chicks uniformity. Usage of 360 degree nipples is strongly recommended.

### AT 6-9 DAYS

- Use the "baby chick" adapter on the machine.
- Select the guide hole that will allow the guillotine blade to make its cut 1-2 mm (3/64"-6/64") distal from the nostril. In most cases, this would be the 4.4 mm (11/64") diameter hole.
- The blade must be sharp, straight and heated to a cherry

red color (600°C or 1112°F). Blades must be replaced after every three hours of use. Do not clean it with metal scrapers.

- Insert the beak squarely into the guide hole. The thumb should be pressed lightly against the back of the chick's head, the forefinger at its throat to hold back the tongue.
- Depress the blade firmly and steadily to achieve a clean cut. Keep the blade down for a count of 2.5 seconds and keep the beak firmly against the blade.
- An automatic cam on the machine can take the guesswork out of that part of the operation and through controlling the blade movement, achieve better, more uniform results.
- Do not cauterize for longer than 2.5 seconds.
- If the cut continues to bleed, stop the machine and re-sharpen or, preferably, replace the blade. Do not increase the blade temperature above that recommended.
- One common cause of permanent beak damage is excessive cauterization.

If the job is done with precision, it will last the flock's lifetime. If done in haste, or without due care, re-trimming may be needed within a few weeks and, at that time, may seriously interrupt growth rate.

At the age of 6-10 days, a skilled operator should beak trim at the rate of 500-600 birds per hour, not faster.

### AT 10-12 WEEKS

- Inspect 100% of the pullets' beaks and select individuals that need a touch-up (birds with beak regrowth, of either upper or lower beak, or with sharp edges or "prongs", that may cause injury during eventual peck behavior). Depending on the precision of the first trimming, it may be even necessary to touch-up the whole flock, which is commonly done.
- Do not use any "baby chick" adapter, but only the beak-support bar (for older birds) on a manually operated beak-trimming machine.
- The blade must be at a higher temperature: 700°C - 1245°F.
- Cut and cauterize the upper and lower beak separately, to achieve ideal shape and size desired for upper and lower beak. A single block cut will give less than optimum results for control of regrowth.
- Cut the upper beak 5 mm from the nostrils, with the pullet's head on an angle 10° to 15° below horizontal. Cauterize the upper beak for 2.5-3.0 seconds, slightly turning the pullet head to the sides, but still pressing the cut edge against the centre of the blade, thus allowing a thorough rounding of the corners of the beak.
- Cut the lower beak at the same length or 1-2 mm longer than the upper beak, but never shorter. Cauterize it rounding the corners.




To ensure a rapid recovery from the inevitable stress caused by trimming, increase the depth of feed in feeders during the day of trimming and subsequent week.  
Provide water soluble Vitamin K and Vitamin C for two days

prior to trimming and three days after.  
Mid-night feeding for a week has provided satisfactory results in body weight recovery after trimming.

### Examples of proper infrared beak treated birds:

Infrared beak-treated chick at day of age



Infrared beak-treated hen at 17 weeks of age



Photos: courtesy off Nova-Tech Engineering, LLC

## INSOLUBLE GRIT

Insoluble grit should be given from the first day where birds have access to litter. It is an aid to feed digestion and also helps birds cope with any fibrous materials that they may ingest such as feathers, litter, etc.  
(Specifications on Table 6)

## VENTILATION

There are 5 main objectives for ventilation:

- to provide fresh air
- to remove stale air
- to control temperature
- to control humidity
- to remove dust

Each of these five must be satisfied if the flock is to perform to its best ability in feed conversion, livability, growth and egg production. (Specifications on Table 7)





TABLE 6

## INSOLUBLE GRIT

Floor	Grit Size	Litter Housing	Cages & Wire Floor
1 - 21 days	Baby Chick (1.0-1.5 mm diameter)	1 handful/200 birds/week	1 handful/200 birds/week
22 - 70 days	Grower (2.0-2.5 mm diameter)	500 g (1 lb.) / 100 birds/week	500 g (1 lb.) / 100 birds/month
71 days to end of lay	Layer (3.0-4.0 mm diameter)	500 g (1 lb.) / 100 birds/month	500 g (1 lb.) / 100 birds/month

NOTE: Amounts are approximate. Surplus grit in an automatic feeding system may cause damage.

TABLE 7

## VENTILATION RATES

Type of house	Fan Design Specifications at 2.0 mm (0.08 in) Static Pressure			
	Maximum Air Requirement/Bird		Minimum Air Requirement/Bird	
	m <sup>3</sup> /hr	C.F.M.	m <sup>3</sup> /hr	C.F.M.
Brooder House (0-6 wks)	3.5	2.1	0.5	0.3
Grower House (7-17 wks)	7.7	4.6	1.0	0.6
Layer House (17 wks-end of lay)	11.0	6.5	1.0	0.6

# THE GROWING PERIOD

## REARING SPACE REQUIREMENTS

The requirements for feeder, drinker and floor space are shown in Table 8. These are minimum requirements and should be increased during periods of hot weather and where ventilation or feed quality is marginal.

## MONITORING DEVELOPMENT

Important records to keep during the growing period include:

- body weight
- water intake
- temperature
- mortality
- feed intake
- hours of light
- vaccination

Other important information to record includes:

- type of feed (protein and energy values)
- any change of attendant

- equipment failures
- disease outbreaks
- beak trimming

Complete records, analyzed correctly, will provide the opportunity to make good management decisions to improve the profitability of current and future flocks.

Every time the records are updated, compare the actual information with the objective for that age. Any deviation from the expected target indicates that some form of management change is necessary to bring the flock back to the correct line of performance.

To develop a large framed, strong, lean pullet for optimum laying results, it is necessary to grow the pullet to the correct weight for age. To achieve and maintain target body weights, monitor growth by individually weighing a representative sample of the flock.



TABLE 8

## REARING SPACE REQUIREMENTS 6-18 WEEKS

	Cages	Litter	Litter & Slats
Floor	350 cm <sup>2</sup> (54 sq.in.) / bird	10 birds/m <sup>2</sup> (1.07 sq.ft./bird)	
Feeder    Trough Pans	7.5 cm (3") / bird -	7.5 cm (3") / bird 4/100 birds	
Drinker    Trough Birds/Round Bell Drinker	5.0 cm (2") / bird -	5.0 cm (2") / bird 90	
Cups or Nipples	maximum 10 birds/cup or nipple a minimum of 2 cups or nipples per cage	maximum 10 birds/cup or nipple	

### NOTES:

1. With all types of equipment, do not exceed the manufacturer's recommendations for minimum floor, feeder and drinker space.
2. These recommendations reflect general practice. In some countries, legislation dictates greater minimum space allowances. Comply with the law.
3. In some countries, floor, feeder and drinker allowances are less than required for optimum performance. Under these conditions, lower performance is expected and accepted. Bovans birds perform competitively under all circumstances.

### WEIGHING

Start weighing at 4 weeks of age.

In a cage house, select the cages containing the birds to be weighed from various locations and different levels in the house. Include some on both the delivery and the return sides of the automatic feeder. Always weigh every bird in the cage and mark the selected locations so that the same birds are weighed each time. In a litter house, use a lightweight screen to block off two areas in each pen and weigh every bird enclosed.

### GROWTH RATE CONTROL

Compare the bird's average body weight with the standard (Table 9). If on target at 28, 35 and 42 days, it will probably be safe to plan the change from starter to grower ration for the 50-57 day period. If body weights are not reaching the standard, postpone the feed change and/or take other measures to speed growth. In any case, it is normally not recommended to continue with chick starter feed beyond the 10th week.

#### Some Ways To Stimulate Feed Intake And Growth Rate

- a) If the age and health of the flock permit, reduce room temperatures (gradually) by 1 or 2°C (2 or 4°F).
- b) Increase light intensity over drinkers and feeders.

- c) Increase the number of drinkers per pen, or per cage.
- d) Increase the frequency of feeding, or feeder line operation.
- e) Increase depth of feed in the trough.
- f) Add feeders, or decrease bird numbers per pen, or per cage.
- g) Increase the energy level of the diet (adding fat).
- h) Increase the protein level in the chick starter.
- i) Verify the amino acid balance and vitamin/mineral levels in the diet.
- j) In rearing, delay the step-down lighting program or step down to a longer day length (but not more than 12 hours).
- k) An additional 1-hour of light in the middle of the dark period ("mid-night feeding") has also been shown to increase feed consumption.

Underweight flocks should be checked to determine the reason. Poor growth can be the result of a disease outbreak, low energy or low protein feed, heat stress, overcrowding and other environmental factors. Take prompt action to correct deficiencies and to stimulate feed intake.

Consult your feed supplier or nutritionist for further advice.



TABLE 9

## BODY WEIGHTS & APPROXIMATE FEED CONSUMPTION FOR BOVANS WHITE PULLETS

Age		Body weight minimum-maximum		Feed intake per bird per day minimum-maximum		Feed intake per bird cumulative minimum-maximum	
weeks	days	g	Lbs.	g/bird	Lbs./100 birds	g/bird	Lbs./100
1	0 - 7	64 - 66	0.14 - 0.15	6 - 8	1.32 - 1.76	42 - 56	9.3 - 12.4
2	8 - 14	118 - 122	0.26 - 0.27	12 - 14	2.64 - 3.09	126 - 154	27.8 - 33.9
3	15 - 21	185 - 195	0.41 - 0.43	20 - 22	4.41 - 4.85	266 - 308	58.6 - 67.9
4	22 - 28	258 - 272	0.57 - 0.60	29 - 31	6.39 - 6.83	469 - 525	103.4 - 115.7
5	29 - 35	336 - 354	0.74 - 0.78	36 - 38	7.94 - 8.38	721 - 791	158.9 - 174.4
6	36 - 42	419 - 441	0.92 - 0.97	40 - 42	8.82 - 9.26	1001 - 1085	220.7 - 239.2
7	43 - 49	502 - 528	1.11 - 1.16	43 - 45	9.48 - 9.92	1302 - 1400	287.0 - 308.6
8	50 - 56	585 - 615	1.29 - 1.35	45 - 47	9.92 - 10.36	1617 - 1729	356.5 - 381.2
9	57 - 63	668 - 702	1.47 - 1.55	47 - 49	10.36 - 10.80	1946 - 2072	429.0 - 456.8
10	64 - 70	746 - 784	1.64 - 1.73	49 - 51	10.80 - 11.24	2289 - 2429	504.6 - 535.5
11	71 - 77	824 - 866	1.82 - 1.91	51 - 53	11.24 - 11.68	2646 - 2800	583.3 - 617.3
12	78 - 84	902 - 948	1.99 - 2.09	53 - 55	11.68 - 12.12	3017 - 3185	665.1 - 702.2
13	85 - 91	975 - 1025	2.15 - 2.26	55 - 57	12.12 - 12.57	3402 - 3584	750.0 - 790.1
14	92 - 98	1048 - 1102	2.31 - 2.43	57 - 59	12.57 - 13.01	3801 - 3997	838.0 - 881.1
15	99 - 105	1112 - 1169	2.45 - 2.58	60 - 62	13.23 - 13.67	4221 - 4431	930.6 - 976.9
16	106 - 112	1165 - 1225	2.57 - 2.70	64 - 66	14.11 - 14.55	4669 - 4893	1029.3 - 1078.7
17	113 - 119	1204 - 1266	2.65 - 2.79	70 - 72	15.43 - 15.87	5159 - 5397	1137.4 - 1189.8
18	120 - 126	1229 - 1292	2.71 - 2.85	77 - 79	16.97 - 17.42	5698 - 5950	1256.2 - 1311.8

## UNIFORMITY

Commercial poultrymen apply their management techniques to the complete flock as a single unit because it is impractical to manage each bird as an individual. It is for this reason that flock uniformity is so important in obtaining optimum performance and the greatest profitability.

The lighting program, diet changes and vaccinations are applied to the whole flock. If it is uniform in weight and maturity, the flock will derive maximum benefit from these treatments. The better the uniformity in a laying flock, the higher the peak. Better persistency

of egg production, livability and uniformity of egg size can also be expected.

Wherever flock uniformity is less than desired, take action to improve it. Poor uniformity may be caused by disease, poor ventilation, lack of, or poor distribution of equipment, or any other management fault.

A good level of uniformity is when 80% of the birds are within plus or minus 10% of the flock's average weight.



## WEIGH SCALES

The "uniformity" of a flock may work out at 75% if weights are taken on a scale with 5-gram (0.2-oz) increments. If these weights are simultaneously measured on a scale with 20 gram (0.7 oz) increments, uniformity would appear better, in this case about 80%. With still larger increments, 50 grams or about 1/10 lb., the uniformity would appear to be as high as 85%.

Do not compare separate "uniformity" calculations unless the scales measure in the same increments.

At ISA, we recommend using a 5-kg or 11-lb. dial faced scale graduated in either 20 gram or 1/10-lb. increments.

## FEED IN GROWING PERIOD

Suggested ration specifications are provided in Tables 2, 3 and 4. These should be full fed. Where temperatures exceed 30°C (86°F), feed intake may be depressed and body weights fail to reach target. Under these circumstances, a change in diet formulation may be necessary. Consult your nutritionist for such changes. Starter ration should be fed

from week 1 to week 8; Grower 1 ration from 9 to 12 and Grower 2 ration from week 13 to 16 or 17.

## MOVING TO THE LAYING HOUSE

During and following the move from the growing facilities to the laying house, the birds are subject to many varied stresses. Make every effort to keep these factors to an absolute minimum.

Losses of between 5-10% in body weight can be common at this time. Every effort should be made following transfer to regain this weight and attain target weights. Failure to maintain growth in the early production period can result in low peak egg production, inadequate egg size and/or post peak dips in production.

Complete the move by 112 days (16 weeks) of age. This will allow the birds to settle down and become familiar with their new surroundings before production starts. The main vaccination program must be completed before the flock is moved. For the first 2-5 days in the new quarters, keep light intensity high until you are sure all the birds have located feed and water.

# LIGHTING MANAGEMENT

The lighting program, together with the recommended feeding and management programs contained in this guide, are designed to achieve the performance standards of Appendices 3 and 4. The BOVANS WHITE has shown great adaptability in egg size distribution to meet various market requirements. The purpose of controlling daylength during the rearing and laying periods is:

- to adjust maturity and onset of production for the correct age and stage of development
- to achieve the best rate of egg production
- to achieve required egg size
- to achieve adequate body weight

## INTRODUCTION

- An increasing photoperiod stimulates the bird to mature. A decreasing photoperiod retards maturity and will affect egg production.
- Daylength control in the rearing and laying periods is an important management tool in the achievement of economic performance, both in egg number and egg weight.
- The lighting program starts immediately day old chicks arrive.
- For suggested lighting programs, see pages 14 - 16.

TABLE 10

## LIGHTING PROGRAM FOR CONTROLLED ENVIRONMENT HOUSING

Age	Day length (hours)
Day 1 - 7	23
Day 8 - 14	20
Day 15 - 21	17
Day 22 - 28	14
Day 29 - 112	11
Week 17 (113 days)	12
Week 18	13
Week 19	13:30
Week 20	14
Week 21	14:30
Week 22	15
Week 23	15
Week 24	15
Week 25	15
Week 26	15
Week 27	15

NOTE: It is possible to begin light stimulation at 16 weeks of age if pullets achieved above target body weight (1200-1225 grams or 2.64-2.70 lbs.) and good uniformity (no less than 80%).





TABLE 11

### LIGHTING PROGRAM FOR OPEN-SIDED HOUSES AT THE EQUATOR

Age	Day length (hours)
Day 1 - 7	23
Day 8 - 14	20
Day 15 - 21	18
Day 22 - 28	16
Day 29 - 119	12
Week 18 (120 days)	13
Week 19	13:30
Week 20	14
Week 21	14:30
Week 22	15
Week 23	15:30
Week 24	16
Week 25	16
Week 26	16
Week 27	16
Week 28	16

TABLE 12-a

### LIGHTING PROGRAM FOR OPEN-SIDED HOUSES - NORTHERN HEMISPHERE

#### Hatches from April to September

Age	Day length (hours)
Day 1 - 7	22
Day 8 - 14	20
Day 15 - 21	18
Day 22 - 28	16
Day 29 - 112	Constant day length, equivalent to longest natural day in this period
Week 17 (113 days)	+ 1 h
Week 18	+30 min
Week 19	<div style="text-align: center;"> ↓  weekly increases of 30 minutes until reaching a total of 16 hours  ↓ </div>
Week 20	
Week 21	
Week 22	
Week 23	
Week 24	
Week 25	
Week 26	
Week 27	
Week 28	

TABLE 12-b

### NORTHERN HEMISPHERE

#### Hatches from October to March

Age	Day length (hours)
Day 1 - 7	22
Day 8 - 14	21
Day 15 - 21	20
Day 22 - 28	19
Day 29 - 35	18
Day 36 - 42	17
Day 43 - 119	Constant day length, equivalent to longest natural day in this period
Week 18 (120 days)	+ 1 h
Week 19	<div style="text-align: center;"> +30 min  ↓  weekly increases of 30 minutes until reaching a total of 16 hours  ↓ </div>
Week 20	
Week 21	
Week 22	
Week 23	
Week 24	
Week 25	
Week 26	

#### NOTE:

Flocks hatched from April to September will tend to be later into production than flocks hatched from October to March. Therefore, we recommend:

- Hatches from April to September:
  - Quicker step-down program in rearing and an earlier stimulus of one extra hour at week 17.
  - Constant Light between 5 and 16 weeks. Total day length never shorter than longest natural day in the period.
- Hatches from October to March:
  - Longer step-down program in rearing, to ensure proper weight gain.
  - Constant Light between 7 and 17 weeks. Total day length never shorter than longest natural day in the period.



TABLE 13-a

## LIGHTING PROGRAM FOR OPEN-SIDED HOUSES - SOUTHERN HEMISPHERE

### Hatches from April to September

Age	Day length (hours)
Day 1 - 7	22
Day 8 - 14	21
Day 15 - 21	20
Day 22 - 28	19
Day 29 - 35	18
Day 36 - 42	17
Day 43 - 119	Constant day length, equivalent to longest natural day in this period
Week 18 (120 days)	+ 1 h
Week 19	+30 min
Week 20	↓
Week 21	weekly increases of 30
Week 22	minutes until reaching a
Week 23	total of 16 hours
Week 24	↓
Week 25	
Week 26	

TABLE 13-b

## SOUTHERN HEMISPHERE

### Hatches from October to March

Age	Day length (hours)
Day 1 - 7	22
Day 8 - 14	20
Day 15 - 21	18
Day 22 - 28	16
Day 29 - 112	Constant day length, equivalent to longest natural day in this period
Week 17 (113 days)	+ 1 h
Week 18	+30 min
Week 19	↓
Week 20	weekly increases of 30
Week 21	minutes until reaching a
Week 22	total of 16 hours
Week 23	↓
Week 24	
Week 25	
Week 26	
Week 27	
Week 28	

### NOTE:

Flocks hatched from October to March will tend to be later into production than flocks hatched from April to September. Therefore, we recommend:

- Hatches from October to March:
  - Quicker step-down program in rearing and an earlier stimulus of one extra hour at week 17.
  - Constant Light between 5 and 16 weeks. Total day length never shorter than longest natural day in the period.
- Hatches from April to September:
  - Longer step-down program in rearing, to ensure proper weight gain.
  - Constant Light between 7 and 17 weeks. Total day length never shorter than longest natural day in the period.

## THE BASICS OF DAYLENGTH ADJUSTMENT

### REARING

In rearing, use a "Step-down" or a "Constant" lighting pattern, or a combination of the two.

A step-down program tends to retard maturity and increase lifetime average egg size, more so if applied beyond 42 days of age. However, in the early weeks, when chicks are growing rapidly, longer days stimulate higher feed intake and maximum bone growth at the least cost.

In fact, the light program is flexible. A decision not to reduce day length as quickly as planned may be taken if growth is insufficient. However, this may delay sexual maturity and affect egg size.

### CONTROLLED ENVIRONMENT HOUSING

See Table 10

### OPEN-SIDED HOUSING

Birds should be reared on a day length equal to the longest natural photoperiod that they will encounter during the period of 6-18 weeks of age.

An example is provided in Table 11 for the situation encountered at the equator with a constant day length of 12 hours. For recommendations that are more suited to your particular conditions, please consult your local Bovans representative.

### START OF LAY

Never increase day length until the rearing period is over and then time it carefully to match the needs of the flock in terms of body weight and physical maturity. If body size and development have been retarded, the day length increase should be postponed a few days. The timing of photostimulation should be based on body weight. A precocious, well-grown flock could be stimulated safely earlier than the Guide suggests, depending on your particular



egg size requirements. Light stimulation at 16 weeks of age can only be safely employed if birds are on target weight for 17 weeks of age (minimum 1204 g or 2.65 lbs). Obviously, any deficiency in body weight will reduce average egg weight.

In fully blacked out, light proofed houses, where daylight cannot enter, a year round day length of 15 hours is recommended once egg production has peaked (see Table 10).

Sixteen hours will also support maximum production in open sided houses, wherever the longest day of the year is 14 hours or less.

Whenever the longest natural day is more than 15 hours and the house not 100% light proof, day length throughout lay must at least equal the longest natural day. This ensures that the flock will never experience shortening days after mid summer, a factor that could induce neck moult, or a significant drop in egg production.

ON-OFF TIMES & TIME CLOCKS

Light “On” and “Off” times are usually arranged to suit working hours. A popular system is for lights to come on one hour before attendants arrive. “Off” time in the evening allows for a routine inspection of the flock after the lights are out.

The regular, repeated pattern of lights on, lights off is the stimulation for a flock’s egg production response. If the pattern is interrupted or irregular, the full benefit of lighting may be lost. As a result, egg production will suffer, especially in older flocks (over 40 weeks).

Reliable, fully automatic, accurate and properly maintained electric time clocks are strongly recommended.

TABLE 14  
**SUGGESTED LIGHT INTENSITIES**

Age days	Light intensities	
	Lux	Foot candles
0 - 3	20 - 30	2.0 - 3.0
4 - 10	20	2.0
11 - 112	15 - 20	1.5 - 2.0
113 - end of lay	5 - 10	0.50 - 1.0

**LIGHT INTENSITY**

Once accustomed to a particular level of light intensity, a flock will react to any noticeable change. More brightness will increase activity, may stimulate feed intake and will increase the chance of cannibalism. Under bright lights, birds in strange surroundings find water and feed more quickly than with dim lights.

Decreasing brightness will make a flock calmer, although the bird’s initial response to a sudden large reduction in light intensity may be to cease activity and sleep. This happens even when the light level is theoretically adequate. Minimum light intensity is usually put at 10 lux (1.0-foot candle), but may, under some circumstances, be as low as 5.0 lux (0.5-foot candle).

DIMMERS

Dimmers on the light circuits give some advantages. Day to day management may require more or less light at short notice for such jobs as vaccinating, catching, beak trimming and routine inspections. None the less, to avoid possible interruptions in eating or laying habits, long term changes in light intensity should be made slowly. Small daily adjustments over a 7 day period can achieve a change without any negative reaction from the flock.



# THE PRE-LAY AND LAYING PERIODS

## LAYING HOUSE REQUIREMENTS

Adult optimum space requirements are presented in Table 15. In hot areas, allow 20% more space per bird, particularly when ventilation is marginal.

TABLE 15

### ADULT SPACE REQUIREMENT

	Cages	Litter	Litter & Slats
Floor	450 cm <sup>2</sup> (69.8 sq.in.) / bird	6 birds/m <sup>2</sup> 1.8 sq.ft./bird	8 birds/m <sup>2</sup> 1.3 sq.ft./bird
Feeder    Trough Pans	10 cm (4") / bird -	7.5 cm (3") / bird 4/100 birds	
Drinker    Trough Birds/Round Bell Drinker	10 cm (4") / bird -	5.0 cm (2") / bird 90	
Cups or Nipples	maximum 10 birds/cup or nipple a minimum of 2 cups or nipples per cage	maximum 10 birds/cup or nipple	

#### NOTES:

1. With all types of equipment, do not exceed the manufacturer's recommendations for minimum floor, feeder and drinker space.
2. These recommendations reflect general practice. In some countries, legislation dictates greater minimum space allowances. Comply with the law.
3. In some countries, floor, feeder and drinker allowances are less than required for optimum performance. Under these conditions, lower performance is expected and accepted. Bovans birds perform competitively under all circumstances.

## FEEDING DURING PRE-LAYING PERIOD

Bovans White layers should be full fed for maximum production.

During the 2-3 weeks prior to first egg, the liver and reproductive systems increase in size in preparation for egg production. At this time, calcium reserves are built up.

Layer I ration should be given to the birds as soon as the first secondary signs of sexual maturity appear (combs, wattles). Ideally, at least one week before expected first egg.

In order to avoid feeding a costly layer ration when the birds are still not laying, some programs utilize a pre-lay ration to replace the 17-18% crude protein grower. A pre-lay ration is similar to a laying ration, except for 2.0-2.5% total calcium. Furthermore, the linoleic acid content of a pre-lay ration does not need to be over 1.0%. Just as with the layer, the pre-lay Energy level must be higher than that of the grower ration. Pre-lay ration can only be fed until first egg is

reached, and never after. Failure to feed laying hens with a complete Layer ration may result in less than optimum shell quality later in production.

To obtain the best results, pullets should be housed no later than 16 weeks of age, or one week prior to light stimulation.

## FEEDING AT ONSET OF PRODUCTION

At first egg, birds must be already on a layer ration (Table 4). Recommended vitamin and trace mineral levels are found in Table 3.

In certain circumstances, the flock might not reach 100 g/ bird (22 lb/100) daily feed intake before peak production. In such cases, a high nutrient density layer ration may be required to ensure the birds receive the required nutrients for sustained production and early egg size increase. It is also recommended to stimulate feed intake, through suggestions on page 12.



## PROTEIN

Satisfying the crude protein requirement is no assurance that the bird will attain its genetic potential. The daily intake of essential amino acids in the proper proportions is essential if maximum performance is to be obtained. Specific amino acid daily intake recommendations are shown in Appendix No.2.

TABLE 16

### BODY WEIGHT IN THE ADULT PERIOD

Age in Weeks	Grams	Pounds
19	1320	2.91
20	1380	3.04
21	1450	3.20
22	1510	3.33
23	1550	3.42
24	1570	3.46
25	1585	3.49
26	1595	3.52
27	1605	3.49
28	1615	3.54
29	1622	3.58
30	1628	3.59
40	1660	3.66
50	1672	3.69
65	1690	3.73
75	1702	3.75
90	1720	3.79

## ENERGY

Energy intake of the hen is often more limiting than protein or amino acid intake. This is especially true during warm periods (25°C (77°F) and above), and also at the onset of production when feed intake is low. The energy level, as well as density of all other nutrients in the ration, should be adjusted in accordance with actual intake of feed. Appendix 2 can be used to determine correct nutrient densities based on actual feed consumption.

## BODY WEIGHT, EGG SIZE AND PRODUCTION INCREASE

During the 10 weeks after the flock reaches 5% daily egg production, weekly egg weights and body weights should be taken (see Appendix 3 and Table 16). The aim is for weekly increases in body weight and egg weight. Body weight should increase by 300 grams (0.66 lb.) and egg weight by 14.5 grams (11.5 lb. per 30 dozen case) during this 10-week period. If the flock fails to increase body weight in any one week, egg weight may also soon fail to increase. Loss in egg numbers could follow if effective remedial action is not taken promptly.

Low gain in egg and body weight is usually indicative of sub-optimal nutrient intake. Stimulate feed intake as necessary (see "Some Ways to Stimulate Feed Intake", page 12) and consult a nutritionist for ration changes. Disease conditions can adversely influence nutrient absorption. Check with a veterinarian or poultry pathologist when necessary.

### ADJUSTED DENSITY FEEDING

Market conditions vary throughout the world. Premiums are paid for eggs of a certain size in some markets, while in others they are sold by weight. The "ideal size" from an economic point of view varies with market area.

Egg weight tends to increase throughout the life of a flock. This trend can be modified by regulating nutrient intake, as well as modifications in lighting and other management techniques.

Several formulae for the laying period are given in Table 4. Energy densities between 2820 kcal/kg or 11.80 MJ/kg (1280 kcal/lb.) and 2900 kcal/kg or 12.13 MJ/kg (1320 kcal/lb.) are suitable. The change from one ration to another should be based not only on percent production, but also on egg mass output. Page 6 illustrates how this output is calculated. Do not shift from Layer I to II before daily egg mass starts declining (usually at about 56.3 g). The change from Layer II to III should not be made before daily egg mass is down to 54 g. Change from Layer III to IV before daily egg mass goes below 50 g.

The formulae suggested differ in the level of protein and other nutrients. The protein must provide all required amino acids in the proper balance. Make sure that the recommended amino acid levels are met.

Decide which formula should be used, based on the actual feed intake and on daily protein requirements (see Appendix 2).





Protein and amino acid requirements are greatest from the onset of production up to peak egg mass. This is the period when body weight, egg weight and egg production are all increasing.

While production is over 85%, feed 19-20 g of protein/bird/day.

The Adjusted Density Feeding Program is designed to provide adequate levels of all nutrients early in the laying cycle to encourage good early egg size.

## POST PEAK PROTEIN AND ENERGY

After the peaks in both egg numbers and daily egg mass have been reached, daily nutrient requirements are lower. The protein/energy level in the ration can be adjusted to reflect this lower requirement.

Do not decrease daily protein intake in any one week by more than 0.5 g/bird/day and energy by 50 kcal/kg (23 kcal/lb). Intervals between subsequent reductions must be at least 3-4 weeks.

## POST PEAK BODY WEIGHTS

From 36 weeks of age to the end of lay, body weight should remain relatively constant with only a slight gain (Table 16). Investigate any decrease in body weight. The cause, if not corrected, may lead to reduced egg production.

## EGG SIZE

Egg size may be manipulated by adjusting nutrient intake or the lighting program. If satisfactory egg size is not obtained, check your critical amino acid levels, particularly that of methionine. If this is not adequate, correct it by either increasing protein intake, and/or adding a feed grade form of methionine. The level of linoleic acid should also be checked to make sure that it is over 1.3%.

Small egg size can be caused by low energy intake, as well as low protein intake which, in turn, may be caused by extremes in temperature, excessive feed control and/or inadequate feeder space.

Egg size may be increased by switching to a 3 hour on and 3 hour off lighting program. A small loss in egg numbers may be experienced at this time, but egg size is increased and shell quality and colour are also improved. Returning to a regular lighting program will return egg size to normal. This program should only be used in light-tight housing.

It must be noted that limiting protein/energy in an attempt to reduce egg size may adversely affect egg numbers. Changing the protein/energy level to control egg size must be done carefully and slowly.

## TEMPERATURE AND VENTILATION

For optimum bird health and performance, laying house temperatures should be between 21-24.5°C (70-76°F). This is the range within which normal metabolic heat production is balanced by heat loss. However, in temperate climates, laying houses are often kept at 24-28°C (75-82°F) for maximum feed efficiency.

## CALCIUM

Calcium is one of the most critical nutrients for laying hens. Calcium absorption is more efficient when it is provided both as ground limestone and granular limestone or marine shells. From the commencement of lay to 40 weeks of age, it is recommended that two thirds of the total calcium be provided as ground limestone powder and one third in a granular form.

After 40 weeks of age, the amount of granular calcium should be increased so that it represents two thirds of the calcium, while the powder is one third.

The efficiency of calcium absorption may become progressively lower after 40 weeks of age. In addition, the increase in egg size raises the amount of calcium required for a strong shell, thus a higher daily intake is necessary.

Recommended daily calcium allowances are shown in Appendix 2. Appendix 2 can be used to determine the calcium level required in the feed to meet the recommended calcium allowance.

## PHOSPHORUS

Phosphorus requirements vary slightly during the laying period from about 0.45% of Available Phosphorus early in the production cycle to 0.37% at the end. Do not feed over 0.5 g of Available Phosphorus per bird per day. Excess phosphorus has been shown to be detrimental to egg shell quality.



# PULLET AND HEN HEALTH

Bird health results from the interface between adequate biosecurity, animal welfare, poultry husbandry, nutrition, immunization, and general disease prevention, control and monitoring. This section is intended to serve as a general guide for maintaining healthy flocks. Details on actual procedures should be provided by a poultry health professional.

## BIOSECURITY

The concept of biosecurity may be complex and difficult to generalize or adapt to every possible circumstance. However, the important notion and objective of biosecurity is to prevent infectious disease from affecting otherwise healthy flocks. Among the many strategies used to prevent infectious disease, some of the most effective ones include: a) limit access of unnecessary visitors; b) avoiding visits to multiple farms in the same day; c) shower in and out of any poultry facility; d) even when showering is not possible, it is imperative to wear clean clothes or coveralls, footwear and hairnets that should not leave the farm being visited; e) establish, maintain and monitor adequate programs for rodent and insect control. In addition, it is important to consider all major risks in terms of biosecurity such as moving birds into and from the farm; sales, maintenance, equipment and construction personnel; manure removal personnel, vehicles and equipment. Service personnel should not visit any flocks after having been in contact with flocks with known, suspect or obvious signs of disease caused by agents such as MG, MS, ILT or IBV.

## WELFARE AND POULTRY HUSBANDRY

Overall bird health is relatively easy to maintain by simply applying good husbandry. The health and productivity of chickens is closely related to their welfare, which in turn depends on the use of adequate biosecurity and husbandry practices. In many areas, official regulations dictate specific requirements related to animal welfare and it is important to ensure compliance with regulatory agencies. Local or national poultry associations and Government institutions are usually a good source of welfare guidelines that are relevant for each geographical area.

## DISEASE PREVENTION BY IMMUNIZATION

An individual becomes “immune”, “immunized” or resistant to a specific disease after inoculation of a specific vaccine, or after exposure to a disease agent in the field. Vaccination programs should be designed to “immunize” flocks against diseases of economic importance; and against disease agents that could potentially compromise food safety. The entire disease control program relies on sound and well-designed vaccination programs and adequate biosecurity, husbandry and nutrition. At the same time, vaccinations should be administered at times or ages when their detrimental impact should be minimal, and at times when the best possible benefit can be obtained from them.

Most vaccination programs are intended to immunize chickens against diseases that affect the immune system; cause tumors in chickens; affect the respiratory, urinary or reproductive tracts; affect the nervous system; induce disease in the intestinal tract; or represent a food safety concern. Fortunately there are vaccines and vaccination methods available to protect chickens against most of these groups of conditions.

Prior to using any vaccines, ensure that their use is legal and that it will not disqualify specialty flocks because of the type of preservatives contained in the vaccines.

## TYPES OF VACCINES

There are many types of vaccines available for commercial poultry. It is important to become familiar with their basic characteristics related to their potential for protection, safety, ease of administration, relative cost, reactivity, compatibility with other vaccines, etc. Following is a list of some of the most important types of vaccines:

- Live virus vaccines
- Recombinant virus vaccines
- Live bacterial vaccines
- Inactivated bacterial vaccines (bacterins)
- Gene modified and deletion mutant live bacterial vaccines
- Autogenous inactivated bacterial vaccines
- Autogenous inactivated viral vaccines
- Live coccidiosis vaccines
- Live Mycoplasma vaccines
- Inactivated Mycoplasma vaccines (bacterins)
- Recombinant Mycoplasma vaccines
- Competitive exclusion products



## VACCINATION METHODS

It is important to know the characteristics of each vaccine and to use each product according to the manufacturer's recommendations. Vaccines are designed and approved for individual or mass application methods.

Individual vaccination methods include:

- Ocular (eye drop)
- Beak dipping or intranasal
- Subcutaneous injection
- Intramuscular injection
- Transcutaneous injection (wing web)
- Vent brush application

Mass vaccination methods include:

- In ovo injection
- Drinking water vaccination
- Spray vaccination

### OCULAR (EYE DROP), BEAK DIPPING AND INTRANASAL VACCINATION

Eye drop vaccination is commonly used to protect chickens against respiratory viruses, *Mycoplasma* and occasionally against infectious bursal disease. Ocular vaccination is most suitable for delivery of live vaccines against diseases or agents such as (but not exclusively) Newcastle disease, infectious bronchitis, infectious laryngotracheitis, avian metapneumovirus and *Mycoplasma gallisepticum*. Eye drop vaccination is likely the most effective and safest method for respiratory viruses. Direct contact of the vaccine with the mucosa of the eye will result in stimulation of the Harderian gland and a strong local immune response. Despite being highly effective, eye drop vaccination is labor intensive and time consuming and thus it is usually limited to application of vaccines that must be administered via the ocular route and by no other method, such as some (but not all) live MG vaccines and live attenuated vaccines against ILT. Intranasal and beak dipping application of vaccines has the same objectives as with the ocular route. Intranasal application is popular in some countries but beak dipping is rarely used. The vaccine is administered by depositing a drop (usually 30  $\mu$ l or 0.03 ml) of reconstituted vaccine directly on the eye or into the nostrils. The advantage of eye drop application is that if applied properly, every bird receives a similar dose of vaccine and is thus likely to be immunized (protected) against the disease, as opposed to mass application methods, which unavoidably result in suboptimal coverage since not every bird received an equally immunizing dose. Because eye drop vaccination requires individual handling of birds, biosecurity is most important and the vaccination crews must follow strict biosecurity procedures not to bring infectious diseases to the flock being vaccinated. For the beak dipping method

to be successful, both nostrils must be immersed in the vaccine. This method is suitable only for chicks up to 7 days of age and is used for immunization against NDV or IBDV. It is used in areas or farms where even vaccine uptake is not possible using the drinking water or spray methods, or with the objective of minimizing vaccine reactions.

### SUBCUTANEOUS AND INTRAMUSCULAR INJECTION

Injection via the intramuscular and subcutaneous routes is reserved primarily for inactivated vaccines and bacterins. The vaccination equipment should be sterile and the needles used should be of the proper caliber and length for the age of the bird and also for the type of product being injected. The needles should be replaced with sterile needles at least every 500 injections to prevent infections with bent or blunt needles, and to avoid transmission of some diseases from infected to non-infected chickens. Most inactivated (killed) vaccines are administered at approximately 12-14 weeks of age. Should it be necessary to vaccinate younger chickens with inactivated products it should be kept in mind that handling and administration of inactivated vaccines or bacterins between 6 and 11 weeks of age might delay or alter the development of the pullets. Inactivated viral vaccines are usually available in a water-in-oil (WO) or water-in-oil-in-water (WOW) emulsion, which are typically not very reactive. Thus, such products can be injected with confidence intramuscularly or subcutaneously, provided the injection is done in the proper site and without depositing any of the vaccine product in the cavity or directly into the internal organs. Inactivated products containing *Mycoplasmas* and/or bacteria such as *Salmonella* may be quite reactive and every effort should be made to minimize the local vaccine reactions that can be derived from the injections. For subcutaneous injections, it is especially important to avoid the thymus by injecting the vaccine in the middle line (avoiding the sides of the neck), and by not injecting too close to the head or the base of the neck. For intramuscular injections (in the breast muscle), every effort should be made to avoid injecting the product into the cavity. Vaccinations in the thigh may contribute to reduce adverse reactions but care must be exercised to minimize injuries resulting in lameness.

### TRANSCUTANEOUS INJECTION (WING WEB)

Transcutaneous (wing web) application is used almost exclusively to vaccinate chickens against poxvirus (POX). For convenience, manufacturers of vaccines have added other agents such as chicken infectious anemia virus (CAV) and avian encephalomyelitis virus (AE) to POX vaccines and thus it is possible to vaccinate pullets simultaneously against AE, POX and CAV in one injection. The latter (CAV)





is only necessary in layer breeders but AE and POX are routinely used in commercial layers. In addition, there are recombinant vaccines with a poxvirus as a vector carrying genes that express proteins from ILTV or MG. Thus, such products can also be administered by wing web application.

## VENT BRUSH VACCINATION

Vent brush vaccination was developed decades ago to protect chickens against ILTV using vaccine strains that were extremely reactive and caused vaccine-induced ILT. The procedure involves dipping a rough brush into the reconstituted ILTV vaccine vial and brushing harshly the mucosa of the vent. This procedure is still used with relative success in some countries for administration of live attenuated vaccines against ILTV.

## IN OVO INJECTION

In ovo vaccination is a mass-application procedure that is reserved for vaccination of embryos in the hatchery and is typically done at 17 to 19 days of incubation. The procedure was designed for immunization against Marek's disease virus (MDV). With the advent of recombinant vaccines, in ovo vaccination can now be used to protect chickens against diseases such as Marek's disease, fowl poxvirus, infectious laryngotracheitis, infectious bursal diseases (Gumboro) and Newcastle disease. In addition, coccidiosis vaccines are now registered and approved for in ovo administration.

## DRINKING WATER (ORAL) VACCINATION

Vaccination via the drinking water is a suitable method to vaccinate pullets against hardy viruses such as infectious bursal disease virus (IBDV) and CIAV, but it can be used to immunize against diseases such as Newcastle, infectious bronchitis, colibacillosis, salmonellosis and other diseases. Along with spray vaccination and in ovo vaccination, administration of live vaccines via the drinking water is considered a mass-application method. Although practical, mass application methods usually result in less-than-optimal vaccine coverage and thus protection might be suboptimal compared with individual vaccination methods. Vaccination via the drinking water should be used in birds one week old or older because water consumption in younger pullets might be too irregular. Oral vaccination can be done by directly adding the vaccine into the water reservoirs supplying water to the barn to be vaccinated; it can also be accomplished by using "medicators" or "dosifiers" that can be connected to the main water pipelines feeding the drinkers.

The method relies on the preparation of a stock solution of vaccine that is to be placed in a container (a clean bucket) from which the medicator draws small quantities of vaccine that is mixed automatically with fresh incoming water in the

water pipelines. For example, 1 ounce of stock solution of vaccine (approx. 28.5 ml) is drawn by the medicator and mixed with every 1 gallon (approx. 3.78 liters) of fresh water to be consumed. This method requires that the birds to be vaccinated be thirsty so that water consumption helps to consume the vaccine rapidly (within approximately 60 minutes). Thus, access to water by the birds should be interrupted for approximately 2 hours (depending on the temperature, humidity, age of the birds, etc.) prior to vaccination. Because the stock solution is drawn in a pulse manner, and because it must be mixed with incoming water automatically before it is delivered it is not possible to achieve an even vaccination in all birds.

## VACCINATION THROUGH A MEDICATOR

Vaccination through a medicator is one of the methods of vaccination with live virus vaccines which is least recommended, albeit it is a suitable method for administration of drugs, vitamins, etc. Coccidiosis vaccination using a medicator should be avoided because the Eimeria oocysts will tend to settle and the actual dose of oocysts per bird will vary greatly, and so giving very poor results.

## WATER VACCINATION

Water vaccination can also be accomplished using a water pump to "inject" or "force" the vaccine into the water lines, which is a popular and very effective method of mass application using the drinking water for delivery of live vaccines. Water pump vaccination requires a closed water system (nipple drinker lines) and can be used successfully for delivery of vaccines against diseases or disease agents such as IBDV or CIAV. As with other methods involving water delivery, this one requires that the birds be thirsty prior to delivering the vaccine to them. Wherever possible, the drinker lines are raised high enough so as to prevent drinking by the birds in the 2-3 hours prior to vaccination.

Water vaccination requires flushing the drinker lines with fresh water to minimize the amount of unwanted residues. Commercial products can be used to clean the drinker lines thoroughly prior to vaccination. Even after the use of commercial products, it is recommended to flush the lines with clean fresh water before vaccinating the flock. This is particularly important in operations that have hard water, or in operations that have used antibacterial drugs or other products that may have formed a film in the drinker lines. Prior to vaccination, it is important to water-starve the pullets to be vaccinated such that most will consume vaccine upon its administration. Check the drinkers or nipple drinkers to ensure they are clean and operational and shut down all water sanitizing systems. Allow the birds to become thirsty by interrupting their access to water. The amount of time



required for the birds to become thirsty will depend on their age, environmental temperature, feed formulation, etc. The goal should be for all pullets to consume the vaccine in a matter of 60 minutes. If the birds consume the vaccine in less time, it would mean they were too thirsty. On the other hand, if it takes the birds more than one hour to fully consume the vaccine this would be an indication that the water was not removed long enough prior to vaccination.

A few essential steps for water vaccination are listed as follows:

- Clean and flush the water lines.
- Turn off the water sanitation system.
- Ensure proper functioning of the drinker system.
- Water-starve the pullets enough for them to consume the vaccine in less than one hour.
- Verify that the vaccine to be administered has been stored according to the manufacturer's recommendations; that it is still viable (before expiration); and maintain a record of the type of vaccine, serial (lot) number, number of doses per vial and number of vials used, as well as the expiration date.
- Reconstitute the vaccine in an aseptic manner and verify that the number of vials used matches the number of doses to be given. The amount of vaccine to be consumed in volume should be equivalent to approximately 1/7 the total water consumed the previous day.
- Use a commercially produced vaccine stabilizer or powdered skim milk to help protect the vaccine viruses. Follow closely the recommendations of the manufacturer of the vaccine stabilizer. If skimmed milk is used, approximately 2.5 g of well-dissolved skim milk per liter of water plus vaccine is enough to protect the vaccine from any residual chemicals or minerals in the drinking water. Keep the reconstituted vaccine cool and away from exposure to the sunlight.
- Deliver the vaccine into the drinkers and drinker lines. To ensure a complete fill out of the drinker lines (pipes) add a visual aid such as commercial blue dye and let the vaccine be flushed to the end of the lines until blue dye is seen at the end of the lines. At this time close the end of the water lines and allow the birds to drink. If the vaccine is delivered into open water systems, it is important to walk slowly through the house to stimulate water consumption and to help distribute the birds in the house.
- Check at least 100 birds throughout the barn to verify that they have consumed the vaccine. If enough dye was used, it should be easy to observe a blue coloring of the tongue, head feathers and occasionally the crop, which is visible through the skin. Vaccine coverage of at least 90% should be a realistic goal.

## SPRAY VACCINATION

Spray vaccination is used primarily for immunization against respiratory viruses such as Newcastle disease virus (NDV) and infectious bronchitis virus (IBV). However, it should be noted that spray vaccination should involve the less invasive forms or strains of viruses, such as the B1B1 strain of Newcastle, or H120 of infectious bronchitis. In general, the more invasive the virus, the better the protection against disease but the harsher the vaccine reactions, especially in flocks infected with Mg or some strains of MS. Coccidiosis vaccines are sometimes sprayed on the feed of layer breeders in some areas. Some live *Mycoplasma gallisepticum* vaccines (but not all) can be sprayed directly on chickens in the field. Each type of equipment intended for spray vaccinations may be different and the operator must be thoroughly familiar with each piece of equipment and its spray patterns, pressure and particle size. For example, pressurized sprayers are excellent to deliver vaccine to the respiratory tract but because of the small particle size that they produce the vaccine will tend to remain suspended in the air or it may be sucked towards the house fans if they are not turned off prior to vaccinating the birds. With some types of sprayers the equipment must be located not more than 50 cm over those birds to be vaccinated. This method is therefore not practical for mass application over chickens on the ground. Rather, sprayers intended for horticultural use or pesticide application in the horticultural industry have proved very popular and effective for application of live respiratory vaccines in the field. The particle size will range between 100 and 300 µm, which is suitable for most respiratory viruses. In general, spray vaccination is used for protection against respiratory viruses and *Mycoplasma* in pullets; and for protection against respiratory viruses in hens in production.

A few essential considerations for spray vaccination are listed as follows:

- Prior to choosing spray vaccination to immunize chickens against respiratory viral diseases, consider all possible options. Be aware that spray vaccination against Newcastle disease and Infectious Bronchitis generally provides better protection than water vaccination, but vaccine reactions can be harsh, particularly in *Mycoplasma*-positive chickens. Spray vaccination against Infectious Laryngotracheitis should be avoided and must never be done in chickens in production. Vaccinate only healthy chickens.
- For adult flocks, verify the flock antibody titers prior to vaccination. If antibody titers are low, vaccine reactions may be harsh.
- Ensure that the vaccination equipment has been thoroughly cleaned, disinfected and rinsed to remove all traces of vaccine and disinfectant.



- Drive the birds (if reared on the floor) to an area of the barn where they can be vaccinated without them flying or moving freely away from the vaccination equipment.
- Calculate the total number of doses and the total volume of diluent (distilled deionized water) required to vaccinate all chickens. The water used should not be chlorinated and should have a pH of 5.5 to 7.0.
- Turn off the lights, brooders and ventilation system while ensuring the birds do not overheat or suffocate. The flock should be relatively calm at the moment of spraying the vaccine on them.
- Reconstitute the vaccine aseptically and in the shade, and only immediately prior to vaccinating the flock.
- Use appropriate personal protective equipment (PPE), including protective mask and goggles.
- Adjust the spray nozzle to a proper droplet size. Coarse sprays (>80-120 microns) are recommended for priming vaccinations and also for invasive vaccines. Fine sprays (50-60 microns) are recommended for boost vaccinations in older chickens, but only after they have been primed with similar viruses.
- MG-infected chickens tend to react too severely to spray vaccinations, particularly if the droplet size is too small.
- Use distilled water to dilute the vaccine (the amount should be adjusted to every situation). If a pressurized spray apparatus is used, it should be kept in mind that this type of equipment delivers droplets with a diameter range of 50-1000 microns, and thus only part of the vaccine will be inhaled. Thus, it is necessary to spray the vaccine at a distance not larger than 50 cm from the chickens. This type of equipment typically requires a relatively large volume per chicken house (15-20 liters). For situations where a controlled-droplet application apparatus is used, the droplet size is considerably more uniform (~50-150 microns). Although the droplet size is more uniform with this type of equipment, some of the droplets are too small and may remain in suspension for quite some time after the vaccine is sprayed. This may represent a problem because a vaccine that stays in suspension a long time may decrease in virus titer before it is inhaled and much of the vaccine ends up on house and equipment surfaces but not in the chickens. In addition, if much of the vaccine remains in suspension (in the form of a mist), re-activating the ventilation system will draw the vaccine out of the house through the exhaust fans.
- Spray-vaccinate only healthy birds. Avoid spraying birds that are infected with MG.
- Adjust the nozzle to obtain the desired droplet size.
- Wear a mask and goggles for personal protection when spray-vaccinating.
- Make sure the sprayer to be used is clean and has no residual disinfectant. The vaccine containers of the spray apparatus should be rinsed with distilled water prior and after every use.
- Use only one dose per bird or less.
- Reconstitute the vaccine only immediately prior to use.
- Close up the house including curtains and doors and shut the ventilation system and dim the lights while the birds are being vaccinated and if possible, during 20-30 minutes post-vaccination (provided the air quality and temperature allow for a temporary shut down without compromising the flock integrity). If the flock is in a high temperature area, vaccinate birds at night or early in the morning. Make sure the ventilation system is not running at the time the vaccine is being applied or that it runs at a minimum power. Dim the lights to a minimum to settle the birds.
- Spray the birds evenly and thoroughly at least twice and ensure that all calculated doses are used evenly. The heads and upper body of the sprayed birds should appear wet after vaccination.
- Make a point about not leaving the farm without making sure the ventilation system and the lights have been re-engaged. Ventilation should be restored approximately 20 minutes after the initiation of the vaccination process.
- Rinse, clean, disinfect and rinse again the vaccination equipment before leaving the farm.
- Destroy all residual vaccine and vaccine vials by incineration. Follow local regulations regarding adequate disposal of vaccines, vaccine vials and biological materials.

## PARASITE CONTROL

The most common internal parasites in laying hens include coccidia, Histomonas ("black head"), Capillaria worms, round (Ascaridia) worms, and cecal (Heterakis) worms. External parasites frequently seen in layer operations include the Northern fowl mite (Ornithonyssus), the red mite or roost mite (Dermanyssus) and poultry lice are less frequent but can occur in commercial operations. The Northern fowl mite completes its entire life cycle on the birds, whereas the red mite feeds on the birds only at night. As a whole, mites are external parasites that must be controlled to avoid drops in egg production, dermatitis around the vent, restless birds, increased mortality and farm employee discomfort. Some mites are known to carry other disease agents and can induce anemia if the infestation is severe and thus must be controlled.

## VACCINATION AGAINST COCCIDIOSIS

Pullets reared in battery cages do not ordinarily experience significant internal parasitic diseases. However, if they have access to droppings in the hen house because of the type of equipment and manure removal systems, outbreaks of coccidiosis could potentially occur. Regardless of the type of operation, it is important to ensure immunity against coccidiosis, which can be accomplished by using one of two





common methods. Where legal, pullets reared on the floor may be treated with anticoccidial drugs for 8-12 weeks to allow for a gradual acquisition of immunity. Commonly used drugs for this purpose include (not exclusively) amprolium and salinomycin. However, other anticoccidial drugs have been used successfully. Perhaps the best approach to control coccidiosis in pullets reared on the floor is vaccination. Pullets can be vaccinated by spray at the hatchery with one of the various commercially available vaccines. It is important to use a commercial product that will contain at least *E. acervulina*, *E. maxima*, *E. tenella* and *E. necatrix*. Coccidiosis vaccines for broiler chickens do not contain *E. necatrix*, an essential component of coccidiosis vaccines for long-lived birds. When coccidiosis vaccines are used it is critical not to medicate the flock with any drug that coccidia would be sensitive to in order to allow at least two complete coccidial cycles, which normally occurs at approximately 14-16 days of age, depending on various factors including litter moisture, bird density, environmental temperature, etc. It is also important to allow vaccinated birds to remain in the brood chamber for the duration of at least 2 complete coccidial cycles before allowing them to occupy the entire barn. If vaccinated pullets are given the entire barn prior to the second cycle being completed, many of them will not be properly immunized and might develop coccidiosis at a later age, with the significant consequences of increased mortality, delayed growth, poor uniformity and the need for treating the flock. Specialty flocks may not be treated with drugs of any kind and thus it is critical to ensure proper coccidiosis control with the use of vaccines.

## HISTOMONIASIS AND ROUND WORMS

*Histomonas meleagridis* (HM) is the causative agent of histomoniasis ("black head") and affects almost exclusively pullets reared on the floor, particularly in premises with dirt floors. The condition, which can be devastating, has made a comeback after the ban of many anti-parasitic drugs and is difficult to control since there are no drugs that can be used legally in many areas of the world. Because the microscopic parasite depends to some extent on the life cycle of cecal worms and earthworms, one of the strategies for control involves the control of worms. Early administration of drugs against round worms might contribute to maintain HM under control. The layer industry uses most commonly piperazine and anti-worm compounds in the family of the benzimidazole drugs such as levamisole or albendazole. Controlling worms reduces the challenge posed by HM. Where legal, HM infection may be treated with drugs such as nitarsone, but even this drug is only partially effective. Control of HM involves not only treating birds against worms, but also proper cleaning and disinfection, adequate husbandry, and proper coccidiosis control, particularly of *E. tenella*.

## MITES

The best form of prevention for mite infestation is biosecurity. Infested flocks should never be visited before visiting mite-free flocks. Mites can be mechanically carried from farm to farm in clothes, footwear, on people, equipment, egg boxes, etc. Such parasites usually thrive in sexually mature flocks and thus most treatments become necessary while the infested flocks are in production. Effective control requires direct application of "acaricide" products. A variety of products can be used for mite control, including pyrethroids, organophosphates, carbamates, mineral-based products, vegetable oils, citrus concentrated extracts and other products. Some of such products can be administered by dry (dust) spray, or as a wet spray. Prior to using any of these products it is critical to determine whether they are approved for use in hens in production, and also whether the personnel applying the products requires personal protective equipment. Some products are poorly effective if applied dry but quite effective when applied wet directly on the birds, which requires considerable more time than the application of dry products. Treating birds against mites is frequently expensive and it may be necessary to treat an infested flock more than once. After the affected flock is removed, thorough cleaning and disinfection and chemical treatment of the premises and equipment is necessary. The best approach is to exercise adequate biosecurity and never to transit from infested flocks to clean flocks, or to share equipment and egg boxes between infested and clean premises.

## CONTROLLING GROUPS OF DISEASES BY VACCINATION

Infectious diseases can be grouped by the organ system they affect. Thus, infectious diseases can affect the respiratory, digestive, nervous, urinary, reproductive and immune systems among others. Other diseases tend to affect the integument (skin or cutaneous tissues) and yet some others are considered a concern for food safety.

## RESPIRATORY DISEASES

Respiratory diseases of major concern in commercial layers include Newcastle disease, infectious bronchitis, avian influenza, avian metapneumovirus infection (swollen head syndrome), avian mycoplasmosis (MG and MS), infectious coryza, avian pasteurellosis (fowl cholera) and *Gallibacterium anatis* infection. All such diseases or disease agents can be prevented or controlled by using a combination of biosecurity and vaccination. In general, vaccination against respiratory viruses is done with live vaccines followed by killed (inactivated) vaccines. Live attenuated avian influenza



vaccines are not available, but recombinant vaccines and killed vaccines are. Bacterial diseases (infectious coryza, fowl cholera and Gallibacterium infection) are typically prevented by means of inactivated (killed) vaccines or bacterins, which are given once or twice during the rearing period. Bacterins are usually administered by intramuscular or subcutaneous injection at approximately 10-14 weeks of age. Live vaccines against viral respiratory diseases may be administered by spray or in the drinking water once or multiple times while the flocks are in production.

## PERITONITIS IN LAYERS

Peritonitis in layers is frequently caused by E. coli strains that are unrelated to E. coli strains affecting cattle or humans. However, they can induce severe economic losses if there is no adequate control. Colibacillosis associated with peritonitis in layers is not strictly a respiratory condition, but E. coli can penetrate via the respiratory tract (descending infection). E. coli can also penetrate via an ascending route (via the reproductive tract), or possibly from the intestinal tract, a mechanism that has not yet been confirmed. Peritonitis in layers should be controlled by a variety of approaches, including maintaining proper husbandry practices, adequate ventilation, and vaccination against E. coli among other strategies. Vaccination against E. coli in layers is a very effective method of control and is commonly done by using live vaccines by spray or in the drinking water twice during rearing, once at hatch and once a few weeks later. Live E. coli vaccines can also be given safely in flocks in production or soon before onset of production if they were not vaccinated during rearing.

## DISEASES OF THE DIGESTIVE SYSTEM

Diseases of the digestive system that are preventable by vaccination include the parasitic disease coccidiosis. Coccidiosis vaccines are typically administered at the hatchery in ovo or by spray, or by spray on the feed during the first week of life.

## DISEASES AFFECTING THE NERVOUS SYSTEM

Diseases affecting the nervous system such as avian encephalomyelitis (AE) require effective vaccination for prevention. Flocks may be vaccinated via the drinking water or by transcutaneous injection in the wing web, usually along with POX vaccination at approximately 10-12 weeks of age. AE vaccines should not be given for the first time before 10 weeks of age or too soon before the flock initiates egg production because they can induce disease or drops in egg production.

## DISEASES AFFECTING THE URINARY AND REPRODUCTIVE TRACTS

Diseases affecting the urinary and reproductive tracts are represented typically by infectious bronchitis. Prevention of infectious bronchitis requires vaccination at various ages with the same or similar serotypes of virus circulating in the field. It may be necessary to vaccinate 3-4 times the pullets with live viruses during rearing and once with a killed vaccine containing at least the same or similar serotypes circulating in the field. Still, in many instances it might be necessary to vaccinate flocks in production by spray several times in order to maintain a healthy urinary, respiratory and reproductive tract.

## DISEASES AFFECTING THE IMMUNE SYSTEM

Diseases affecting the immune system can be numerous. Well-known diseases affecting the immune system include infectious bursal disease (IBDV, or Gumboro disease), chicken infectious anemia (CIAV), and Marek's disease (MDV), the latter being a disease that also causes tumors and mortality. IBDV can be prevented by vaccination with live attenuated vaccines, immune complex vaccines, or recombinant vaccines. Live attenuated vaccines are becoming less popular because of the frequent need to give them multiple times in order to control IBDV effectively during the rearing period. Still, they have contributed very positively to the effective control of IBDV in the field, particularly in floor rearing operations. Live attenuated IBDV vaccines are given 3-4 times during the first 8 weeks of age, beginning with an initial application at approximately 14 days of age. It is not necessary to vaccinate commercial layers against CIAV because they are only susceptible to this immunosuppressive agent during the first 3 weeks of life, and the layer parents should provide protection after being exposed and/or vaccinated themselves. All layer pullets should be vaccinated against MDV to prevent losses to mortality, immunosuppression and tumors.

## DISEASE AGENTS OF CONCERN FOR FOOD SAFETY

Salmonella control requires a very complex approach, part of which involves vaccination. Where legal, vaccination against Salmonella is one of the most effective means of control and is usually done with live attenuated or genetically modified vaccines against S. typhimurium, followed by killed vaccines against S. enteritidis or containing other Salmonella serovars that may be residents in a particular area or operation. It is recommended to use two live Salmonella vaccines and at least one killed vaccine containing SE and other serovars to reduce the probability of infection in the field.



# APPENDIX 1

## SUGGESTED DAILY NUTRIENT INTAKE FOR BOVANS WHITE COMMERCIAL PULLETS

### Per caged bird

Age in weeks	M.E.	Protein	Lysine	Methionine	Methionine & Cystine	Tryptophan	Threonine	Calcium	Available Phosphorus
	Kcal	g	mg	mg	mg	mg	mg	mg	mg
1	21	1.5	77	34	57	16	52	74	33
2	38	2.8	143	62	107	30	96	137	61
3	62	4.5	231	101	172	48	155	221	99
4	93	6.4	304	141	234	64	211	336	150
5	107	7.4	352	163	270	74	244	389	174
6	119	8.2	390	180	299	82	271	431	193
7	128	8.6	418	194	321	88	290	462	207
8	133	8.7	437	198	336	92	304	483	216
9	134	9.1	442	206	346	101	317	504	226
10	140	9.4	460	215	360	105	320	525	235
11	146	9.6	478	224	374	109	333	546	244
12	151	9.8	497	232	389	113	340	567	254
13	157	10.1	504	235	392	115	353	616	269
14	160	10.4	510	244	406	119	360	638	278
15	168	10.7	537	256	427	125	366	671	293
16	179	11.4	572	273	455	133	390	715	312
17	204	12.8	618	334	554	156	497	2840	334
18	228	14.2	687	371	616	174	553	3160	367

### Instructions for the use of Appendix 1 and 2

1. The Daily feed requirement is determined by dividing the daily energy requirement by the level of the ration, e.g. assuming a daily requirement of 300 kcal ME and a ration containing 2850 kcal ME / kg:  $300 / 2850 = 0.105$  kg or 105 g/bird/day
2. Add 3.5 kcal ME/bird/day for each 1°C below 21°C.  
Subtract 3.5 kcal ME/bird/day for each 1°C above 21°C.
3. Add 2 kcal ME/bird/day for each 1 g of egg mass above the figures quoted in Appendix 3.  
Subtract 2 kcal ME/bird/day for each 1 g of egg mass below the figures quoted in Appendix 3.
4. Add 6 kcal ME/bird/day for each 50 g of body weight in excess of the mean weight quoted in Table 9 and Table 16.



## APPENDIX 2

### SUGGESTED DAILY NUTRIENT ALLOWANCES FOR BOVANS WHITE COMMERCIAL

Per caged bird

Age in weeks	M.E.	Lysine	Methionine	Methionine & Cystine	Tryptophan	Threonine	Arginine	Calcium	Available Phosphorus
	Kcal	mg	mg	mg	mg	mg	mg	g	mg
19	231	784	384	640	176	560	768	3.3	368
20	246	833	408	680	187	595	816	3.5	391
21	266	902	442	736	202	644	883	3.8	423
22	275	931	456	760	209	665	912	3.9	437
23	283	960	470	784	216	686	941	4.1	451
24	286	970	475	792	218	693	950	4.1	455
25	286	970	475	792	218	693	950	4.1	455
26	286	970	475	792	218	693	950	4.1	455
27	286	970	475	792	218	693	950	4.1	455
28	286	970	475	792	218	693	950	4.1	455
29	286	970	475	792	218	693	950	4.1	455
30	286	970	475	792	218	693	950	4.1	455
31	286	970	475	792	218	693	950	4.1	455
32	286	970	475	792	218	693	950	4.1	455
33	286	970	475	792	218	693	950	4.1	455
34	286	970	475	792	218	693	950	4.1	455
35	286	970	475	792	218	693	950	4.1	455
36	286	970	475	792	218	693	950	4.1	455
37	286	970	475	792	218	693	950	4.1	455
38	286	970	475	792	218	693	950	4.1	455
39	289	949	455	768	192	667	929	4.3	444
40	289	949	455	768	192	667	929	4.3	444
41	289	949	455	768	192	667	929	4.3	444
42	289	949	455	768	192	667	929	4.3	444
43	289	949	455	768	192	667	929	4.3	444
44	289	949	455	768	192	667	929	4.3	444
45	289	949	455	768	192	667	929	4.3	444
46	289	949	455	768	192	667	929	4.3	444
47	289	949	455	768	192	667	929	4.3	444
48	289	949	455	768	192	667	929	4.3	444
49	289	949	455	768	192	667	929	4.3	444
50	289	949	455	768	192	667	929	4.3	444



## SUGGESTED DAILY NUTRIENT ALLOWANCES FOR BOVANS WHITE COMMERCIAL

Per caged bird

Age in weeks	M.E.	Lysine	Methionine	Methionine & Cystine	Tryptophan	Threonine	Arginine	Calcium	Available Phosphorus
	Kcal	mg	mg	mg	mg	mg	mg	g	mg
51	289	949	455	768	192	667	929	4.3	444
52	294	927	433	742	185	649	906	4.5	422
53	294	927	433	742	185	649	906	4.5	422
54	294	927	433	742	185	649	906	4.5	422
55	294	927	433	742	185	649	906	4.5	422
56	294	927	433	742	185	649	906	4.5	422
57	294	927	433	742	185	649	906	4.5	422
58	294	927	433	742	185	649	906	4.5	422
59	294	927	433	742	185	649	906	4.5	422
60	294	927	433	742	185	649	906	4.5	422
61	294	927	433	742	185	649	906	4.5	422
62	294	927	433	742	185	649	906	4.5	422
63	294	927	433	742	185	649	906	4.5	422
64	294	927	433	742	185	649	906	4.5	422
65	293	886	402	700	175	628	865	4.6	402
66	293	886	402	700	175	628	865	4.6	402
67	293	886	402	700	175	628	865	4.6	402
68	293	886	402	700	175	628	865	4.6	402
69	293	886	402	700	175	628	865	4.6	402
70	293	886	402	700	175	628	865	4.6	402
71	293	886	402	700	175	628	865	4.6	402
72	293	886	402	700	175	628	865	4.6	402
73	293	886	402	700	175	628	865	4.6	402
74	293	886	402	700	175	628	865	4.6	402
75	293	886	402	700	175	628	865	4.6	402
76	293	886	402	700	175	628	865	4.6	402
77	293	886	402	700	175	628	865	4.6	402
78	291	845	371	639	165	587	824	4.8	361
79	291	845	371	639	165	587	824	4.8	361
80	291	845	371	639	165	587	824	4.8	361
81-90	291	845	371	639	165	587	824	4.8	361





## APPENDIX 3

### BOVANS WHITE EGG PRODUCTION

Age in weeks	% Hen day production	% Liveability	Cumulative eggs/hen housed	Average egg weight			Daily egg mass	Cumulative egg mass/hen housed			Feed intake		Body weight	
				g/egg	oz/doz	lbs/case		g	kg	lbs	bird/day g	100/day lbs	g	lbs
19	4.0	99.9	0	43.7	18.5	34.7	1.7	0.0	0.0	0.0	80	17.6	1320	2.91
20	40.0	99.8	3	46.5	19.7	36.9	18.6	0.1	0.3	0.3	85	18.7	1380	3.04
21	64.0	99.7	8	49.5	21.0	39.3	31.7	0.4	0.8	0.8	92	20.3	1450	3.20
22	80.0	99.6	13	51.7	21.9	41.0	41.4	0.7	1.4	1.4	95	20.9	1510	3.33
23	87.0	99.6	19	53.7	22.7	42.6	46.7	1.0	2.2	2.2	98	21.6	1550	3.42
24	90.5	99.5	25	54.6	23.1	43.3	49.4	1.3	2.9	2.9	100	22.0	1570	3.46
25	93.0	99.4	32	56.1	23.7	44.5	52.2	1.7	3.7	3.7	101	22.3	1585	3.49
26	94.5	99.3	39	57.2	24.2	45.4	54.1	2.1	4.5	4.5	103	22.7	1595	3.52
27	95.5	99.2	45	58.5	24.8	46.4	55.9	2.4	5.4	5.4	104	22.9	1605	3.54
28	96.0	99.2	52	59.0	25.0	46.8	56.6	2.8	6.3	6.3	105	23.1	1615	3.51
29	96.0	99.1	58	59.5	25.2	47.2	57.1	3.2	7.1	7.1	106	23.4	1622	3.56
30	96.0	99.0	65	59.9	25.4	47.5	57.5	3.6	8.0	8.0	107	23.6	1628	3.59
31	96.0	98.9	72	60.3	25.5	47.9	57.9	4.0	8.9	8.9	107	23.6	1635	3.60
32	95.8	98.8	78	60.6	25.7	48.1	58.1	4.4	9.8	9.8	107	23.6	1640	3.62
33	95.6	98.8	85	60.8	25.7	48.3	58.1	4.8	10.7	10.7	107	23.6	1645	3.63
34	95.4	98.7	92	61.0	25.8	48.4	58.2	5.2	11.6	11.6	108	23.8	1650	3.64
35	95.2	98.6	98	61.1	25.9	48.5	58.2	5.6	12.4	12.4	108	23.8	1652	3.64
36	95.0	98.5	105	61.2	25.9	48.6	58.1	6.0	13.3	13.3	108	23.8	1655	3.65
37	94.8	98.4	111	61.4	26.0	48.7	58.2	6.4	14.2	14.2	108	23.8	1656	3.65
38	94.6	98.4	118	61.5	26.0	48.8	58.2	6.8	15.1	15.1	108	23.8	1657	3.65
39	94.4	98.3	124	61.6	26.1	48.9	58.2	7.2	16.0	16.0	109	24.0	1659	3.66
40	94.1	98.2	131	61.7	26.1	49.0	58.1	7.6	16.9	16.9	109	24.0	1660	3.66
41	93.8	98.1	137	61.8	26.2	49.0	58.0	8.0	17.7	17.7	109	24.0	1661	3.66
42	93.5	98.0	144	62.0	26.2	49.2	58.0	8.4	18.6	18.6	109	24.0	1662	3.66
43	93.2	98.0	150	62.2	26.3	49.4	58.0	8.8	19.5	19.5	109	24.0	1663	3.67
44	92.9	97.9	156	62.4	26.4	49.5	58.0	9.2	20.4	20.4	109	24.0	1665	3.67
45	92.6	97.8	163	62.6	26.5	49.7	58.0	9.6	21.2	21.2	109	24.0	1666	3.67
46	92.3	97.7	169	62.7	26.5	49.8	57.9	10.0	22.1	22.1	109	24.0	1667	3.68
47	92.0	97.6	175	62.8	26.6	49.8	57.8	10.4	23.0	23.0	109	24.0	1668	3.68
48	91.6	97.6	182	62.9	26.6	49.9	57.6	10.8	23.8	23.8	109	24.0	1669	3.68
49	91.2	97.5	188	63.1	26.7	50.1	57.5	11.2	24.7	24.7	109	24.0	1671	3.68
50	90.8	97.4	194	63.2	26.8	50.2	57.4	11.6	25.6	25.6	109	24.0	1672	3.69



## BOVANS WHITE EGG PRODUCTION

Age in weeks	% Hen day production	% Liveability	Cumulative eggs/hen housed	Average egg weight			Daily egg mass	Cumulative egg mass/hen housed		Feed intake		Body weight	
				g/egg	oz/doz	lbs/case	g	kg	lbs	bird/day g	100/day lbs	g	lbs
51	90.4	97.3	200	63.3	26.8	50.2	57.2	12.0	26.4	110	24.3	1673	3.69
52	90.0	97.2	206	63.4	26.8	50.3	57.1	12.4	27.3	110	24.3	1674	3.69
53	89.6	97.2	212	63.5	26.9	50.4	56.9	12.8	28.1	110	24.3	1675	3.69
54	89.2	97.1	218	63.5	26.9	50.4	56.6	13.2	29.0	110	24.3	1677	3.70
55	88.8	97.0	224	63.6	26.9	50.5	56.5	13.5	29.8	110	24.3	1678	3.70
56	88.4	96.9	230	63.6	26.9	50.5	56.2	13.9	30.7	110	24.3	1679	3.70
57	87.9	96.8	236	63.7	27.0	50.6	56.0	14.3	31.5	110	24.3	1680	3.70
58	87.4	96.8	242	63.7	27.0	50.6	55.7	14.7	32.3	110	24.3	1681	3.71
59	86.9	96.7	248	63.7	27.0	50.6	55.4	15.0	33.2	110	24.3	1683	3.71
60	86.4	96.6	254	63.8	27.0	50.6	55.1	15.4	34.0	110	24.3	1684	3.71
61	85.9	96.5	260	63.8	27.0	50.6	54.8	15.8	34.8	110	24.3	1685	3.71
62	85.4	96.4	266	63.9	27.0	50.7	54.6	16.2	35.6	110	24.3	1686	3.72
63	84.9	96.4	271	63.9	27.0	50.7	54.3	16.5	36.4	110	24.3	1687	3.72
64	84.4	96.3	277	64.0	27.1	50.8	54.0	16.9	37.2	110	24.3	1689	3.72
65	83.9	96.2	283	64.0	27.1	50.8	53.7	17.3	38.0	110	24.3	1690	3.73
66	83.4	96.1	288	64.1	27.1	50.9	53.5	17.6	38.8	110	24.3	1691	3.73
67	82.9	96.0	294	64.1	27.1	50.9	53.1	18.0	39.6	110	24.3	1692	3.73
68	82.4	96.0	299	64.1	27.1	50.9	52.8	18.3	40.4	110	24.3	1693	3.73
69	81.9	95.9	305	64.2	27.2	51.0	52.6	18.7	41.2	110	24.3	1695	3.74
70	81.4	95.8	310	64.2	27.2	51.0	52.3	19.0	41.9	110	24.3	1696	3.74
71	80.9	95.7	316	64.3	27.2	51.0	52.0	19.4	42.7	110	24.3	1697	3.74
72	80.4	95.6	321	64.3	27.2	51.0	51.7	19.7	43.5	110	24.3	1698	3.74
73	79.9	95.6	326	64.4	27.3	51.1	51.5	20.1	44.2	110	24.3	1699	3.75
74	79.4	95.5	332	64.4	27.3	51.1	51.1	20.4	45.0	110	24.3	1701	3.75
75	78.9	95.4	337	64.5	27.3	51.2	50.9	20.7	45.7	110	24.3	1702	3.75
76	78.4	95.3	342	64.5	27.3	51.2	50.6	21.1	46.5	110	24.3	1703	3.75
77	77.8	95.2	347	64.6	27.3	51.3	50.3	21.4	47.2	110	24.3	1704	3.76
78	77.2	95.2	353	64.6	27.3	51.3	49.9	21.8	48.0	110	24.3	1705	3.76
79	76.6	95.1	358	64.7	27.4	51.3	49.6	22.1	48.7	111	24.5	1707	3.76
80	76.0	95.0	363	64.8	27.4	51.4	49.2	22.4	49.4	111	24.5	1708	3.77
81	75.4	94.9	368	64.9	27.5	51.5	48.9	22.7	50.1	111	24.5	1709	3.77
82	74.8	94.8	373	64.9	27.5	51.5	48.5	23.1	50.8	111	24.5	1710	3.77
83	74.2	94.7	378	65.0	27.5	51.6	48.2	23.4	51.5	111	24.5	1711	3.77
84	73.6	94.6	383	65.1	27.6	51.7	47.9	23.7	52.2	111	24.5	1713	3.78
85	73.0	94.6	387	65.2	27.6	51.7	47.6	24.0	52.9	111	24.5	1714	3.78
86	72.4	94.5	392	65.2	27.6	51.7	47.2	24.3	53.6	111	24.5	1715	3.78
87	71.8	94.4	397	65.3	27.6	51.8	46.9	24.6	54.3	111	24.5	1716	3.78
88	71.2	94.3	402	65.4	27.7	51.9	46.6	24.9	55.0	111	24.5	1717	3.78
89	70.6	94.2	406	65.5	27.7	52.0	46.2	25.2	55.6	111	24.5	1719	3.79
90	70.0	94.1	411	65.6	27.8	52.1	45.9	25.5	56.3	111	24.5	1720	3.79



## APPENDIX 4

### BOVANS WHITE EGG WEIGHT DISTRIBUTION (%) – CANADIAN SYSTEM

Age in weeks	Weekly						Cumulative					
	Jumbo over 70 g	Extra Large 64-70 g	Large 56-64 g	Medium 49-56 g	Small 42-49 g	Peewee under 42 g	Jumbo over 70 g	Extra Large 64-70 g	Large 56-64 g	Medium 49-56 g	Small 42-49 g	Peewee under 42 g
19	0.0	0.0	0.0	6.5	62.2	31.3	0.0	0.0	0.0	6.5	62.2	31.3
20	0.0	0.0	0.5	24.5	63.6	11.3	0.0	0.0	0.5	22.9	63.5	13.1
21	0.0	0.0	5.0	50.0	42.1	2.9	0.0	0.0	3.2	38.9	50.8	7.1
22	0.0	0.1	14.8	59.4	24.7	1.0	0.0	0.1	8.1	47.6	39.7	4.5
23	0.0	0.8	28.8	56.7	13.4	0.3	0.0	0.3	14.6	50.5	31.4	3.2
24	0.0	1.5	35.9	52.6	9.8	0.2	0.0	0.6	19.9	51.0	26.0	2.4
25	0.1	3.8	47.0	43.4	5.6	0.1	0.0	1.3	25.4	49.5	21.9	2.0
26	0.3	6.6	53.5	36.0	3.6	0.0	0.1	2.2	30.2	47.2	18.8	1.6
27	0.7	11.3	58.3	27.5	2.1	0.0	0.2	3.5	34.3	44.3	16.3	1.4
28	1.0	13.5	59.3	24.5	1.7	0.0	0.3	4.8	37.5	41.8	14.5	1.2
29	1.4	15.9	59.7	21.7	1.4	0.0	0.4	6.1	40.0	39.5	13.0	1.1
30	1.8	17.9	59.6	19.6	1.1	0.0	0.5	7.3	42.0	37.4	11.8	1.0
31	2.2	19.9	59.2	17.7	1.0	0.0	0.7	8.4	43.6	35.6	10.8	0.9
32	2.6	21.5	58.7	16.3	0.8	0.0	0.9	9.5	44.9	34.0	9.9	0.8
33	2.9	22.6	58.3	15.4	0.8	0.0	1.0	10.6	45.9	32.5	9.2	0.7
34	3.3	23.7	57.8	14.6	0.7	0.0	1.2	11.5	46.8	31.3	8.6	0.7
35	3.4	24.2	57.5	14.2	0.7	0.0	1.3	12.4	47.5	30.1	8.1	0.6
36	3.6	24.8	57.2	13.8	0.6	0.0	1.5	13.1	48.1	29.1	7.6	0.6
37	4.0	25.8	56.6	13.0	0.6	0.0	1.6	13.9	48.6	28.1	7.2	0.6
38	4.2	26.4	56.3	12.6	0.5	0.0	1.8	14.6	49.0	27.3	6.8	0.5
39	4.4	26.9	55.9	12.3	0.5	0.0	1.9	15.2	49.4	26.5	6.5	0.5
40	4.6	27.4	55.5	11.9	0.5	0.0	2.0	15.8	49.7	25.8	6.2	0.5
41	4.9	28.0	55.1	11.6	0.5	0.0	2.2	16.4	50.0	25.1	5.9	0.5
42	5.3	29.0	54.3	10.9	0.4	0.0	2.3	16.9	50.2	24.5	5.7	0.4
43	5.8	30.0	53.5	10.2	0.4	0.0	2.5	17.5	50.3	23.9	5.5	0.4
44	6.4	31.0	52.6	9.6	0.4	0.0	2.6	18.1	50.4	23.3	5.2	0.4
45	7.0	32.0	51.6	9.0	0.3	0.0	2.8	18.6	50.4	22.7	5.1	0.4
46	7.3	32.5	51.1	8.8	0.3	0.0	3.0	19.1	50.5	22.2	4.9	0.4
47	7.6	33.0	50.6	8.5	0.3	0.0	3.1	19.6	50.5	21.7	4.7	0.4
48	7.9	33.4	50.1	8.2	0.3	0.0	3.3	20.1	50.5	21.3	4.6	0.3
49	8.6	34.3	49.1	7.7	0.3	0.0	3.5	20.6	50.4	20.8	4.4	0.3
50	8.9	34.8	48.6	7.5	0.2	0.0	3.6	21.0	50.4	20.4	4.3	0.3



## BOVANS WHITE EGG WEIGHT DISTRIBUTION (%) – CANADIAN SYSTEM

Age in weeks	Weekly						Cumulative					
	Jumbo over 70 g	Extra Large 64-70 g	Large 56-64 g	Medium 49-56 g	Small 42-49 g	Peewee under 42 g	Jumbo over 70 g	Extra Large 64-70 g	Large 56-64 g	Medium 49-56 g	Small 42-49 g	Peewee under 42 g
51	9.3	35.2	48.0	7.2	0.2	0.0	3.8	21.5	50.3	20.0	4.2	0.3
52	9.7	35.6	47.5	7.0	0.2	0.0	4.0	21.9	50.2	19.6	4.0	0.3
53	10.0	36.0	46.9	6.8	0.2	0.0	4.2	22.3	50.1	19.2	3.9	0.3
54	10.0	36.0	46.9	6.8	0.2	0.0	4.3	22.7	50.0	18.9	3.8	0.3
55	10.4	36.4	46.4	6.6	0.2	0.0	4.5	23.0	49.9	18.5	3.7	0.3
56	10.4	36.4	46.4	6.6	0.2	0.0	4.6	23.4	49.8	18.2	3.6	0.3
57	10.8	36.8	45.8	6.3	0.2	0.0	4.8	23.7	49.7	17.9	3.6	0.3
58	10.8	36.8	45.8	6.3	0.2	0.0	4.9	24.0	49.6	17.7	3.5	0.3
59	10.8	36.8	45.8	6.3	0.2	0.0	5.1	24.3	49.5	17.4	3.4	0.3
60	11.2	37.2	45.2	6.1	0.2	0.0	5.2	24.6	49.4	17.1	3.3	0.3
61	11.2	37.2	45.2	6.1	0.2	0.0	5.4	24.9	49.3	16.9	3.3	0.2
62	11.6	37.6	44.7	5.9	0.2	0.0	5.5	25.2	49.2	16.6	3.2	0.2
63	11.6	37.6	44.7	5.9	0.2	0.0	5.6	25.5	49.1	16.4	3.1	0.2
64	12.1	37.9	44.1	5.7	0.2	0.0	5.8	25.7	49.0	16.2	3.1	0.2
65	12.1	37.9	44.1	5.7	0.2	0.0	5.9	26.0	48.9	16.0	3.0	0.2
66	12.5	38.3	43.5	5.5	0.2	0.0	6.0	26.2	48.8	15.8	2.9	0.2
67	12.5	38.3	43.5	5.5	0.2	0.0	6.1	26.4	48.7	15.6	2.9	0.2
68	12.5	38.3	43.5	5.5	0.2	0.0	6.3	26.6	48.6	15.4	2.8	0.2
69	12.9	38.6	42.9	5.4	0.2	0.0	6.4	26.9	48.5	15.2	2.8	0.2
70	12.9	38.6	42.9	5.4	0.2	0.0	6.5	27.1	48.4	15.0	2.7	0.2
71	13.4	38.9	42.3	5.2	0.1	0.0	6.6	27.3	48.3	14.9	2.7	0.2
72	13.4	38.9	42.3	5.2	0.1	0.0	6.7	27.5	48.2	14.7	2.7	0.2
73	13.9	39.2	41.8	5.0	0.1	0.0	6.8	27.7	48.1	14.6	2.6	0.2
74	13.9	39.2	41.8	5.0	0.1	0.0	6.9	27.8	48.0	14.4	2.6	0.2
75	14.3	39.5	41.2	4.8	0.1	0.0	7.1	28.0	47.9	14.3	2.5	0.2
76	14.3	39.5	41.2	4.8	0.1	0.0	7.2	28.2	47.8	14.1	2.5	0.2
77	14.8	39.8	40.6	4.7	0.1	0.0	7.3	28.4	47.7	14.0	2.5	0.2
78	14.8	39.8	40.6	4.7	0.1	0.0	7.4	28.5	47.6	13.8	2.4	0.2
79	15.3	40.1	40.0	4.5	0.1	0.0	7.5	28.7	47.5	13.7	2.4	0.2
80	15.8	40.3	39.4	4.4	0.1	0.0	7.6	28.9	47.4	13.6	2.4	0.2
81	16.3	40.6	38.8	4.2	0.1	0.0	7.7	29.0	47.3	13.4	2.3	0.2
82	16.3	40.6	38.8	4.2	0.1	0.0	7.9	29.2	47.2	13.3	2.3	0.2
83	16.8	40.8	38.2	4.1	0.1	0.0	8.0	29.3	47.0	13.2	2.3	0.2
84	17.3	41.0	37.6	3.9	0.1	0.0	8.1	29.5	46.9	13.1	2.3	0.2
85	17.9	41.2	37.0	3.8	0.1	0.0	8.2	29.6	46.8	13.0	2.2	0.2
86	17.9	41.2	37.0	3.8	0.1	0.0	8.3	29.8	46.7	12.9	2.2	0.2
87	18.4	41.4	36.4	3.7	0.1	0.0	8.5	29.9	46.6	12.7	2.2	0.2
88	19.0	41.6	35.8	3.5	0.1	0.0	8.6	30.0	46.4	12.6	2.2	0.2
89	19.5	41.7	35.2	3.4	0.1	0.0	8.7	30.2	46.3	12.5	2.1	0.2
90	20.1	41.9	34.7	3.3	0.1	0.0	8.8	30.3	46.2	12.4	2.1	0.2



## APPENDIX 5

### BOVANS WHITE EGG WEIGHT DISTRIBUTION (%) – U.S.A. SYSTEM

Age in weeks	Weekly						Cumulative					
	Jumbo over 30 oz/doz	Extra Large 27-30 oz/doz	Large 24-27 oz/doz	Medium 21-24 oz/doz	Small 18-21 oz/doz	Peewee under 18 oz/doz	Jumbo over 30 oz/doz	Extra Large 27-30 oz/doz	Large 24-27 oz/doz	Medium 21-24 oz/doz	Small 18-21 oz/doz	Peewee under 18 oz/doz
19	0.0	0.0	0.0	4.6	58.9	36.6	0.0	0.0	0.0	4.6	58.9	36.6
20	0.0	0.0	0.3	19.9	65.7	14.1	0.0	0.0	0.3	18.5	65.0	16.2
21	0.0	0.0	3.4	45.5	47.2	3.9	0.0	0.0	2.1	34.5	54.4	8.9
22	0.0	0.2	11.2	58.1	29.3	1.3	0.0	0.1	6.0	44.5	43.7	5.7
23	0.0	0.9	23.3	58.8	16.5	0.5	0.0	0.3	11.5	49.0	35.1	4.0
24	0.0	1.7	29.8	55.8	12.3	0.3	0.0	0.7	16.0	50.7	29.5	3.1
25	0.0	4.3	40.4	47.9	7.3	0.1	0.0	1.4	20.9	50.2	25.0	2.5
26	0.1	7.3	46.9	40.8	4.8	0.1	0.0	2.4	25.4	48.6	21.5	2.1
27	0.4	12.5	52.1	32.2	2.8	0.0	0.1	3.9	29.3	46.2	18.8	1.8
28	0.6	14.9	53.2	29.0	2.3	0.0	0.2	5.3	32.4	43.9	16.7	1.5
29	0.8	17.5	53.9	25.9	1.9	0.0	0.2	6.7	34.8	41.9	15.0	1.4
30	1.1	19.7	54.0	23.6	1.6	0.0	0.3	8.0	36.8	40.0	13.6	1.2
31	1.4	22.0	53.8	21.4	1.3	0.0	0.4	9.3	38.3	38.3	12.5	1.1
32	1.7	23.8	53.5	19.9	1.2	0.0	0.5	10.5	39.6	36.8	11.5	1.0
33	1.9	25.0	53.2	18.9	1.1	0.0	0.6	11.7	40.7	35.4	10.7	0.9
34	2.1	26.2	52.8	17.9	1.0	0.0	0.7	12.7	41.6	34.1	10.0	0.9
35	2.2	26.8	52.6	17.5	0.9	0.0	0.8	13.7	42.3	33.0	9.4	0.8
36	2.4	27.4	52.3	17.0	0.9	0.0	0.9	14.5	42.9	32.0	8.9	0.8
37	2.7	28.6	51.8	16.1	0.8	0.0	1.0	15.3	43.4	31.1	8.4	0.7
38	2.8	29.2	51.5	15.7	0.8	0.0	1.1	16.1	43.9	30.2	8.0	0.7
39	3.0	29.8	51.2	15.3	0.7	0.0	1.2	16.8	44.3	29.4	7.6	0.7
40	3.1	30.4	50.9	14.8	0.7	0.0	1.3	17.5	44.6	28.7	7.3	0.6
41	3.3	31.0	50.6	14.4	0.7	0.0	1.4	18.1	44.9	28.0	6.9	0.6
42	3.6	32.2	49.9	13.6	0.6	0.0	1.5	18.8	45.1	27.4	6.7	0.6
43	4.0	33.4	49.2	12.9	0.6	0.0	1.6	19.4	45.3	26.8	6.4	0.5
44	4.4	34.5	48.4	12.2	0.5	0.0	1.7	20.0	45.4	26.2	6.2	0.5
45	4.9	35.7	47.5	11.5	0.5	0.0	1.9	20.6	45.5	25.6	5.9	0.5
46	5.1	36.2	47.1	11.1	0.4	0.0	2.0	21.2	45.5	25.1	5.7	0.5
47	5.3	36.8	46.7	10.8	0.4	0.0	2.1	21.7	45.6	24.6	5.5	0.5
48	5.6	37.3	46.2	10.5	0.4	0.0	2.2	22.3	45.6	24.1	5.4	0.4
49	6.1	38.4	45.3	9.9	0.4	0.0	2.3	22.8	45.6	23.6	5.2	0.4
50	6.4	38.9	44.8	9.6	0.4	0.0	2.5	23.3	45.6	23.2	5.0	0.4



## BOVANS WHITE EGG WEIGHT DISTRIBUTION (%) – U.S.A. SYSTEM


Age in weeks	Weekly						Cumulative					
	Jumbo over 30 oz/doz	Extra Large 27-30 oz/doz	Large 24-27 oz/doz	Medium 21-24 oz/doz	Small 18-21 oz/doz	Peewee under 18 oz/doz	Jumbo over 30 oz/doz	Extra Large 27-30 oz/doz	Large 24-27 oz/doz	Medium 21-24 oz/doz	Small 18-21 oz/doz	Peewee under 18 oz/doz
51	6.7	39.4	44.3	9.3	0.3	0.0	2.6	23.8	45.5	22.7	4.9	0.4
52	7.0	39.9	43.8	9.0	0.3	0.0	2.7	24.3	45.5	22.3	4.8	0.4
53	7.3	40.4	43.3	8.7	0.3	0.0	2.9	24.8	45.4	21.9	4.6	0.4
54	7.3	40.4	43.3	8.7	0.3	0.0	3.0	25.2	45.4	21.6	4.5	0.4
55	7.6	40.9	42.8	8.5	0.3	0.0	3.1	25.6	45.3	21.2	4.4	0.4
56	7.6	40.9	42.8	8.5	0.3	0.0	3.2	26.0	45.2	20.9	4.3	0.4
57	7.9	41.3	42.3	8.2	0.3	0.0	3.3	26.4	45.2	20.6	4.2	0.3
58	7.9	41.3	42.3	8.2	0.3	0.0	3.5	26.8	45.1	20.3	4.1	0.3
59	7.9	41.3	42.3	8.2	0.3	0.0	3.6	27.1	45.0	20.0	4.0	0.3
60	8.2	41.8	41.8	7.9	0.3	0.0	3.7	27.4	44.9	19.7	3.9	0.3
61	8.2	41.8	41.8	7.9	0.3	0.0	3.8	27.8	44.9	19.4	3.8	0.3
62	8.5	42.2	41.3	7.7	0.3	0.0	3.9	28.1	44.8	19.2	3.8	0.3
63	8.5	42.2	41.3	7.7	0.3	0.0	4.0	28.4	44.7	18.9	3.7	0.3
64	8.9	42.7	40.7	7.5	0.2	0.0	4.1	28.7	44.6	18.7	3.6	0.3
65	8.9	42.7	40.7	7.5	0.2	0.0	4.2	29.0	44.6	18.5	3.6	0.3
66	9.2	43.1	40.2	7.2	0.2	0.0	4.3	29.2	44.5	18.3	3.5	0.3
67	9.2	43.1	40.2	7.2	0.2	0.0	4.4	29.5	44.4	18.0	3.4	0.3
68	9.2	43.1	40.2	7.2	0.2	0.0	4.5	29.7	44.3	17.8	3.4	0.3
69	9.6	43.5	39.7	7.0	0.2	0.0	4.5	30.0	44.2	17.7	3.3	0.3
70	9.6	43.5	39.7	7.0	0.2	0.0	4.6	30.2	44.2	17.5	3.3	0.3
71	10.0	43.9	39.2	6.8	0.2	0.0	4.7	30.5	44.1	17.3	3.2	0.3
72	10.0	43.9	39.2	6.8	0.2	0.0	4.8	30.7	44.0	17.1	3.2	0.3
73	10.4	44.3	38.6	6.5	0.2	0.0	4.9	30.9	43.9	16.9	3.1	0.2
74	10.4	44.3	38.6	6.5	0.2	0.0	5.0	31.1	43.8	16.8	3.1	0.2
75	10.7	44.7	38.1	6.3	0.2	0.0	5.1	31.3	43.7	16.6	3.0	0.2
76	10.7	44.7	38.1	6.3	0.2	0.0	5.2	31.5	43.6	16.4	3.0	0.2
77	11.1	45.0	37.5	6.1	0.2	0.0	5.3	31.7	43.5	16.3	2.9	0.2
78	11.1	45.0	37.5	6.1	0.2	0.0	5.3	31.9	43.5	16.1	2.9	0.2
79	11.5	45.4	37.0	5.9	0.2	0.0	5.4	32.1	43.4	16.0	2.9	0.2
80	12.0	45.7	36.4	5.7	0.2	0.0	5.5	32.3	43.3	15.9	2.8	0.2
81	12.4	46.0	35.9	5.6	0.2	0.0	5.6	32.5	43.2	15.7	2.8	0.2
82	12.4	46.0	35.9	5.6	0.2	0.0	5.7	32.7	43.1	15.6	2.7	0.2
83	12.8	46.3	35.4	5.4	0.2	0.0	5.8	32.9	43.0	15.4	2.7	0.2
84	13.3	46.6	34.8	5.2	0.1	0.0	5.9	33.0	42.9	15.3	2.7	0.2
85	13.7	46.9	34.3	5.0	0.1	0.0	6.0	33.2	42.8	15.2	2.6	0.2
86	13.7	46.9	34.3	5.0	0.1	0.0	6.1	33.4	42.7	15.1	2.6	0.2
87	14.2	47.1	33.7	4.9	0.1	0.0	6.2	33.5	42.6	14.9	2.6	0.2
88	14.7	47.4	33.2	4.7	0.1	0.0	6.3	33.7	42.4	14.8	2.6	0.2
89	15.1	47.6	32.6	4.5	0.1	0.0	6.4	33.9	42.3	14.7	2.5	0.2
90	15.6	47.8	32.1	4.4	0.1	0.0	6.5	34.0	42.2	14.6	2.5	0.2





## BODYWEIGHT REARING GRAPH


Bovans White commercial layers





# PRODUCTION RECORDING GRAPH

Bovans White commercial layers








## EGG MASS & FEED CONSUMPTION GRAPH

Bovans White commercial layers





**Institut de Sélection Animale BV**  
Villa 'de Körver'  
Spoorstraat 69, 5831 CK Boxmeer  
P.O. Box 114, 5830 AC Boxmeer  
The Netherlands-EU  
T +31 485 319 111  
F +31 485 319 112  
[www.isapoultry.com](http://www.isapoultry.com)

**ISA North America**  
650 Riverbend Drive, Suite C  
Kitchener, Ontario  
Canada N2K 3S2  
  
T +1 519 578 2740  
F +1 519 578 1870  
[www.isapoultry.com](http://www.isapoultry.com)