


Exam Questions – Chapter 6 + 7 Trig (A2)







The shape *ABCDEA*, as shown in Figure 2, consists of a right-angled triangle *EAB* and a triangle *DBC* joined to a sector *BDE* of a circle with radius 5 cm and centre *B*.

The points A, B and C lie on a straight line with BC = 7.5 cm.

Angle $EAB = \frac{\pi}{2}$ radians, angle EBD = 1.4 radians and CD = 6.1 cm.

(a) Find, in cm^2 , the area of the sector *BDE*.

(b) Find the size of the angle DBC, giving your answer in radians to 3 decimal places.

(c) Find, in cm², the area of the shape *ABCDEA*, giving your answer to 3 significant figures.

(5)

(2)

(2)

(Total 9 marks)

رکیا مگرم Online Maths Teaching

Q2.

The depth of water, D metres, in a harbour on a particular day is modelled by the formula

 $D = 5 + 2\sin(30t)^\circ \quad 0 \le t < 24$

where *t* is the number of hours after midnight.

A boat enters the harbour at 6:30 am and it takes 2 hours to load its cargo. The boat requires the depth of water to be at least 3.8 metres before it can leave the harbour.

- (a) Find the depth of the water in the harbour when the boat enters the harbour.
- (b) Find, to the nearest minute, the earliest time the boat can leave the harbour. (Solutions based entirely on graphical or numerical methods are not acceptable.)

(4)

(1)

(Total for question = 5 marks)

Q3.

On a roller coaster ride, passengers travel in carriages around a track.

On the ride, carriages complete multiple circuits of the track such that

- the maximum vertical height of a carriage above the ground is 60 m
- a carriage starts a circuit at a vertical height of 2 m above the ground
- the ground is horizontal

The vertical height, H m, of a carriage above the ground, t seconds after the carriage starts the first circuit, is modelled by the equation

$$H = a - b(t - 20)^2$$

where a and b are positive constants.

(a) Find a complete equation for the model.

(3)

(b) Use the model to determine the height of the carriage above the ground when t = 40

(1)

In an alternative model, the vertical height, H m, of a carriage above the ground, t seconds after the ^{Teaching} carriage starts the first circuit, is given by

 $H = 29\cos(9t + \alpha)^\circ + \beta \qquad 0 \le \alpha < 360^\circ$

where α and β are constants.

(c) Find a complete equation for the alternative model.

Given that the carriage moves continuously for 2 minutes,

(d) give a reason why the alternative model would be more appropriate.

(Total for question = 7 marks)

Q4.

(a) Prove that

(b) Hence, or otherwise, solve, for $-\pi \le x < \pi$,

 $6\cot 2x + 3\tan x = \csc^2 x - 2$

 $2\cot 2x + \tan x \equiv \cot x$ $x \neq \frac{n\pi}{2}, n \in \mathbb{Z}$

Give your answers to 3 decimal places.

(Solutions based entirely on graphical or numerical methods are not acceptable.)

(6)

(4)

(Total for question = 10 marks)

وریا م Online Maths the^{Teaching}

(1)

ريسا مالندو Maths Teaching

Q5.

 $g(\theta) = 4\cos 2\theta + 2\sin 2\theta$

Given that $g(\theta) = R \cos(2\theta - \alpha)$, where R > 0 and $0 < \alpha < 90^{\circ}$,

(a) find the exact value of *R* and the value of α to 2 decimal places.

(b) Hence solve, for $-90^{\circ} < \theta < 90^{\circ}$,

 $4\cos 2\theta + 2\sin 2\theta = 1$

giving your answers to one decimal place.

Given that *k* is a constant and the equation $g(\theta) = k$ has no solutions,

(c) state the range of possible values of k.

(2)

(3)

(5)

(5)

(3)

(Total for question = 10 marks)

Q6.

(a) Express $2\cos\theta - \sin\theta$ in the form $R\cos(\theta + a)$, where *R* and *a* are constants, R > 0 and $0 < a 90^{\circ}$ Give the exact value of *R* and give the value of *a* to 2 decimal places.

(b) Hence solve, for $0 \le \theta < 360^\circ$,

$$\frac{2}{2\cos\theta - \sin\theta - 1} = 15$$

Give your answers to one decimal place.

(c) Use your solutions to parts (a) and (b) to deduce the smallest positive value of θ for which

$$\frac{2}{2\cos\theta + \sin\theta - 1} = 15$$

Give your answer to one decimal place.

(2)

(Total for question = 10 marks)

In this question you must show all stages of your working.

Solutions relying entirely on calculator technology are not acceptable.

(a) Show that

$$\frac{1}{\cos\theta} + \tan\theta \equiv \frac{\cos\theta}{1 - \sin\theta} \qquad \theta \neq (2n+1)90^{\circ} \quad n \in \mathbb{Z}$$

Given that $\cos 2x \neq 0$

(b) solve for $0 < x < 90^{\circ}$ $\frac{1}{\cos 2x} + \tan 2x = 3\cos 2x$ giving your answers to one decimal place.

_ /	-	Λ.
	~	- 1
	\smile	

(3)

(Total	for	question = 8 I	marks
٩			940001011 - 01	mai no

Q8.

(a) Show that the equation

 $\cos^2 x = 8\sin^2 x - 6\sin x$

can be written in the form

 $(3\sin x - 1)^2 = 2$

(b) Hence solve, for $0 \le x < 360^\circ$,

$$\cos^2 x = 8\sin^2 x - 6\sin x$$

giving your answers to 2 decimal places.

(5)

(3)

(Total for question = 8 marks)

وکریا مانبه Maths Teaching

Q9.

(a) Prove that

 $\sin 2x - \tan x \equiv \tan x \cos 2x$, $x \neq (2n + 1)90^\circ$, $n \in \mathbb{Z}$

(b) Given that $x \neq 90^{\circ}$ and $x \neq 270^{\circ}$, solve, for $0 \le x < 360^{\circ}$,

 $\sin 2x - \tan x = 3\tan x \sin x$

Give your answers in degrees to one decimal place where appropriate. (Solutions based entirely on graphical or numerical methods are not acceptable.)

(5)

(4)

(Total for question = 9 marks)

Q10.

(i) Solve, for $0 \le \theta < 360^{\circ}$, the equation

 $9\sin(\theta + 60^\circ) = 4$

giving your answers to 1 decimal place. You must show each step of your working.

(ii) Solve, for $-\pi \le x < \pi$, the equation

 $2\tan x - 3\sin x = 0$

giving your answers to 2 decimal places where appropriate. [Solutions based entirely on graphical or numerical methods are not acceptable.]

(5)

(4)

(Total 9 marks)

Q11.

In this question you must show detailed reasoning.

Solutions relying entirely on calculator technology are not acceptable.

(a) Show that the equation

 $4\tan x = 5\cos x$

can be written as

$$5\sin^2 x + 4\sin x - 5 = 0$$

(3)

(b) Hence solve, for $0 < x \le 360^{\circ}$

$4\tan x = 5\cos x$

giving your answers to one decimal place.

(c) Hence find the number of solutions of the equation

$$4\tan 3x = 5\cos 3x$$

in the interval $0 < x \le 1800^\circ$, explaining briefly the reason for your answer.

(2)

(Total for question = 9 marks)




Figure 1

The triangle XYZ in Figure 1 has XY = 6 cm, YZ = 9 cm, ZX = 4 cm and angle ZXY = a. The point W lies on the line XY.

The circular arc ZW, in Figure 1 is a major arc of the circle with centre X and radius 4 cm.

(a) Show that, to 3 significant figures, a = 2.22 radians.

(b) Find the area, in cm^2 , of the major sector *XZWX*.

The region enclosed by the major arc ZW of the circle and the lines WY and YZ is shown shaded in Figure 1.

Calculate

Q12.

7

- (c) the area of this shaded region,
- (d) the perimeter ZWYX of this shaded region.

(Total 12 marks) www.onlinemathsteaching.co.uk



(4)

(2)

(3)

(3)

(4)