



Magnum Compounding LLC
490 W 84 St, Hialeah, FL 33014
Phone: (786) 622-2301
Fax: (786) 622-2319
Toll Free: (855) 403-6028

**LET'S DO THIS
TOGETHER**

COMPOUNDED INJECTABLE FORMULA INFORMATION SHEET

Product Name: Super Lipo (MIC / B12 / B5 / Leucine Complex)

Formulation Type: Sterile Injectable Solution (Intramuscular Use / Subcutaneous Use)

Vial Size: 30 mL

ACTIVE INGREDIENTS (per mL)

Ingredients	Strength (mg / mL)	Role & Benefits
Methionine (L)	12	Lipotropic agent – supports fat metabolism and liver function.
Inositol FCC	25	Lipotropic and metabolic support.
Choline Chloride	25	Hepatoprotective, fat emulsifier.
Cyanocobalamin (Vitamin B12)	1	Supports nerve and red blood cell health.
Leucine (L)	5	Branched-chain amino acid – supports muscle recovery.
Dexpanthenol (Vitamin B5)	1.5	Skin and nerve health; metabolic support.

OTHER INGREDIENTS:

- Benzyl Alcohol, NF – Preservative (1%)
- Sterile Water for Injection, USP – Diluent/Base
- pH adjusted with Sodium Hydroxide and/or Hydrochloric

DESCRIPTION: Super Lipo is a compounded sterile injectable solution composed of lipotropic nutrients, vitamins, and amino acids. It is designed to support liver detoxification, fat metabolism, energy production, and overall metabolic function.

INDICATIONS:

- Adjunct in weight management programs
- Metabolic and liver support
- Nutrient repletion (e.g., B12 deficiency)
- General wellness and fatigue support

DOSAGE & ADMINISTRATION:

- Route: Subcutaneous (SC) or Intramuscular (IM) injection, per provider discretion
- Typical Dosage: 1–3 mL weekly or as directed by the healthcare provider
- Frequency: Once or twice weekly, depending on clinical needs

*This product is to be administered under licensed medical supervision.

If you have any questions or concerns regarding your medication, please do not hesitate to contact us.



Magnum Compounding LLC
490 W 84 St, Hialeah, FL 33014
Phone: (786) 622-2301
Fax: (786) 622-2319
Toll Free: (855) 403-6028

**LET'S DO THIS
TOGETHER**

STORAGE: Store at room temperature (59°F to 77°F). Protect from light. Do not freeze.

WARNINGS & PRECAUTIONS:

- For intramuscular and subcutaneous use
- Contraindicated in patients with known hypersensitivity to any ingredient
- Use caution in hepatic or renal impairment
- Monitor vitamin B12 levels and liver function with long-term use

POTENTIAL SIDE EFFECTS:

- Injection site reactions (pain, redness)
- Dizziness or mild gastrointestinal discomfort
- Rare allergic reactions

DRUG DISPOSAL: Unused or expired compounded medications should be disposed of properly according to local regulations. Do not flush medications down the toilet or pour them into a drain unless instructed to do so. Consult with your pharmacy or local waste disposal authority for appropriate disposal methods. Magnum Compounding LLC can provide guidance or accept returned compounds for proper disposal when applicable.

INJECTION SAFETY CONSIDERATIONS (CDC GUIDANCE):

- Medications should be drawn up in a designated clean medication preparation area using a new sterile syringe and sterile needle.
- Prepare an injection as close as possible to the time of administration to the patient.
- Do NOT leave the needle inserted into a medication vial septum for multiple uses.
- Vials labeled as single-dose or single-use should be used for only a single patient and entered only once.
- Do NOT combine (pool) leftover contents of single-dose or single-use vials or store them for later use.
- If a single-dose or single-use vial has been opened or accessed (e.g., needle-punctured), it should be discarded according to pharmacy policy or at the end of the procedure, whichever is sooner.
- Discard any medication vial if sterility is compromised or questionable.
- Multi-dose vials should be dedicated to a single patient whenever possible.

REFERENCE:

1. United States Pharmacopeia (USP)
2. Clinical Compounding Standards
3. Peer-reviewed literature on lipotropic injection therapy

If you have any questions or concerns regarding your medication, please do not hesitate to contact us.