



THE COASTAL NC DAYLILY SOCIETY NEWSLETTER

JANUARY 2023

No. 121

PRESIDENT'S CORNER



Beverly Ambrose

HAPPY NEW YEAR!! Hope everyone had a great holiday season with family and friends.

If you missed our December meeting; it can be viewed on our Facebook page, Coastal North Carolina Daylily Society. We're getting pretty good at going Live! We will be switching things up a little for 2023 to avoid conflicts with family events. Dates for this year's meetings will be **15 Jan, 19 Mar, 21 May, 16 Jul, 17 Sep** and **19 Nov**. All meetings will begin at **2 PM**, locations to be determined.

If you are planning to join us for 2023 but haven't yet paid your dues you can send a check made out to CNCDS to Pat Mansfield, 503 W Bear Swamp Rd, Hertford NC 27944 or PayPal Family and Friends to CoastalNCDaylilies@gmail.com. Dues are \$10 for a single membership and \$15 for a family. For those members of CNCDS that are not members of AHS/ADS, the club voted to pay for your memberships!!

I have only received 1 response to club colors for the new t-shirts. After reviewing the colors available for all 3 styles, Mint Green, Royal and Sapphire are available for all. Please take a look (email 20 Dec) and let us know what you think at our 15 Jan meeting if not before.

We need to make a decision on what hybridizer we would like to get our *Plants for Members* from for this year. At our last meeting Darlene suggested getting them from Patrick and Grace Stamile and I agree. They sent great plants to the National Convention last year and they performed and bloomed beautifully while our club was growing them. Check them out at distinctly.on.ca/stamiledaylilies and let us know what you think. Other suggestions are welcome. I'm thinking we should also have a backup chosen just in case our first choice isn't able to accommodate our request.

Our **January 15 meeting** will be held at **Great Hope Baptist Church**, 131 Great Hope Church Rd, Hertford NC at 2PM.



Ken Ferguson

STARTING THE NEW YEAR...

This is the best I could come up with -

THE OFFICIAL COMPREHENSIVE
IDENTIFICATION GUIDE FOR
DAYLILIES



DAYLILY



DAYLILY



DAYLILY



DAYLILY



DAYLILY



DAYLILY



DAYLILY



DAYLILY



DAYLILY

Fewer moths, more flies: Complex relationships between plants and pollinators have changed dramatically

by [Helmholtz Association of German Research Centres](#)



Volucella bombylans. Credit: [gailhampshire](#) from Cradley, Malvern, U.K./Wikimedia Commons

Their service is invaluable. The army of insects and other animals that pollinates the numerous plants of this earth has an essential function. Without these flower visitors, numerous wild plants could reproduce only poorly—or not at all. Ecosystems would thus no longer be able to function in their current form. More than three quarters of the most important crops depend on pollinators in order to be able to produce a high yield and good quality. A loss of pollinators would therefore also lead to economic losses amounting to billions.

It remains unclear whether the pollinators will still be able to provide their usual service. That's because plants and their pollinators are intertwined in an elaborate network in which the distribution and abundance of the species involved as well as their seasonal occurrence, physiology, and behavior are finely tuned. Even small changes could throw everything out of balance. Experts thus fear that human influences such as changes in climate and [land use](#) could lead to less effective pollination services.

However, because few studies have investigated the interaction between plants and different pollinator groups over longer periods of time, it is difficult to say whether and to what extent such developments are already under way. This makes the more than 120-year-old data from Finland on which the new study is based all the more exciting. Between 1895 and 1900, in the vicinity of Kittilä (a village that lies about 120 km north of the Arctic Circle), forester Frans Silén systematically recorded which insects visited which flowers and how often.

"I am passionate about working with historical datasets like this," says Prof. Tiffany Knight from the UFZ. "If you repeat the historic studies again today, it's often the only way to learn about long-term ecological processes." For her, such work also challenges the imagination. "I am trying to understand what motivated the people who collected the data in the past and what challenges they faced," she explains. "This information can then be used to plan a comparable modern study."

The scientists thus first looked around Kittilä for sites where Silén had also made observations—and where the 17 plant species he studied best still grow today. At these sites, the team repeated the pollinator census in 2018 and 2019. The area remains sparsely populated, and little has changed in terms of land use. However, it has not escaped the consequences of climate change. "We have noticed drastic changes in the networks of pollinators," says Leana Zoller from the MLU. Only 7% of the flower visits observed involved the same species of insects and plants as back then. "That is surprisingly little," says Zoller.

In Silén's time, this species was the most frequent visitor to the Arctic raspberry (*Rubus arcticus*) and the woodland geranium (*Geranium sylvaticum*). The bumblebee hoverfly was likely able to effectively transfer the pollen of these species from one plant to the next.

Also moths use a physical advantage during pollination: with their long proboscis, they can also reach the nectar from the base of tubular flowers. This is why they used to be the most frequent visitors of the fringed pink (*Dianthus superbus*) and bladder campion (*Silene vulgaris*), both of which have such flowers.

While these insects have become rarer, the flowers around Kittilä are now getting considerably more visits from bumblebees and certain flies. Whether these animals work as effectively as the earlier pollinators is not yet known. However, one trend in particular concerns the researchers. There are now considerably fewer insects that are specialists for certain flower shapes. These have been replaced by flies of the genus *Thricops*, which visit many different plants. Such generalists are often more robust when it comes to environmental changes; if one of their host plants is lacking, they can easily switch to others. But they also carry the pollen of various other [plant species](#) onto a flower, thereby potentially providing a less effective pollination service than the specialists.

"So far, the pollinator network in our study area still seems to be working well," says Zoller. "There is no evidence so far that the plants are getting too little pollen and are thus less capable of reproducing." But according to the scientists, this can change in the future if changes in the insect communities continues. So far, the flies there seem to be coping with the rising temperatures. But further north in the high Arctic, one study has revealed a massive decrease in the number of flies.

"If this also happens in our study area, it could become a problem," says Zoller. Because at some point, the plants will no longer be able to compensate for the losses in their pollinator network.

More information: Leana Zoller et al, Plant–pollinator network change across a century in the subarctic, *Nature Ecology & Evolution* (2023). [DOI: 10.1038/s41559-022-01928-3](https://doi.org/10.1038/s41559-022-01928-3)



IN SEARCH OF CONTRIBUTORS

Have you read an article that you would like to share with the group? Or, do you want to share some of your experiences with daylilies? Become a contributor to our NEWSLETTER by sending your story to Ken at kferguson@mchsi.com.

We'll even make a custom meme for you at no charge!



Enough said.

See you at the **January 15 meeting, Great Hope Baptist Church**, 131 Great Hope Church Rd, Hertford NC at **2PM**.