



**Tree Survey**

**At**

**Land at Britannia Terrace,  
St. Clears**

*Inspected by:-  
Julian Wilkes BSc.For, MSc.Land Man, MIC.For, TechArborA  
Treescene Ltd  
The Walled Garden  
Old Coedarhydyglyn  
St Nicholas  
Cardiff  
CF5 6SG  
Tel No. 029 20599300*

**14<sup>th</sup> August, 2023**

**Registered Office: Treescene Limited  
The Walled Garden, Old Coedarhydyglyn, St. Nicholas, Cardiff CF5 6SG  
Tel. 029 205 99300 Email. [trees@treescene.co.uk](mailto:trees@treescene.co.uk)**

## **Brief**

I have been instructed by Mr. G Powys Jones to carry out a survey on trees at Land at Britannia Terrace, St. Clears.

## **Scope of Report**

This Tree Survey has been undertaken within the recommendations of British Standards 5837:2012 and current good arboricultural practice.

The survey entailed a visual inspection from ground level of all trees.

Each tree has been numbered and, where instructed, have been tagged using small durable metal or plastic tags.

Due to variations of existing ground levels through the site, height dimensions are estimated and are given in metres.

Trunk/stem diameters are measured at 1.5 metres above ground level, or immediately above the root flare for multi-stemmed trees.

Estimated branch spread is taken in metres from the centre of the trunk, at the four cardinal points of a compass, to achieve an accurate representation of crown shape.

An assessment of a tree's age classification is made in terms of its maturity within the site's landscape.

An assessment of a tree's physiological condition is made as good, fair, poor, dead.

Data on the structural condition of the tree has been entered, e.g., collapsing, leaning and the presence of any decay or physical defect has been noted.

Preliminary management recommendations include further investigation of suspected defects that require more detailed assessment or potential for wildlife habitat.

An assessment of a tree's future life expectancy is made as <10, 10-20, 20-40 or >40 etc.

Table 1 – Cascade chart for tree quality assessment

Category and definition	Criteria (including subcategories where appropriate)		
<u>Category U</u> Those in such a condition that they cannot realistically be retained as living trees in the context of the current land use for longer than 10 years	<ul style="list-style-type: none"> <li>Trees that have a serious, irremediable, structural defect, such that their early loss is expected due to collapse, including those that will become unviable after removal of other U category trees (i.e. where, for whatever reason, the loss of companion shelter cannot be mitigated by pruning)</li> <li>Trees that are dead or are showing signs of significant, immediate, and irreversible overall decline</li> <li>Trees infected with pathogens of significance to the health and/or safety of other trees nearby, or very low quality trees suppressing adjacent trees of better quality</li> </ul> NOTE Category U trees can have existing or potential conservation value which it might be desirable to preserve; see 4.5.7		
	1 Mainly Arboricultural values	2 Mainly landscape values	3 Mainly cultural values, including conservation
<u>Category A</u> Those of high quality with an estimated remaining life expectancy of at least 40 years	Trees that are particularly good examples of their species, especially if rare or unusual, or essential components of groups, or of formal or semi-formal arboricultural features (e.g. the dominant and/or principal trees within an avenue)	Trees, groups or woodlands of particular visual importance as Arboricultural and/or landscape features	Trees, groups or woodlands of significant conservation; historical, commemorative or other value (e.g. veteran trees or wood-pasture)
<u>Category B</u> Those of moderate quality with an estimated remaining life expectancy of at least 20 years	Trees that might be included in category A, but are downgraded because of impaired condition (e.g. presence of significant though remediable defects, including unsympathetic past management and storm damage), such that they are unlikely to be suitable for retention for beyond 40 years; or trees lacking the special quality necessary to merit the category A designation	Trees present in numbers, usually growing as groups or woodlands, such that they attract a higher collective rating than they might as individuals; or trees occurring as collectives but situated so as to make little visual contribution to the wider locality	Trees with material conservation or other cultural benefits
<u>Category C</u> Those of low quality with an estimated remaining life expectancy of at least 10 years, or young trees with a stem diameter below 150mm	Unremarkable trees of very limited merit or such impaired condition that they do not qualify in higher categories	Trees present in groups or woodlands, but without this conferring on them significantly greater collective landscape value, and/or trees offering low or only temporary/transient landscape benefits	Trees with no material conservation or other cultural value

BRITISH STANDARD BS 5837:2012

Tree No.	Species	Height(m)	Single/Multi Stemmed	Stem Diameter(m)	Branch Spread(m)				Height of Crown(m)	Age	Physiological Condition	Structural Condition	Prel. Man. Recommendations	Est. Remaining Contribution	Category
					N	E	S	W							
G1	Group of: Field Maple ( <i>Acer campestre</i> ), Hazel ( <i>Corylus avellana</i> ), Blackthorn ( <i>Prunus spinosa</i> ), Hawthorn ( <i>Crataegus monogyna</i> ), Oak ( <i>Quercus robur</i> )	1	Single and Multi	0.05	0.25	0.25	0.25	0.25	0	Young	Fair	Newly planted hedgerow on raised bank.	No action required at this time.	>40	C
G2	Group of: Field Maple ( <i>Acer campestre</i> ), Hazel ( <i>Corylus avellana</i> ), Blackthorn ( <i>Prunus spinosa</i> ), Hawthorn ( <i>Crataegus monogyna</i> ), Oak ( <i>Quercus robur</i> ), Elder ( <i>Sambucus nigra</i> )	1	Single and Multi	0.05	0.25	0.25	0.25	0.25	0	Young	Fair	Newly planted hedgerow on raised bank.	No action required at this time.	>40	C
T3	Oak ( <i>Quercus robur</i> )	11	Single	0.28	5	6	2	1	3	Middle aged	Fair	Hedgerow tree of reasonable form and vigour.	No action required at this time.	>40	B

Tree No.	Species	Height(m)	Single/Multi Stemmed	Stem Diameter(m)	Branch Spread(m)				Height of Crown(m)	Age	Physiological Condition	Structural Condition	Prel. Man. Recommendations	Est. Remaining Contribution	Category
					N	E	S	W							
T4	Oak ( <i>Quercus robur</i> )	12	Single	0.48	5	3	4	6	2	Middle aged	Fair	Prominent hedgerow tree of reasonable form and vigour. Some mechanical damage on southern side of main stem which is not considered significant at this point in time.	Monitor for health.	>40	B
T5	Oak ( <i>Quercus robur</i> )	11	Single	0.45	5	5	5	3	1	Middle aged	Good	Prominent hedgerow tree of good form and upright habit. Dense ivy colonisation on main stem and mid crown prevents full inspection.	No action required at this time.	>40	B
T6	Oak ( <i>Quercus robur</i> )	11	Single	0.31	4	4	4	4	3	Middle aged	Good	Prominent hedgerow tree of good form and vigour.	No action required at this time.	>40	B
G7	Group of: Goat Willow ( <i>Salix caprea</i> ), Hazel ( <i>Corylus avellana</i> ), Hawthorn ( <i>Crataegus monogyna</i> ), Blackthorn ( <i>Prunus spinosa</i> )	6	Single and Multi	0.15 (avg.)	3	2	2	2	0	Middle aged	Fair	Scrubby specimens forming gappy hedgerow. Specimens of Goat Willow are vulnerable to structural failure as they mature.	Monitor for stability.	20-40	C
T8	Yew ( <i>Taxus baccata</i> )	6	Multi	0.2	1	3	1	1	0	Young	Fair to poor	Hedgerow tree of variable form that has suffered some recent root loss on southeastern side.	Monitor for stability.	10-20	C
T9	Oak ( <i>Quercus robur</i> )	11	Single	0.34	4	3	4	4	1	Middle aged	Good	Prominent hedgerow tree of reasonable upright habit that has recently suffered some root loss on southeastern side.	Monitor for stability.	>40	B

Tree No.	Species	Height(m)	Single/Multi Stemmed	Stem Diameter(m)	Branch Spread(m)				Height of Crown(m)	Age	Physiological Condition	Structural Condition	Prel. Man. Recommendations	Est. Remaining Contribution	Category
					N	E	S	W							
G10	Group of: Oak ( <i>Quercus robur</i> )	12	Single	0.3 (avg.)	4	4	4	3	2	Middle aged	Good	Prominent hedgerow trees of reasonable form and vigour.	No action required at this time.	>40	B
T11	Goat Willow ( <i>Salix caprea</i> )	6	Single	0.21	2	6	1	1	2	Middle aged	Fair to poor	Heavily suppressed specimen leaning to the east. Evidence of slight thinning of foliage.	Monitor for health and stability.	10-20	C
T12	Oak ( <i>Quercus robur</i> )	14	Multi	0.6	6	7	6	6	2	Middle aged	Fair	Prominent hedgerow tree of reasonable form exhibiting some slight thinning of foliage within crown.	Monitor for health.	>40	B
T13	Yew ( <i>Taxus baccata</i> )	10	Multi	0.4	3	3	2	2	0	Middle aged	Fair	Heavily suppressed evergreen tree located on hedge bank.	No action required at this time.	20-40	B
T14	Oak ( <i>Quercus robur</i> )	15	Single	0.77	7	9	8	7	2	Mature	Fair	Prominent hedgerow tree of reasonable upright form that exhibits some slight thinning of foliage in upper crown.	Monitor for health.	>40	B
G15	Group of 2: Ash ( <i>Fraxinus excelsior</i> )	13	Single	0.33	3	3	3	2	2	Middle aged	Poor	Trees located on edge of ditch. Both specimens severely infected with Ash Dieback Disease.	Remove.	<10	U
T16	Oak ( <i>Quercus robur</i> )	12	Single	0.28	5	2	2	4	3	Middle aged	Fair	Slightly suppressed tree located on edge of drainage ditch. Crown more heavily developed on northern side. Some evidence of slight thinning of crown on western side.	Monitor for health.	20-40	C
T17	Ash ( <i>Fraxinus excelsior</i> )	12	Single	0.21	0	1	1	0	3	Middle aged	Poor	Hedgerow tree severely infected with Ash Dieback Disease. This specimen is in a moribund condition.	Remove.	<10	U

Tree No.	Species	Height(m)	Single/Multi Stemmed	Stem Diameter(m)	Branch Spread(m)				Height of Crown(m)	Age	Physiological Condition	Structural Condition	Prel. Man. Recommendations	Est. Remaining Contribution	Category
					N	E	S	W							
T18	Oak ( <i>Quercus robur</i> )	17	Multi	0.85	8	4	8	5	3	Mature	Poor	Prominent hedgerow tree severely infected with Ash Dieback Disease. This specimen is in a moribund condition.	Remove.	<10	U
T19	Oak ( <i>Quercus robur</i> )	11	Single	0.19	2	2	2	2	3	Young	Fair	Hedgerow tree of reasonable form.	No action required at this time.	>40	B
T20	Dead														
G21	Group of: Blackthorn ( <i>Prunus spinosa</i> ), Hawthorn ( <i>Crataegus monogyna</i> ), Hazel ( <i>Corylus avellana</i> ), Elder ( <i>Sambucus nigra</i> ), Oak ( <i>Quercus robur</i> ), Holly ( <i>Ilex aquifolium</i> )	7	Single and Multi	0.15	2	2	2	2	0	Middle aged	Fair	Scrubby specimens forming gappy field boundary hedgerow containing some selected specimens of young Oak.	No action required at this time.	>40	C
T22	Ash ( <i>Fraxinus excelsior</i> )	10	Multi	1.0	4	5	2	4	2	Middle aged	Poor	Multi-stemmed hedgerow tree severely infected with Ash Dieback Disease.	Remove.	<10	U
T23	Oak ( <i>Quercus robur</i> )	17	Multi	0.75	11	11	6	7	2	Mature	Fair	Prominent hedgerow tree of variable form with crown more heavily developed on northeastern side. Main stem and mid crown heavily colonised by ivy thus preventing full inspection. Evidence of slight thinning of foliage in upper crown.	Monitor for health.	20-40	B

Tree No.	Species	Height(m)	Single/Multi Stemmed	Stem Diameter(m)	Branch Spread(m)				Height of Crown(m)	Age	Physiological Condition	Structural Condition	Prel. Man. Recommendations	Est. Remaining Contribution	Category
					N	E	S	W							
G24	Group of 5: Ash ( <i>Fraxinus excelsior</i> )	14	Single and Multi	0.35	4	2	2	1	2	Middle aged	Poor	Hedgerow trees that have died as a result of Ash Dieback Disease.	Remove.	<10	U
G25	Group of 2: Oak ( <i>Quercus robur</i> )	18	Single	0.51	10	6	9	4	2	Middle aged	Fair	Prominent hedgerow trees of variable form and average vigour. Main stems colonised by ivy thus preventing full inspection.	No action required at this time.	>40	B
T26	Dead														
G27	Group of: Holly ( <i>Ilex aquifolium</i> ), Hawthorn ( <i>Crataegus monogyna</i> ), Hazel ( <i>Corylus avellana</i> ), Blackthorn ( <i>Prunus spinosa</i> ), Goat Willow ( <i>Salix caprea</i> )	6	Single and Multi	0.25 (avg.)	2	2	2	2	0	Middle aged	Fair	Scrubby specimens forming gappy field boundary hedgerow.	No action required at this time.	>40	C
T28	Oak ( <i>Quercus robur</i> )	17	Single	0.72	4	9	8	4	2	Mature	Fair	Prominent hedgerow tree slightly suppressed on western side by more dominant Oak. A tree of variable form but reasonable vigour. Main stem and mid crown colonised by ivy thus preventing full inspection.	No action required at this time.	>40	B



Tree No.	Species	Height(m)	Single/Multi Stemmed	Stem Diameter(m)	Branch Spread(m)				Height of Crown(m)	Age	Physiological Condition	Structural Condition	Prel. Man. Recommendations	Est. Remaining Contribution	Category
					N	E	S	W							
T29	Oak ( <i>Quercus robur</i> )	19	Single	0.91	10	13	9	10	2	Mature	Fair	Prominent hedgerow tree of good form and reasonable vigour. Some historic storm damage on eastern side of mid crown has led to some branches becoming hung up within this and adjacent tree's crown. Main stem colonised by ivy thus preventing full inspection. Evidence of slight thinning of foliage throughout crown.	Remove hung up branches. Monitor for health.	>40	B
T30	Ash ( <i>Fraxinus excelsior</i> )	13	Single	0.36	2	3	5	3	3	Middle aged	Poor	Hedgerow tree severely infected with Ash Dieback Disease.	Remove.	<10	U
T31	Oak ( <i>Quercus robur</i> )	10	Single	0.58	2	5	5	5	2	Middle aged	Fair	Scrubby hedgerow tree heavily suppressed by more dominant Ash to the north.	No action required at this time.	20-40	C
T32	Ash ( <i>Fraxinus excelsior</i> )	20	Single	0.72	6	6	6	6	4	Mature	Poor	Prominent hedgerow tree severely infected with Ash Dieback Disease. This specimen is in a moribund condition.	Remove.	<10	U
T33	Oak ( <i>Quercus robur</i> )	13	Single	0.44	5	5	1	4	2	Middle aged	Fair	Hedgerow tree of variable form with crown more heavily developed on northern side.	Monitor for stability.	20-40	B
G34	Group of 3: Ash ( <i>Fraxinus excelsior</i> )	13	Single and Multi	0.4 (avg.)	4	5	5	5	2	Middle aged	Poor	Prominent hedgerow trees infected with Ash Dieback Disease.	Remove.	<10	U
T35	Oak ( <i>Quercus robur</i> )	10	Single	0.47	5	8	4	6	3	Mature	Fair	Hedgerow tree of variable form with crown more heavily developed on eastern side.	Monitor for stability.	20-40	B
T36	Ash ( <i>Fraxinus excelsior</i> )	11	Multi	0.35	1	3	1	0	4	Middle aged	Poor	Hedgerow tree infected with Ash Dieback Disease.	Remove.	<10	U

Tree No.	Species	Height(m)	Single/Multi Stemmed	Stem Diameter(m)	Branch Spread(m)				Height of Crown(m)	Age	Physiological Condition	Structural Condition	Prel. Man. Recommendations	Est. Remaining Contribution	Category
					N	E	S	W							
G37	Group of: Hazel ( <i>Corylus avellana</i> ), Hawthorn ( <i>Crataegus monogyna</i> ), Blackthorn ( <i>Prunus spinosa</i> ), Ash ( <i>Fraxinus excelsior</i> ), Goat Willow ( <i>Salix caprea</i> )	1	Single and Multi	0.05	0.25	0.25	0.25	0.25	0	Young	Fair	Young hedge plants forming sporadic sections of hedgerow on raised bank. This feature is dominated by bracken that is suppressing hedge plants.	Monitor for health.	10-20	C
T38	Oak ( <i>Quercus robur</i> )	12	Single	0.57	7	7	7	7	4	Mature	Fair	Prominent hedgerow tree of good form and well-balanced crown. Some evidence of slight thinning of foliage within crown, particularly on northern side.	Monitor for health.	>40	B

Tree No.	Species	Height(m)	Single/Multi Stemmed	Stem Diameter(m)	Branch Spread(m)				Height of Crown(m)	Age	Physiological Condition	Structural Condition	Prel. Man. Recommendations	Est. Remaining Contribution	Category
					N	E	S	W							
G39	Group of: Hazel ( <i>Corylus avellana</i> ), Ash ( <i>Fraxinus excelsior</i> ), Oak ( <i>Quercus robur</i> ), Holly ( <i>Ilex aquifolium</i> ), Hawthorn ( <i>Crataegus monogyna</i> ), Sycamore ( <i>Acer pseudoplatanus</i> ), Goat Willow ( <i>Salix caprea</i> )	1	Single and Multi	0.1	0.5	0.5	0.5	0.5	0	Middle aged	Fair	Scrubby specimens forming gappy field boundary hedgerow.	No action required at this time.	20-40	C
T40	Oak ( <i>Quercus robur</i> )	12	Multi	0.44	6	6	5	5	3	Middle aged	Good	Prominent multi-stemmed tree of reasonable form and vigour.	No action required at this time.	>40	B

Tree No.	Species	Height(m)	Single/Multi Stemmed	Stem Diameter(m)	Branch Spread(m)				Height of Crown(m)	Age	Physiological Condition	Structural Condition	Prel. Man. Recommendations	Est. Remaining Contribution	Category
					N	E	S	W							
G41	Group of: Hazel ( <i>Corylus avellana</i> ), Ash ( <i>Fraxinus excelsior</i> ), Oak ( <i>Quercus robur</i> ), Holly ( <i>Ilex aquifolium</i> ), Hawthorn ( <i>Crataegus monogyna</i> ), Sycamore ( <i>Acer pseudoplatanus</i> ), Goat Willow ( <i>Salix caprea</i> )	1	Single and Multi	0.1	0.5	0.5	0.5	0.5	0	Middle aged	Fair	Scrubby specimens forming gappy field boundary hedgerow.	No action required at this time.	20-40	C
T42	Oak ( <i>Quercus robur</i> )	19	Single	1.17	9	8	7	9	2	Mature	Good	Prominent boundary tree of good form and well-balanced crown.	No action required at this time.	>40	A
T43	Oak ( <i>Quercus robur</i> )	14	Single	0.43	5	1	6	6	3	Middle aged	Fair	Hedgerow tree of variable form with crown developed on western side only.	No action required at this time.	20-40	B
G44	Group of: Goat Willow ( <i>Salix caprea</i> ), Oak ( <i>Quercus robur</i> )	9	Single and Multi	0.3	1	3	4	2	0	Middle aged	Fair to poor	Scrubby specimens of variable form and vigour suppressed by more dominant central Ash.	Monitor for stability.	10-20	C
T45	Ash ( <i>Fraxinus excelsior</i> )	12	Single	0.36	3	3	3	2	2	Middle aged	Poor	Hedgerow tree severely infected with Ash Dieback Disease.	Remove.	<10	U

Tree No.	Species	Height(m)	Single/Multi Stemmed	Stem Diameter(m)	Branch Spread(m)				Height of Crown(m)	Age	Physiological Condition	Structural Condition	Prel. Man. Recommendations	Est. Remaining Contribution	Category
					N	E	S	W							
G46	Group of: Plum ( <i>Prunus</i> spp.), Hawthorn ( <i>Crataegus monogyna</i> ), Hazel ( <i>Corylus avellana</i> ), Goat Willow ( <i>Salix caprea</i> ), Holly ( <i>Ilex aquifolium</i> ), Blackthorn ( <i>Prunus spinosa</i> )	2	Multi	0.15	1	1	1	1	0	Middle aged	Fair	Scrubby specimens forming dense field boundary hedgerow.	No action required at this time.	>40	C
G47	Group of: Ash ( <i>Fraxinus excelsior</i> )	15	Single and Multi	0.35 (avg.)	6	5	4	5	2	Middle aged	Poor	Hedgerow trees severely infected with Ash Dieback Disease. These specimens are in a moribund condition.	Remove.	<10	U
T48	Oak ( <i>Quercus robur</i> )	19	Single	0.53	7	6	7	9	5	Mature	Good	Prominent boundary tree of good form and well-balanced crown.	No action required at this time.	>40	A
G49	Group of: Ash ( <i>Fraxinus excelsior</i> )	17	Single and Multi	0.3 (avg.)	3	3	2	1	2	Middle aged	Poor	Hedgerow tree severely infected with Ash Dieback Disease. These specimens are in a moribund condition.	Remove.	<10	U

Tree No.	Species	Height(m)	Single/Multi Stemmed	Stem Diameter(m)	Branch Spread(m)				Height of Crown(m)	Age	Physiological Condition	Structural Condition	Prel. Man. Recommendations	Est. Remaining Contribution	Category
					N	E	S	W							
G50	Group of: Goat Willow ( <i>Salix caprea</i> ), Sycamore ( <i>Acer pseudoplatanus</i> ), Hawthorn ( <i>Crataegus monogyna</i> ), Hazel ( <i>Corylus avellana</i> ), Oak ( <i>Quercus robur</i> )	6	Single and Multi	0.2	2	2	2	2	0	Middle aged	Fair	Scrubby specimens forming gappy field boundary hedgerow.	No action required at this time.	20-40	C
T51	Goat Willow ( <i>Salix caprea</i> )	14	Multi	0.6	6	8	4	8	0	Middle aged	Poor	Multi-stemmed hedgerow tree exhibiting significant decay throughout main stem and major limbs. This specimen is at risk of structural failure.	Remove.	<10	U
T52	Oak ( <i>Quercus robur</i> )	17	Single	0.71	7	7	7	9	3	Mature	Good	Prominent boundary tree of good form and upright habit.	No action required at this time.	>40	A
T53	Oak ( <i>Quercus robur</i> )	10	Single	0.45	4	5	3	4	3	Middle aged	Fair	Hedgerow tree of slightly scrubby habit.	No action required at this time.	>40	B
G54	Group of: Ash ( <i>Fraxinus excelsior</i> )	18	Single and Multi	0.5 (avg.)	7	4	6	4	2	Middle aged	Poor	Numerous hedgerow trees severely infected with Ash Dieback Disease.	Remove.	<10	U

Tree No.	Species	Height(m)	Single/Multi Stemmed	Stem Diameter(m)	Branch Spread(m)				Height of Crown(m)	Age	Physiological Condition	Structural Condition	Prel. Man. Recommendations	Est. Remaining Contribution	Category
					N	E	S	W							
G55	Group of: Hazel ( <i>Corylus avellana</i> ), Hawthorn ( <i>Crataegus monogyna</i> ), Blackthorn ( <i>Prunus spinosa</i> ), Goat Willow ( <i>Salix caprea</i> )	4	Single and Multi	0.15	1	1	1	1	0	Middle aged	Fair	Scrubby specimens forming gappy hedgerow.	No action required at this time.	>40	C
T56	Oak ( <i>Quercus robur</i> )	14	Single	0.36	8	2	5	4	3	Middle aged	Fair	Prominent hedgerow tree of variable form with crown more heavily developed on northwestern side. Main stem and mid crown heavily colonised by ivy thus preventing full inspection.	No action required at this time.	>40	B
G57	Group of: Ash ( <i>Fraxinus excelsior</i> )	18	Single and Multi	0.5 (avg.)	7	4	6	4	2	Middle aged	Poor	Numerous hedgerow trees severely infected with Ash Dieback Disease.	Remove.	<10	U
G58	Group of: Hazel ( <i>Corylus avellana</i> ), Hawthorn ( <i>Crataegus monogyna</i> ), Blackthorn ( <i>Prunus spinosa</i> ), Goat Willow ( <i>Salix caprea</i> )	4	Single and Multi	0.15	1	1	1	1	0	Middle aged	Fair	Scrubby specimens forming gappy hedgerow. Some specimens of Goat Willow are partially collapsed.	Remove partially collapsed specimens of Goat Willow.	>40	C

Tree No.	Species	Height(m)	Single/Multi Stemmed	Stem Diameter(m)	Branch Spread(m)				Height of Crown(m)	Age	Physiological Condition	Structural Condition	Prel. Man. Recommendations	Est. Remaining Contribution	Category
					N	E	S	W							
T59	Oak ( <i>Quercus robur</i> )	18	Multi	0.6	7	7	6	4	3	Mature	Good	Prominent twin-stemmed hedgerow tree of reasonable form with crown more heavily developed on northern side.	No action required at this time.	>40	B
G60	Group of: Ash ( <i>Fraxinus excelsior</i> )	18	Single and Multi	0.5 (avg.)	7	4	6	4	2	Middle aged	Poor	Several hedgerow trees severely infected with Ash Dieback Disease.	Remove.	<10	U
T61	Oak ( <i>Quercus robur</i> )	15	Single	0.47	5	1	6	5	3	Middle aged	Fair	Prominent hedgerow tree of variable form suppressed on eastern side by more dominant Ash.	No action required at this time.	>40	B
G62	Group of 3: Oak ( <i>Quercus robur</i> )	18	Single and Multi	0.55 (avg.)	7	7	9	8	2	Mature	Fair	Prominent hedgerow group containing trees of good upright habit. Some evidence of slight thinning of crowns.	Monitor for health.	>40	B
T63	Ash ( <i>Fraxinus excelsior</i> )	15	Single	0.32	8	8	0	0	3	Middle aged	Poor	Hedgerow tree leaning excessively to the northeast. This specimen is severely infected with Ash Dieback Disease.	Remove.	<10	U
T64	Oak ( <i>Quercus robur</i> )	18	Multi	1.0	9	9	8	6	2	Mature	Good	Prominent multi-stemmed hedgerow tree of good form and vigour.	No action required at this time.	>40	A
T65	Oak ( <i>Quercus robur</i> )	17	Single	0.26	2	3	3	2	8	Middle aged	Fair	Hedgerow tree with low stem diameter to height ratio.	Monitor for stability.	>40	B
T66	Dead														
T67	Oak ( <i>Quercus robur</i> )	13	Single	0.31	3	7	2	1	2	Middle aged	Fair to poor	Heavily suppressed hedgerow tree of variable form and declining vigour.	Monitor for health.	10-20	C



Tree No.	Species	Height(m)	Single/Multi Stemmed	Stem Diameter(m)	Branch Spread(m)				Height of Crown(m)	Age	Physiological Condition	Structural Condition	Prel. Man. Recommendations	Est. Remaining Contribution	Category
					N	E	S	W							
T68	Oak ( <i>Quercus robur</i> )	19	Single	0.41	8	7	4	7	4	Mature	Fair	Prominent hedgerow tree of good upright habit. Evidence of slight thinning of foliage within crown.	Monitor for health.	>40	B
T69	Oak ( <i>Quercus robur</i> )	15	Single	0.41	1	3	9	3	2	Middle aged	Fair	Hedgerow tree leaning excessively to the south.	Monitor for stability.	20-40	B
G70	Group of: Ash ( <i>Fraxinus excelsior</i> )	18	Single and Multi	0.5 (avg.)	7	4	6	4	2	Middle aged	Poor	Several hedgerow trees severely infected with Ash Dieback Disease.	Remove.	<10	U
G71	Group of: Hazel ( <i>Corylus avellana</i> ), Hawthorn ( <i>Crataegus monogyna</i> ), Blackthorn ( <i>Prunus spinosa</i> ), Goat Willow ( <i>Salix caprea</i> ), Elder ( <i>Sambucus nigra</i> ), Holly ( <i>Ilex aquifolium</i> )	4	Single and Multi	0.15	1	1	1	1	0	Middle aged	Fair	Scrubby specimens forming gappy hedgerow.	No action required at this time.	>40	C
T72	Oak ( <i>Quercus robur</i> )	11	Single	0.28	0	9	7	0	2	Middle aged	Poor	Heavily suppressed specimen of poor form leaning excessively to the southeast. This specimen is vulnerable to structural failure and is unsuitable for retention.	Remove.	<10	U
T73	Oak ( <i>Quercus robur</i> )	17	Single	0.44	4	4	3	3	1	Middle aged	Fair	Prominent hedgerow tree of slightly variable form.	No action required at this time.	>40	B

Tree No.	Species	Height(m)	Single/Multi Stemmed	Stem Diameter(m)	Branch Spread(m)				Height of Crown(m)	Age	Physiological Condition	Structural Condition	Prel. Man. Recommendations	Est. Remaining Contribution	Category
					N	E	S	W							
T74	Oak ( <i>Quercus robur</i> )	17	Single	0.89	12	5	4	12	4	Mature	Fair	Offsite tree of variable form with crown developed on northern and western side only. Evidence of some thinning and dieback in upper crown. Some storm damaged branches hung up in lower crown.	Prune to remove hung up storm damaged branches. Monitor for health.	>40	B
T75	Ash ( <i>Fraxinus excelsior</i> )	11	Multi	0.35	1	1	1	1	2	Middle aged	Poor	This specimen is severely infected with Ash Dieback Disease and is now in a moribund condition.	Remove.	<10	U
T76	Oak ( <i>Quercus robur</i> )	11	Single	0.74	7	6	7	7	4	Mature	Good	Prominent boundary tree of good form and well-balanced crown. Main stem colonised by ivy thus preventing full inspection.	No action required at this time.	>40	A

Tree No.	Species	Height(m)	Single/Multi Stemmed	Stem Diameter(m)	Branch Spread(m)				Height of Crown(m)	Age	Physiological Condition	Structural Condition	Prel. Man. Recommendations	Est. Remaining Contribution	Category
					N	E	S	W							
G77	Group of: Hazel ( <i>Corylus avellana</i> ), Blackthorn ( <i>Prunus spinosa</i> ), Goat Willow ( <i>Salix caprea</i> ), Hawthorn ( <i>Crataegus monogyna</i> ), Sycamore ( <i>Acer pseudoplatanus</i> ), Ash ( <i>Fraxinus excelsior</i> ), Oak ( <i>Quercus robur</i> ), Holly ( <i>Ilex aquifolium</i> ), Elm ( <i>Ulmus</i> spp.)	9	Single and Multi	0.25	2	2	2	2	0	Middle aged	Fair	Scrubby specimens forming gappy field boundary hedgerow containing some diseased Ash.	Remove diseased Ash trees.	>40	C
T78	Oak ( <i>Quercus robur</i> )	17	Multi	0.8	8	10	7	10	3	Mature	Fair	Prominent boundary tree of reasonable form and vigour. Some storm damaged branches in lower crown. Evidence of slight thinning of foliage within crown.	Prune to remove storm damaged branches. Monitor for health.	>40	B
T79	Oak ( <i>Quercus robur</i> )	17	Multi	0.7	4	7	6	7	3	Middle aged	Fair	Twin-stemmed specimen of reasonable form.	No action required at this time.	>40	B
T80	Goat Willow ( <i>Salix caprea</i> )	4	Multi	0.25	0	0	0	6	0	Middle aged	Poor	This specimen has collapsed.	Remove.	<10	U
T81	Ash ( <i>Fraxinus excelsior</i> )	17	Multi	0.5	4	5	4	3	2	Middle aged	Poor	This specimen is infected with Ash Dieback Disease.	Remove.	<10	U

Tree No.	Species	Height(m)	Single/Multi Stemmed	Stem Diameter(m)	Branch Spread(m)				Height of Crown(m)	Age	Physiological Condition	Structural Condition	Prel. Man. Recommendations	Est. Remaining Contribution	Category
					N	E	S	W							
T82	Elm ( <i>Ulmus</i> spp.)	3	Multi	0.2	4	0	1	2	0	Middle aged	Poor	Scrubby heavily suppressed specimen leaning to the north that is unsuitable for retention.	Remove.	<10	U
G83	Group of: Oak ( <i>Quercus robur</i> ), Holly ( <i>Ilex aquifolium</i> ), Sycamore ( <i>Acer pseudoplatanus</i> ), Elm ( <i>Ulmus</i> spp.)	Up to 16	Single and Multi	0.25 (avg.)	3	4	3	5	2	Middle aged	Fair	Trees of generally variable form suppressed by more dominant trees to the east.	No action required at this time.	20-40	C
T84	Oak ( <i>Quercus robur</i> )	20	Single	0.72	7	9	9	9	4	Mature	Fair	Prominent boundary tree of reasonable form and vigour. Evidence of historical splitting on major limb in central crown. Evidence of slight thinning of foliage in upper crown.	Monitor for safety.	>40	B
G85	Group of 3: Oak ( <i>Quercus robur</i> )	18	Single	0.3	4	4	3	4	3	Middle aged	Fair	Boundary trees of reasonable form.	No action required at this time.	20-40	B
G86	Group of 3: Ash ( <i>Fraxinus excelsior</i> )	17	Single	0.25	4	4	3	5	4	Middle aged	Poor	These specimens are infected with Ash Dieback Disease.	Remove.	<10	U
T87	Sycamore ( <i>Acer pseudoplatanus</i> )	18	Multi	0.6	5	6	5	5	2	Middle aged	Good	Prominent twin-stemmed specimen of good form and vigour.	No action required at this time.	>40	B
T88	Oak ( <i>Quercus robur</i> )	18	Multi	0.65	1	8	9	6	3	Mature	Good	Prominent twin-stemmed specimen of reasonable form with crown more heavily developed on southern side.	No action required at this time.	>40	B

Tree No.	Species	Height(m)	Single/Multi Stemmed	Stem Diameter(m)	Branch Spread(m)				Height of Crown(m)	Age	Physiological Condition	Structural Condition	Prel. Man. Recommendations	Est. Remaining Contribution	Category
					N	E	S	W							
G89	Group of: Cherry Laurel ( <i>Prunus laurocerasus</i> )	2	Multi	0.1	1	1	1	1	0	Middle aged	Fair	Evergreen boundary hedgerow.	No action required at this time.	10-20	C