

# NutriZMC Flo

NPKS 3-0-0-0 +  
Manganese, Zinc & Copper

10 LITRE



## WARNING

H302+H312+H332 Harmful if swallowed, in contact with skin, or if inhaled.  
H320 Causes eye irritation. P234 Keep only in original container. P264  
Wash exposed skin thoroughly after handling. P280 Wear protective gloves/  
protective clothing/eye protection/face protection. P273 Avoid release to the environment.  
P301+330+331 IF SWALLOWED: Rinse mouth. Do Not induce vomiting.



# NutriZMC Flo

ANALYSIS	AUSTRALIA (W/V%)	INTERNATIONAL (W/W%)
Nitrogen (N)	3.0	1.6
Zinc (Zn)	25.0	13.2
Manganese (Mn)	30.0	15.8
Copper (Cu)	10.0	5.3

SG: 1.89 - 1.91

## DIRECTIONS FOR USE:

Agitate contents well before dilution.

Suitable for application by:



## Benefits of NutriZMC Flo

- Accelerated early vegetative growth
- Promotion of root systems to access nutrients from soils
- Controlled particle morphology to delivering sustained uptake over 4 - 6 weeks

**COMPATIBILITY: NutriZMC Flo** is compatible with a wide range of commonly used spray chemical. When mixing with other chemicals always mix a small quantity and check for precipitation (fall out) in mixture. If precipitation occurs, DO NOT proceed with application. If mixture does not precipitate and it is the first time you have used this combination of products, test a small area of the crop prior to treatment of the whole crop.

**STORAGE AND HANDLING:** Store in a cool, well-ventilated area, out of direct sunlight, avoiding temperature extremes. Agitate this container prior to use and replace lid after use.

**CONDITIONS OF SALE:** To the extent permitted by law, Agrichem its distributors and agents, liability (including negligence) for any direct or indirect loss arising in any way from use of the product is limited to replacing the product.

**NOTE:** The suggested rates of application are designed for typical growing conditions and should be used as a guide only. Each farmer's climatic conditions, water quality, soil types, application processes and practices may differ and therefore necessitate corrections to ensure optimum results. Good agricultural practice requires that application be avoided under extreme weather conditions such as temperatures over 28°C, high humidity, frost, rain etc. It is recommended that when applying to a crop or area for the first time, or in combination with other chemicals, a small test area should be sprayed and observed prior to the total spray. Where possible, it is recommended that regular leaf (sap) tests are conducted to determine actual plant nutrient availability during each growth cycle. Soil tests at least once per year are essential.

**GENERAL PRECAUTIONS AND SAFETY:** Do not swallow. Avoid skin and eye contact, ingestion and contact with or inhalation of spray mist. When handling this product, wear suitable gloves, goggles, hat and protective clothing. Should the product be ingested or symptoms or signs develop after inhaling or coming into contact with the product, contact a doctor or Poisons Information Centre 13 11 26.



# agrichem®

CROP	Foliar L/ha	Fertigation L/ha	COMMENTS
<b>BROADACRE:</b> Such as Barley, Canola, Cotton, Grain legumes, Maize, Oats, Rice, Sorghum, Triticale, Wheat & Pasture crops	0.3 - 1.5 in a minimum of 15 - 60 L final spray volume.*	0.5 - 1	Best applied at 3-4 true leaf, may be used at other growth stages. For maintenance, use the higher rate. Best applied at late cabbage stage, may be used at other stages in Canola. For Cotton, apply to boost growth early in season & post waterlogging events. *Aerial application: use maximum practicable water rates. For cotton and pastures, increase dilution rates to 1:60.
<b>DECIDUOUS TREE CROPS:</b> Such as Apple, Almond, Cherry, Nectarine, Peach, Pear, Pistachio and Walnut	1-3 in a minimum of 100 - 300L final spray volume.	2-3	Spray at early bud, post petal fall. Apply post harvest at higher rates of 3L/ha. Note: Avoid applications during flowering. For Stone Fruit: Dormancy spray only. Do not apply to foliage or fruit.
<b>EVERGREEN TREE CROPS:</b> Such as Avocado, Citrus, Macadamia, Lychee	1-3 in a minimum of 100 - 300L final spray volume.	2-3	Apply to recently hardened spring flush or during active growing period & post harvest.
<b>FRUITING VEGETABLES:</b> Such as Capsicum, Cucurbits, Eggplant, Tomatoes, Pumpkins	1-3 in a minimum of 100 - 300L final spray volume.	2-3	Apply at regular intervals from 5th leaf until 21 days pre harvest. Best applied pre flowering. Apply with compatible crop protection sprays.
<b>LEAFY VEGETABLES:</b> Such as Endive, Fennel Lettuce, Broccoli, Cabbage, Cauliflower, Kale and Herbs	1-3 in a minimum of 100 - 300L final spray volume.	2-3	Best applied at 3-4 true leaf, may be used at other growth stages. For maintenance, use the higher rate. Apply with compatible crop protection sprays.
<b>ROOT VEGETABLES:</b> Such as Beetroot, Carrot, Leek, Onion, Potato, Radish, Sweet Potato	1-3 in a minimum of 25 - 300L final spray volume.	2-3	Apply regularly, may substitute for urea or sulphate of ammonia. Can be used in organic production to supply N. Apply with compatible crop protection sprays.
<b>VINE AND BERRY CROPS:</b> Such as Blueberry, Strawberry, Raspberry, Wine and Table Grapes	2-4 in a minimum of 500 - 1000L final spray volume.	2-3	Apply 1-2 applications prior to flowering. Do not exceed 2x concentration or 2x the per hectare rate. Use the higher fertigation rate post harvest. Note: foliar spray, do not use at concentrations less than 1:50.

Fertigation rates are dependent on seasonal nutrient demand. Agitate contents well prior to application.

## SEED DRESSING

CROP	Rate L/Tonne	MIN DILUTION	COMMENTS
<b>BROADACRE</b> Barley, Cotton, Oats, Triticale, Wheat	4-6	Mix sufficient water to ensure adequate coating of seed	If using lower rates follow up tissue tests may be required to determine the need for foliar applications post emergence. If applied without dilution uneven seed coverage may occur. Apply 1-3 L water tonne seed, depending on seed moisture content & ambient air temperature. Some of the water may be substituted with Kelpak (a plant growth promoter) at 1-2 L/tonne of seed, with the upper rate being applied to small seeds such as Canola. Kelpak assists the germination process & encourages root development. Nutri ZMC Flo is not compatible with rhizobia inoculant products.
Canola	20		
Grain legumes	4-6		
Maize, Rice	5-8		
Potato Seed piece	30	1:66	Apply in minimum of 200L water/ha with compatible plant protection products and seed piece spray.

⚠️ MINIMUM DILUTION: A dilution of 1 : 100 means 1 part product : 100 parts water. In hot weather use a higher dilution rate. AERIAL APPLICATION: Use maximum practical water rates

Made in Australia by agrichem

2 Hovey Road, Yatala, QLD 4207 Australia.

Ph: +61 7 3451 0000 Fx: +61 7 3451 0093 Free Call: 1800 654 758

www.agrichem.com.au

