

NutriZinc Flo

65% Zinc
(1.8-0-0-0)

10 LITRE



NutriZinc Flo

ANALYSIS	AUSTRALIA (W/V%)	INTERNATIONAL (W/W%)
Nitrogen (N)	1.8	1
Zinc (Zn)	65	38.4
SG: 1.70		

DIRECTIONS FOR USE:

Agitate contents well before dilution.

Suitable for application by:



Foliar



Fertigation

COMPATIBILITY: NutriZinc Flo is compatible with a wide range of commonly used spray chemicals. When mixing with other chemicals, always mix a small quantity and check for precipitate (fall out) in mixture. Precipitation occurs. DO NOT proceed with application. If mixture does not precipitate and it's the first time you have used this combination of products, test a small area of the crop prior to treatment of the whole crop.

STORAGE AND HANDLING: Store in a cool, well-ventilated area, out of direct sunlight, avoiding temperature extremes. Agitate this container prior to use and replace lid after use.

CONDITIONS OF SALE: To the extent permitted by law, Agrichem's distributors and agents, liability (including negligence) for any direct or indirect loss arising in any way from use of the product is limited to replacing the product.

NOTE: The suggested rates of application are designed for typical growing conditions and should be used as a guide only. Each farmer's climatic conditions, water quality, soil types, application processes and practices may differ and therefore necessitate corrections to ensure optimum results. Good agricultural practice requires that application be avoided under extreme weather conditions such as temperatures over 28°C, high humidity, frost, rain etc. It is recommended that when applying to a crop or area for the first time, or in combination with other chemicals, a small test area should be sprayed and observed prior to the total spray. Where possible, it is recommended that regular leaf (sap) tests are conducted to determine actual plant nutrient availability during each growth cycle. Soil tests at least once per year are essential.

GENERAL PRECAUTIONS AND SAFETY: Do not swallow. Avoid skin and eye contact, ingestion and contact with or inhalation of spraymist. When handling this product, wear suitable gloves, goggles, hat and protective clothing. Should the product be ingested or symptoms or signs develop after inhaling or coming into contact with the product, contact a doctor or Poisons Information Centre 13 11 26.

NOT CLASSIFIED AS HAZARDOUS ACCORDING TO THE GLOBALLY HARMONISED SYSTEM OF CLASSIFICATION AND LABELLING OF CHEMICALS (GHS) AND SAFE WORK AUSTRALIA CRITERIA.



agrichem®




CROP	Rate/ha	MIN DILUTION	COMMENTS
AVOCADOS, CITRUS, NUTS, OLIVES & TROPICAL FRUITS	0.25 - 1 L	1:50	Apply to recently hardened spring flush or during active growing period and post harvest.
BANANAS	0.25 - 2 L	1:75	Apply from early growth, at budding or as required.
BROADACRE - Foliar Barley, Canola, Cotton, Legumes, Maize, Oats, Rice, Sorghum, Triticale & Wheat	0.1 - 0.2 L 0.2 - 0.6 L 0.6 - 1.5 L	1:20	Maintenance rate Moderate deficiency Severe deficiency NutriZinc Flo is best applied at the 3 - 4 leaf stage, but can be used at other stages.
- Down the tube with UAN	0.1 - 2 L	1:5	Apply at seeding
- Water Injection	0.25 - 0.5 L	1:20	Zinc rapidly remediates the effects of SU and other herbicides on the crop. Where compatible, NutriZinc Flo makes the combination product safer and also aids the rapid recovery of crops adversely affected by herbicides.
- Herbicide Remediator	0.15 L		
HORTICULTURE/VEGETABLES - Foliar - Fertigation	0.25 - 1 L 0.5 - 1.5 L	1:50	Apply at early stages of growing season when sufficient leaf cover present. Apply with compatible crop protection sprays.
PINEAPPLES	0.25 - 0.75 L	1:50	Apply at 4 - 6 week intervals with compatible protection sprays.
POME/STONE FRUIT	0.25 - 0.75 L	1:75	First application: early bud. Second application: after petal fall. Top up with the higher rate post harvest.
STRAWBERRIES	0.25 - 0.5 L	1:75	Apply at 4 leaf stage and during early stages of growth.
VINES - Table grapes - Wine grapes	0.25 - 0.75 L	1:75	First applications: shoots 10cm. Second application: 5% flowering. Do not use at concentrations less than 1:10. As much as 4L/ha can be used without detriment to the crop.

SEED DRESSING

CROP	Rate/t	MIN DILUTION	COMMENTS
BROADACRE Barley, Cotton, Oats, Triticale, Wheat	2.2 - 4.4		Mix with sufficient water or compatible liquid plant protection products to ensure adequate coating of seed. If using lower rates, follow up tissue tests may be required to determine the need for foliar applications post emergence. If applied without dilution, uneven seed coverage may occur. Apply 1 - 3 L water/tonne seed, depending on seed moisture content & ambient air temperature. Some of the water may be substituted with Kelpak (a plant growth promoter) at 1 - 2 L/tonne of seed, with the upper rate being applied to small seeds such as Canola. Kelpak assists germination process & encourages root development. Nutri Zinc is not compatible with rhizobia inoculant products.
CANOLA	10 - 15 L		
GRAIN LEGUMES	2.8 - 5.5 L		
MAIZE, RICE	2.8 - 6.5		
POTATO SEED PIECE	2.2 - 5.5 L/ha	1:20	Apply in minimum of 200L water/ha with compatible plant protection product, seed piece spray

④ - MINIMUM DILUTION: A dilution of 1:100 means 1 part product : 100 parts water. In hot weather use a higher dilution rate.

Made in Australia by  **agrichem®**

2 Hovey Road, Yatala, QLD 4207 Australia.

Ph: +61 7 3451 0000 Fx: +61 7 3451 0093 Free Call: 1800 654 758

www.agrichem.com.au

